

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 953 639 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

03.11.1999 Bulletin 1999/44

(21) Application number: 98107925.4

(22) Date of filing: 30.04.1998

(51) Int. Cl.⁶: **C12N 15/13**, C07K 16/40,
C07K 16/46, C12N 15/62,
C12N 15/85, C12N 5/10,
C07K 19/00, A61K 47/48,
A61K 51/10, A61K 39/395,
G01N 33/577, G01N 33/574

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(71) Applicant:

Boehringer Ingelheim International GmbH
55216 Ingelheim (DE)

(72) Inventors:

• Park, John Edward
88400 Biberach/Riss (DE)

- Garin-Chesa, Pilar
88400 Biberach/Riss (DE)
- Bamberger, Uwe
88416 Ochsenhausen (DE)
- Leger, Olivier
74100 Annemasse (FR)
- Saldanha, Jose
Enfield Middlesex, EN1 1TE (GB)
- Rettig, Wolfgang J.
88400 Biberach a.d. Riss (DE)

Remarks:

The applicant has subsequently filed a sequence
listing and declared, that it includes no new matter.

(54) **FAPalpha-specific antibody with improved producibility**

(57) Recombinant antibody proteins are provided
that specifically bind fibroblast activation protein alpha
(FAP α) and comprise framework modifications resulting
in the improved producibility in host cells. The invention
also relates to the use of said antibodies for diagnostic
and therapeutic purposes and methods of producing
said antibodies.

EP 0 953 639 A1

BEST AVAILABLE COPY

ATTORNEY DOCKET NUMBER: 11183-003-999
SERIAL NUMBER: 10/524,134
REFERENCE: **B12**

Description**Field of the Invention**

5 [0001] The present invention relates to antibody proteins that specifically bind fibroblast activation protein alpha (FAP α). The invention also relates to the use of said antibodies for diagnostic and therapeutic purposes and methods of producing said antibodies.

Background of the Invention

10 [0002] The invasive growth of epithelial cancers is associated with a number of characteristic cellular and molecular changes in the supporting stroma. A highly consistent molecular trait of the reactive stroma of many types of epithelial cancer is induction of the fibroblast activation protein alpha (from now on referred to as FAP), a cell surface molecule of reactive stromal fibroblasts originally identified with monoclonal antibody F19 (Garin-Chesa P., Old L. J. and Rettig W. J. (1990) Cell surface glycoprotein of reactive stromal fibroblasts as a potential antibody target in human epithelial cancers. *Proc. Natl. Acad. Sci.* 87: 7235). Since the FAP antigen is selectively expressed in the stroma of a range of epithelial carcinomas, independent of location and histological type, a FAP-targeting concept has been developed for imaging, diagnosis and treatment of epithelial cancers and certain other conditions. For this purpose a monoclonal antibody termed F19 that specifically binds to FAP was developed and described in US Patent 5,059,523, which is hereby
15 incorporated by reference in its entirety.

20 [0003] One serious problem that arises when using non-human antibodies for applications *in vivo* in humans is that they quickly raise a human anti-non-human response which reduces the efficacy of the antibody in patients and impairs continued administration. Humanisation of non-human antibodies is commonly achieved in one of two ways: (1) by constructing non-human/human chimeric antibodies, wherein the non-human variable regions are joined to human constant regions (Boulianne G. L., Hozumi N. and Shulman, M. J. (1984) Production of functional chimaeric mouse/human antibody *Nature* 312: 643) or (2) by grafting the complementarity determining regions (CDRs) from the non-human variable regions to human variable regions and then joining these "reshaped human" variable regions to human constant regions (Riechmann L., Clark M., Waldmann H. and Winter G. (1988) Reshaping human antibodies for therapy. *Nature* 332: 323). Chimeric antibodies, although significantly better than mouse antibodies, can still elicit an anti-mouse response in humans (LoBuglio A. F., Wheeler R. H., Trang J., Haynes A., Rogers K., Harvey E. B., Sun L., Ghayeb J. and Khazaeli M. B. (1989) Mouse/human chimeric monoclonal antibody in man: Kinetics and immune response. *Proc. Natl. Acad. Sci.* 86: 4220). CDR-grafted or reshaped human antibodies contain little or no protein sequences that can be identified as being derived from mouse antibodies. Although an antibody humanised by CDR-grafting may still be able to elicit some immune reactions, such as an anti-allotype or an anti-idiotypic response, as seen even with natural
25 human antibodies, the CDR-grafted antibody will be significantly less immunogenic than a mouse antibody thus enabling a more prolonged treatment of patients.

30 [0004] Another serious limitation relating to the commercial use of antibodies for diagnosis, imaging and therapy is their producibility in large amounts. In many instances recombinant expression of native, chimeric and/or CDR-grafted antibodies in cell culture systems is poor. Factors contributing to poor producibility may include the choice of leader sequences and the choice of host cells for production as well as improper folding and reduced secretion. Improper folding can lead to poor assembly of heavy and light chains or a transport incompetent conformation that forbids secretion of one or both chains. It is generally accepted, that the L-chain confers the ability of secretion of the assembled protein. In some instances multiple or even single substitutions can result in the increased producibility of antibodies.

35 [0005] Because of the clinical importance of specific immunological targeting *in vitro* and *in vivo* of specific disease-related antigens for diagnosis and therapy in humans, there is a growing need for antibodies that combine the features of antigen specificity, low immunogenicity and high producibility.

[0006] Therefore, the problem underlying the present invention was to provide antibody proteins that combine the properties of specific binding to FAP, low immunogenicity in humans, and high producibility in recombinant systems.

Disclosure of the invention

50 [0007] The technical problem is solved by the embodiments characterized in the claims.

[0008] The present invention provides new antibody proteins having the complementary determining regions of the monoclonal antibody F19 (ATCC Accession No. HB 8269), said new antibody proteins specifically binding to fibroblast activation protein (FAP), characterised in that they have framework modifications resulting in the improved producibility in host cells as compared to a chimeric antibody having the variable regions of F19 and foreign constant regions.

[0009] As used herein, an "antibody protein" is a protein with the antigen binding specificity of a monoclonal antibody.

[0010] "Complementarity determining regions of a monoclonal antibody" are understood to be those amino acid

sequences involved in specific antigen binding according to Kabat (Kabat E. A., Wu T. T., Perry H. M., Gottesman K. S. and Foeller C. (1991) *Sequences of Proteins of Immunological Interest* (5th Edn). NIH Publication No. 91-3242. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Bethesda, MD.) in connection with Chothia and Lesk (Chothia and Lesk, J. Mol. Biol., 196:901-917 (1987)).

5 [0011] As used herein, the term "framework modifications" refers to the exchange, deletion or addition of single or multiple amino acids in the variable regions surrounding the individual complementarity determining regions. Framework modifications may have an impact on the immunogenicity, producibility or binding specificity of an antibody protein.

[0012] "Fibroblast activation protein (FAP)", also designated fibroblast activation protein alpha (FAP α), is a membrane-bound glycoprotein belonging to the serine protease gene family (WO 97/34927). No shed or secreted form of FAP is known.

[0013] FAP can be characterized by its binding to the monoclonal antibody F19 (F19 is obtainable from the hybridoma cell line with the accession No. HB 8269 deposited at the ATCC).

[0014] The term "fibroblast activation protein specific binding" of an antibody protein is defined herein by its ability to specifically recognise and stably bind FAP-expressing human cells. The binding specificity of the proteins of the invention can be determined by standard methods for the evaluation of binding specificity such as described in an exemplary fashion in example 6, 8 and example 12.

[0015] The term "chimeric antibody" refers to an antibody protein having the light and heavy chain variable regions as described in figures 17 and 18 and foreign constant regions. "Foreign constant regions" as defined herein are constant regions which are different from the constant regions of F19. For comparing an antibody protein of the invention to a chimeric antibody it is to be understood that such a chimeric antibody must contain the same constant regions as said antibody protein. For the purpose of demonstration and comparison alone the human constant heavy and light chains as described in Figures 19 to 22 are used in an exemplary fashion.

[0016] To provide the antibody proteins of the present invention, the nucleic acid sequences of the heavy and light chain genes of the murine antibody designated F19 were determined from RNA extracted from F19 hybridoma cells (ATCC Accession No. HB 8269).

[0017] In one embodiment the present invention relates to antibody proteins having the complementary determining regions of the monoclonal antibody F19 (ATCC Accession No. HB 8269), said new antibody proteins specifically binding to fibroblast activation protein (FAP), characterized in that they have framework modifications resulting in the improved producibility in host cells as compared to a chimeric antibody having the variable regions of F19 and foreign constant regions, wherein said antibody protein is derived from the murine antibody designated F19 (ATCC Accession No. HB 8269).

[0018] To generate humanised FAP-specific antibody proteins a chimeric antibody was constructed, having variable regions of the light and heavy chains of F19 and human light and heavy constant regions, respectively. The construction and production of chimeric mouse/human antibodies is well known (Boulianne et al. (1984), referenced above) and demonstrated in an exemplary fashion in examples 1 and 2.

[0019] Therefore, in a further embodiment the invention relates to antibody proteins according to the invention, characterised in that they have a variable light chain region and a variable heavy chain region, each joined to a human constant region.

[0020] In particular, the variable region of the light chain was joined to a human kappa constant region and the variable region of the heavy chain was joined to a human gamma-1 constant region. Other human constant regions for humanising light and heavy chains are also available to the expert. A human kappa and a human gamma-1 constant regions were used for demonstrating the invention in an exemplary fashion only.

[0021] Therefore, in one particular embodiment the antibody proteins of the invention contain a human kappa constant region.

[0022] Also, in another particular embodiment the antibody proteins of the invention contain a human gamma-1 constant region.

[0023] One particular "chimeric F19 antibody" protein (cF19) consists of the light and heavy chain variable and constant regions described in Figures 17 to 22. cF19 demonstrates specific binding and high avidity to the FAP antigen. As demonstrated in example 2, the expression of cF19 in COS cells is poor, ranging from about 10 to 60 ng/ml, which is at least 10 fold less than most antibodies.

[0024] In an attempt to increase expression levels of cF19, the leader sequence of the F19 V_L region was changed by substitution of Proline to Leucine at position -9.

[0025] This single change in amino acid in the leader sequence resulted in at least doubling the amount of chimeric antibody produced in COS cells. For the expression of this particular chimeric antibody in COS cells the following mutated leader sequence of the light chain: MDSQAQVLMLLLWVSGTCG, and the following leader sequence of the heavy chain: MGWSWVFLFLLSGTAGVLS were used.

[0026] According to the invention the term "improved producibility" in host cells refers to the substantial improvement of expression levels and/or purified antibody yields when compared with the expression levels and/or antibody yields of

a chimeric antibody without framework modifications as defined above. Two particular but not limiting examples for demonstrating improved producibility are exemplified for the COS cell expression system (in examples 2 and 5) and for the CHO cell expression system (in example 10 and 11).

[0027] While the mutation of the leader sequence only lead to the doubling of the expression yield of the chimeric F19 antibody, a substantial improvement as defined herein refers to an improvement in expression level and/or purification yield of at least a factor of 10.

[0028] In a preferred embodiment, the invention refers to antibody proteins, characterised in that their expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 10.

[0029] In more preferred embodiment, the invention refers to antibody proteins, characterised in that their expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 20.

[0030] In a most preferred embodiment, antibody proteins, characterised in that their expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 100.

[0031] Improved producibility of the recombinant antibody proteins of the invention can be demonstrated for eucaryotic cells in general as shown for COS (cells derived from the kidney of an African green monkey) and CHO (Chinese hamster ovary derived cells) eucaryotic cells (see examples 5 and 11). In a further embodiment, the present invention relates to recombinant antibody proteins characterised in that they display improved producibility in eucaryotic cells.

[0032] In a preferred embodiment the present invention relates to antibody proteins, wherein said eucaryotic cell is a chinese hamster ovary cell (CHO cell).

[0033] It was unexpectedly found that certain framework modifications of the light chain variable regions determine the improved producibility of the antibody proteins of the invention. Three versions of reshaped light chain variable regions, designated version A, B, and C, as described in Figures 1 to 6, were prepared.

[0034] Light chain variable region versions A, B, and C demonstrate substantially improved producibility in CHO cells (see example 11). While light chain variable region versions A and C differ from light chain variable region version B by only two common amino acid residues they display an even further substantial improvement in producibility. There is at least another 10 fold difference in antibody secretion levels between the human reshaped F19 light chain version B and versions A or C. Reshaped human F19 light chain version A and B only differ in their amino acid sequences by two residues at positions 36 (Tyr to Phe mutation) and 87 (Tyr to Asp mutation) (nomenclature according to Kabat). This negative effect on the secretory capability of antibodies containing the light chain variable region version B could have been indirect if the Tyr to Asp and Tyr to Phe mutations, considered individually or together, merely caused improper folding of the protein. But this is unlikely to be the case since antigen binding assays show that immunoglobulins containing F19 light chain version B have similar avidities to those paired with F19 light chain version A or C, suggesting that they were not grossly misfolded.

[0035] Residue 87 in reshaped human F19 light chain version B seems particularly responsible for the reduction of secretion when compared to versions A and C.

[0036] In a preferred embodiment, the present invention relates to antibody proteins according to the invention, wherein the amino acid in Kabat position 87 of the light chain region is not asparagine.

[0037] In a more preferred embodiment, the invention relates to antibody proteins according to the invention, wherein the amino acid in Kabat position 87 of the light chain region is selected from aromatic or aliphatic amino acids.

[0038] In a most preferred embodiment, the present invention relates to antibody proteins according to the invention, wherein the aromatic amino acid in Kabat position 87 of the light chain region is a tyrosine or phenylalanine.

[0039] In a further embodiment, the present invention also pertains to antibody proteins according to the invention, wherein the amino acid in Kabat position 36 of the light chain region is selected from aromatic amino acids.

[0040] In a particular embodiment the invention relates to the specific antibody proteins that may be prepared from the individually disclosed reshaped variable regions of the light and heavy chains.

[0041] Especially light chain variable region versions A and C are particularly suitable to practice the invention because of their exceptionally high producibility, while retaining full FAP-binding specificity and achieving low immunogenicity. This holds especially true when compared to the chimeric antibody having the variable regions of F19 and the same constant regions but also when compared to light chain version B.

[0042] Therefore, in one embodiment the present invention relates to antibody proteins that contain the variable region of the light chain as set forth in SEQ ID NO: 2. In a further embodiment the invention also relates to antibody proteins, characterised in that the variable region of the light chain is encoded by a nucleotide sequence as set forth in

SEQ ID NO: 1.

[0043] In one embodiment the present invention relates to antibody proteins that contain the variable region of the light chain as set forth in SEQ ID NO: 6.

[0044] In a further embodiment the invention also relates to antibody proteins characterised in that the variable region

of the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 5.

[0045] The present invention also discloses several different variable regions of the heavy chain that work particularly well with the variable regions of the light chain versions A and C in terms of improved producibility.

[0046] In one embodiment the invention relates to antibody proteins containing a variable region of the heavy chain as set forth in any one of SEQ ID NOs: 8, 10, 12, 14.

[0047] In another embodiment the invention relates to antibody proteins characterised in that the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in any one of SEQ ID NOs: 7, 9, 11, 13.

[0048] In a very particular embodiment the invention relates to antibody proteins containing the variable region of the light chain as set forth in SEQ ID NO: 2 and the variable region of the heavy chain as set forth in SEQ ID NOs: 12.

[0049] In a further particular embodiment the invention relates to antibody proteins characterised in that the variable region of the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 1 and the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 11.

[0050] In a further particular embodiment the invention relates to antibody proteins containing the variable region of the light chain as set forth in SEQ ID NO: 2 and the variable region of the heavy chain as set forth in SEQ ID NOs: 8.

[0051] In a further particular embodiment the invention relates to antibody proteins characterised in that the variable region of the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 1 and the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 7.

[0052] In a further aspect, the present invention relates to nucleic acid molecules containing the coding information for the antibody proteins according to the invention as disclosed above. Preferably, a nucleic acid molecule according to the present invention is a nucleic acid molecule containing a nucleotide sequence selected from SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, or 15.

[0053] A further aspect of the present invention is a recombinant DNA vector containing the nucleotide sequence of any one of the above-mentioned nucleic acids, especially when said nucleotide sequence is operationally linked to an expression control sequence as in expression vectors. Preferred is a recombinant DNA vector, said vector being an expression vector.

[0054] A further aspect of the present invention is a host cell carrying a vector as described, especially an expression vector. Such a host cell can be a procaryotic or eucaryotic cell. Preferably, such a host cell is a eucaryotic cell, a yeast cell, or a mammalian cell. More preferably, said host cell is an CHO (Chinese hamster ovary) cell or a COS cell.

[0055] Accordingly, a still further aspect of the present invention is a method of producing antibody proteins according to the invention. Such a method comprises the steps of:

- (a) cultivating a host cell as described above under conditions where said antibody protein is expressed by said host cell, and
- (b) isolating said antibody protein.

[0056] Mammalian host cells, preferably CHO or COS cells are preferred. Host cells for producing the antibody proteins of the invention may be transfected with a single vector containing the expression units for both, the light and the heavy chain. In one particular embodiment the method of producing antibody proteins according to the invention pertains to host cells, wherein said host cells are cotransfected with two plasmids carrying the expression units for the light and heavy chains respectively.

[0057] The antibody proteins of the invention provide a highly specific tool for targeting therapeutic agents to the FAP antigen. Therefore, in a further aspect, the invention relates to antibody proteins according to the invention, wherein said antibody protein is conjugated to a therapeutic agent. Of the many therapeutic agents known in the art, therapeutic agents selected from the group consisting of radioisotopes, toxins, toxoids, inflammatoric agents, enzymes, anti-sense molecules, peptides, cytokines, and chemotherapeutic agents are preferred.

[0058] Among the radioisotopes gamma, beta and alpha-emitting radioisotopes may be used as a therapeutic agent. β -emitting radioisotopes are preferred as therapeutic radioisotopes. ^{186}Re henium, ^{188}Re henium, ^{131}I odine and ^{90}Y trium have been proven to be particularly useful β -emitting isotopes to achieve localized irradiation and destruction of malignant tumor cells. Therefore, radioisotopes selected from the group consisting of ^{186}Re henium, ^{188}Re henium, ^{131}I odine and ^{90}Y trium are particularly preferred as therapeutic agents conjugated to the antibody proteins of the invention.

[0059] A further aspect of the present invention pertains to antibody proteins according to the invention, characterised in that they are labeled. Such an FAP-specific labeled antibody allows for the localisation and/or detection of the FAP antigen *in vitro* and/or *in vivo*. A label is defined as a marker that may be directly or indirectly detectable. An indirect marker is defined as a marker that cannot be detected by itself but needs a further directly detectable marker specific for the indirect marker. Preferred labels for practicing the invention are detectable markers. From the large variety of detectable markers, a detectable marker selected from the group consisting of enzymes, dyes, radioisotopes, and biotin is most preferred.

[0060] A further aspect of the present invention relates to antibody proteins according to the invention, characterised

in that they are conjugated to an imageable agent. A large variety of imageable agents, especially radioisotopes, are available from the state of the art. For practicing the invention gamma-emitting isotopes are more preferred. Most preferred is ¹²⁵Iodine.

[0051] One aspect of the present invention relates to pharmaceutical compositions containing an antibody protein according to the present invention as described above and a pharmaceutically acceptable carrier useful for treating tumors, wherein said tumors are associated with activated stromal fibroblasts. There are two possible effector principles for an anti-tumor stroma immunotherapy that may act synergistically: (a) An unmodified (unconjugated, 'naked') antibody according to the invention may induce immune destruction or inflammatory reactions in the tumor stroma while (b) an antibody conjugated to a therapeutic agent, such as for example, a radioisotope or other toxic substance, may achieve localized irradiation and destruction of the malignant tumor cells.

[0052] One further embodiment are pharmaceutical compositions containing an antibody protein according to the invention conjugated to a therapeutic agent as described above and a pharmaceutically acceptable carrier useful for treating tumors, wherein said tumors are associated with activated stromal fibroblasts. Another embodiment pertains to pharmaceutical compositions containing an antibody protein according to the present invention conjugated to an imageable agent as described above and a pharmaceutically acceptable carrier useful for imaging the presence of activated stromal fibroblasts in a healing wound, inflamed skin or a tumor, in a human patient. A most preferred embodiment relates to the pharmaceutical compositions mentioned above, wherein said tumors are tumors selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, invasive bladder cancers, pancreatic cancers and cancers metastatic of the brain.

[0053] In an animal or human body, it can prove advantageous to apply the pharmaceutical compositions as described above via an intravenous or other route, e.g. systemically, locally or topically to the tissue or organ of interest, depending on the type and origin of the disease or problem treated, e.g. a tumor. For example, a systemic mode of action is desired when different organs or organ systems are in need of treatment as in e.g. systemic autoimmune diseases, or allergies, or transplantations of foreign organs or tissues, or tumors that are diffuse or difficult to localise. A local mode of action would be considered when only local manifestations of neoplastic or immunologic action are expected, such as, for example local tumors.

[0054] The antibody proteins of the present invention may be applied by different routes of application known to the expert, notably intravenous injection or direct injection into target tissues. For systemic application, the intravenous, intravascular, intramuscular, intraarterial, intraperitoneal, oral, or intrathecal route are preferred.

[0055] A more local application can be effected subcutaneously, intracutaneously, intracardially, intralobally, intramedullary, intrapulmonarily or directly in or near the tissue to be treated (connective-, bone-, muscle-, nerve-, epithelial tissue). Depending on the desired duration and effectiveness of the treatment, pharmaceutical antibody compositions may be administered once or several times, also intermittently, for instance on a daily basis for several days, weeks or months and in different dosages.

[0056] For preparing suitable antibody preparations for the applications described above, the expert may use known injectable, physiologically acceptable sterile solutions. For preparing a ready-to-use solution for parenteral injection or infusion, aqueous isotonic solutions, such as e.g. saline or corresponding plasmaprotein solutions are readily available. The pharmaceutical compositions may be present as lyophilisates or dry preparations, which can be reconstituted with a known injectable solution directly before use under sterile conditions, e.g. as a kit of parts. The final preparation of the antibody compositions of the present invention are prepared for injection, infusion or perfusion by mixing purified antibodies according to the invention with a sterile physiologically acceptable solution, that may be supplemented with known carrier substances or/and additives (e.g. serum albumine, dextrose, sodium bisulfite, EDTA).

[0057] The amount of the antibody applied depends on the nature of the disease.

[0058] Furthermore, one aspect of the present invention relates to the use of the antibody proteins according to the invention for the treatment of cancer. In a preferred embodiment the present invention relates to the use of antibody proteins according to the invention conjugated to a therapeutic agent as described above for the treatment of cancer. In another preferred embodiment the present invention relates to the use of antibody proteins according to the invention conjugated to an imageable agent for imaging activated stromal fibroblasts. In a further preferred embodiment the present invention relates to the use of labeled antibody proteins according to the invention for detecting the presence of activated stromal fibroblasts in a sample.

[0059] One aspect of the invention relates to a method of treating tumors, wherein the tumor is associated with activated stromal fibroblasts capable of specifically forming a complex with antibody proteins according to the invention, present as naked/unmodified antibodies, modified antibody proteins, such as e.g. fusion proteins, or antibody proteins conjugated to a therapeutic agent, which comprises contacting the tumor with an effective amount of said antibodies. In a preferred embodiment the present invention relates to a method of treating tumors as mentioned above, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, invasive bladder cancers, pancreatic cancers and metastatic cancers of the brain. The method of treating tumors as described above may be effected in

in vitro or *in vivo*.

[0070] A further aspect of the invention relates to a method of detecting the presence of activated stromal fibroblasts in wound healing, inflammation or in tumors, characterised in that

- 5 (a) a sample, possibly containing activated stromal fibroblasts, is contacted with an antibody protein according to the invention under conditions suitable for the formation of a complex between said antibody and antigen,
- (b) detecting the presence of said complex, thereby detecting the presence of activated stromal fibroblasts in wound healing, inflammation or a tumor.

10 [0071] In a preferred embodiment, the present invention relates to a method of detecting the presence of activated stromal fibroblasts in a tumor, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain. Most preferred antibody proteins of the invention are those which are characterised in that they are labeled as mentioned above.

15 [0072] A further aspect of the invention relates to a method of imaging the presence of activated stromal fibroblasts in a healing wound, inflamed skin or a tumor, in a human patient, characterised in that

- (a) an antibody protein according to the present invention conjugated to an imageable agent is administered to a human patient under conditions suitable for the formation of an antibody-antigen complex,
- 20 (b) imaging any complex formed in this manner,
- (c) thereby imaging the presence of activated stromal fibroblasts in a human patient.

[0073] In a preferred embodiment the present invention relates to a method of imaging the presence of activated stromal fibroblasts as described above in tumors, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

[0074] In a further aspect the present invention relates to a method of detecting tumor-stroma, characterised in that

- 30 (a) a suitable sample is contacted with an antibody protein according to the present invention, under conditions suitable for the formation of an antibody-antigen complex,
- (b) detecting the presence of any complex so formed,
- (c) relating the presence of said complex to the presence of tumor-stroma.

[0075] Antibody proteins for practicing the invention are preferably labelled with a detectable marker.

35 [0076] In a further aspect the present invention relates to a method of imaging tumor-stroma in a human patient, which comprises

- (a) administering to the patient an antibody according to the invention conjugated to an imageable agent as described above under conditions suitable for the formation of an antibody-antigen complex,
- 40 (b) imaging any complex so formed, and thereby imaging the presence of tumor-stroma in a human patient.

Figure legends

[0077]

45

Fig. 1. DNA sequence of F19 human reshaped light chain variable region version A (hF19L_A) SEQ ID NO:1.

Fig. 2. Amino acid sequence of F19 human reshaped light chain variable region version A (hF19L_A) SEQ ID NO: 2.

50

Fig. 3. DNA sequence of F19 human reshaped light chain variable region version B (hF19L_B) SEQ ID NO: 3. Nucleotides differing from version A are underlined and in bold type.

Fig. 4. Amino acid sequence of F19 human reshaped light chain variable region version B (hF19L_B) SEQ ID NO: 4. Amino acids differing from version A are underlined and in bold type.

55

Fig. 5. DNA sequence of F19 human reshaped light chain variable region version C (hF19L_C) SEQ ID NO:5. Nucleotides differing from version A are underlined and in bold type.

Fig. 6. Amino acid sequence of F19 human reshaped light chain variable region version C (hF19L_C) SEQ ID NO: 6. Amino acids differing from version A are underlined and in bold type.

Fig. 7. DNA sequence of F19 human reshaped variable region heavy chain version A (hF19H_A) SEQ ID NO: 7.

Fig. 8. Amino acid sequence of F19 human reshaped heavy chain variable region version A (hF19H_A) SEQ ID NO: 8

Fig. 9. DNA sequence of F19 human reshaped heavy chain variable region version B (hF19H_B) SEQ ID NO: 9. Nucleotides differing from version A are underlined and in bold type.

Fig. 10. Amino acid sequence of F19 human reshaped heavy chain variable region version B (hF19H_B) SEQ ID NO: 10. Amino acids differing from version A are underlined and in bold type.

Fig. 11. DNA sequence of F19 human reshaped heavy chain variable region version C (hF19H_C) SEQ ID NO: 11. Nucleotides differing from version A are underlined and in bold type.

Fig. 12. Amino acid sequence of F19 human reshaped heavy chain variable region version C (hF19H_C) SEQ ID NO: 12. Amino acids differing from version A are underlined and in bold type.

Fig. 13. DNA sequence of F19 human reshaped heavy chain variable region version D (hF19H_D) SEQ ID NO: 13. Nucleotides differing from version A are underlined and in bold type.

Fig. 14. Amino acid sequence of F19 human reshaped heavy chain variable region version D (hF19H_D) SEQ ID NO: 14. Amino acids differing from version A are underlined and in bold type.

Fig. 15. DNA sequence of F19 human reshaped heavy chain variable region version E (hF19H_E) SEQ ID NO: 15. Nucleotides differing from version A are underlined and in bold type.

Fig. 16. Amino acid sequence of F19 human reshaped heavy chain variable region version E (hF19H_E) SEQ ID NO: 16. Amino acids differing from version A are underlined and in bold type

Fig. 17. Amino acid sequence of F19 chimeric light chain variable region (chF19LC) SEQ ID NO: 17.

Fig. 18. Amino acid sequence of F19 chimeric heavy chain variable region (chF19HC) SEQ ID NO: 18.

Fig. 19. DNA sequence of human kappa light constant chain SEQ ID NO: 19.

Fig. 20. Amino acid sequence of human light constant chain SEQ ID NO: 20.

Fig. 21. DNA sequence of human heavy constant chain SEQ ID NO: 21.

Fig. 22. Amino acid sequence of human heavy constant chain SEQ ID NO: 22.

Fig. 23. Mammalian cell expression vectors used to produce chimeric and reshaped human antibodies with human kappa light chains and human gamma-1 heavy chains.

A. Light chain expression vector: pKN100

B. Heavy chain expression vector: pG1D105

Fig 24. DNA and amino acid sequences of mouse F19 light chain variable region as modified for use in the construction of chimeric F19 light chain. Restriction sites are indicated by bold letters. The Kozak sequence, CDR's 1 to 3 and the splice donor site are underlined.

Fig 25. DNA and amino acid sequences of mouse F19 heavy chain variable region as modified for use in the construction of chimeric F19 heavy chain. Restriction sites are indicated by bold letters. The Kozak sequence and the splice donor site are underlined.

Fig. 26. DNA sequence of F19 chimeric antibody cloned into pKN100 mammalian expression vector. Restriction sites are indicated by bold letters and underlined. CDR's 1 to 3 and the splice donor site are underlined. This is the DNA sequence of the mouse F19 light chain inside the pKN100 eukaryotic expression vector. This vector has a cDNA version of the human kappa constant region gene (allotype Km(3)) terminated by a strong artificial termination sequence. In addition, the Neo selection gene is also terminated by this artificial sequence and is also in the same orientation as the kappa light chain expression cassette.

The essential components of the pKN100 eukaryotic expression vector are:

- 1 - 6 = EcoRI site
- 7 - 1571 = HCMVi promoter/enhancer
- 583 - 587 = TATAA box
- 610 = Start of transcription
- 728 - 736 = Splice donor site
- 731 = Beginning of intron
- 1557 = End of intron
- 1544 - 1558 = Splice acceptor site
- 1590 - 1598 = Kozak sequence
- 1599 - 1658 = peptide leader sequence
- 1659 - 1997 = mouse F19 light chain
- 1996 - 2004 = splice donor site
- 2011 - 2657 = cDNA copy of human Kappa constant region (Km(3)) gene
- 2664 - 2880 = Artificial spaC2 termination sequence
- 2887 - 7845 = This is the pSV2neo vector DNA fragment comprising of the Amp-resistance gene (in the opposite orientation), the ColEI and SV40 origins of replication and the Neo-resistance gene (in the same orientation as the HCMVi-KCT cassette)
- 7852 - 8068 = Artificial spaC2 termination signal

This sequence ends immediately upstream of the EcoRI site (position 1-6) at the beginning of the sequence. As a vector this DNA sequence would be circular.

Fig. 27. DNA sequence of F19 chimeric antibody cloned into pg1d105 mammalian expression vector. Restriction sites are indicated by bold letters and underlined. CDR's 1 to 3 and the splice donor site are underlined. This is the DNA sequence of the eukaryotic expression vector pG1D105 containing the mouse F19 heavy chain variable region. This vector contains a cDNA version of the human gamma-1 constant region (allotype G1m^{Non-a}).

The essential components of the construct are:

- 1 - 2501 = pBR322 based sequence including Ampicillin resistance gene and ColEI origin plus the SV40 origin and the crippled SV40 early promoter
- 2502 - 3226 = dhfr gene
- 3233 - 4073 = SV40 poly A sequence etc.
- 4074 - 4079 = ligated BamHI and BglII site (BstYI)
- 4080 - 4302 = SPA site plus C2 termination signal
- 4303 - 5867 = HCMVi promoter
- 5879 - 5885 = unique HindIII restriction site for cloning of immunoglobulin variable genes
- 5886 - 5894 = Kozak sequence
- 5895 - 5951 = signal peptide
- 5952 - 6323 = mouse F19 heavy chain
- 6323 - 6330 = splice donor site
- 6331 - 6336 = unique BamHI restriction site for cloning of immunoglobulin variable genes
- 6337 - 7388 = cDNA copy of human gamma-1 constant regions preceded by a 62 bp intron
- 7389 - 7709 = Arnie termination sequence

The human gamma-1 constant region used in this construct has a G1m^{Non-a} allotype which is defined by a Glutamic acid (E) residue at position 356 (according to Eu numbering) and a Methionine (M) residue at position 358 (according to Eu numbering). These two residues are underlined in the sequence above.

Fig. 28. PCR-based method for the construction of human reshaped F19 light chain. This figure provides a schematic overview of the strategy of construction. The dotted lines indicate a complementary sequence of at least 21

bases between the primers.

Fig. 29. Nucleotide and deduced amino acid sequences of reshaped human F19 light chain variable regions version A, B and C. Nucleotide and deduced amino acid sequences are aligned and compared with that of version A, dashes indicate nucleotide identity, dots indicate amino acid identity with this sequence. Amino acids are numbered according to Kabat *et al.* (1991). The locations of CDRs are indicated in boxes.

Fig. 30. DNA sequence of F19 L_A (human reshaped light chain version A) cloned into pKN100 mammalian expression vector. Restriction sites are indicated by bold letters and underlined. CDR's 1 to 3 and the splice donor site are underlined. This is the DNA sequence of the reshaped F19 light chain version A cloned into pKN100 eukaryotic expression vector. This vector has a cDNA version of the human kappa constant region gene (allotype Km(3)) terminated by a strong artificial termination sequence. In addition, the Neo selection gene is also terminated by this artificial sequence and is also in the same orientation as the kappa light chain expression cassette.

The components of the vector are:

7 - 1571	= HCMVi promoter/enhancer
583 - 587	= TATAA box.
610	= Start of transcription.
728 - 736	= Splice donor site.
731	= Beginning of intron.
1557	= End of intron.
1544 - 1558	= Splice acceptor site.
1590 - 1598	= Kozak sequence
1599 - 1658	= peptide leader sequence
1659 - 1997	= reshaped F19 light chain version A
1996 - 2004	= splice donor site
2011 - 2657	= cDNA copy of human kappa constant region (Km(3)) gene.
2664 - 2880	= Artificial spaC2 termination sequence.
2887 - 7845	= This is the pSV2neo vector DNA fragment comprising of the Amp-resistance gene (in the opposite orientation), the ColEI and SV40 origins of replication and the Neo-resistance gene (in the same orientation as the HCMVi-KCT cassette).
7852 - 8068	= Artificial spaC2 termination signal.

This sequence ends immediately upstream of the EcoRI site (position 1-6) at the beginning of the sequence below. As a vector this DNA sequence would be circular.

Fig. 31. PCR-based method for the construction of human reshaped F19 heavy chain. This figure provides a schematic overview of the strategy of construction. The dotted lines indicate a complementary sequence of at least 21 bases between the primers.

Fig. 32. Nucleotide and deduced amino acid sequences of reshaped human F19 heavy chain variable region versions a to e. Nucleotide and deduced amino acid sequences are aligned and compared with that of version A, dashes indicate nucleotide identity, dots indicate amino acid identity with this sequence. Amino acids are numbered according to Kabat *et al.* (1991). The location of CDRs is indicated by boxes.

Fig. 33. DNA sequence of F19Ha (human reshaped heavy chain version a) cloned into pg1d105 mammalian expression vector. Restriction sites are indicated by bold letters and underlined. CDR's 1 to 3 and the splice donor site are underlined. This is the DNA sequence of the eukaryotic expression vector pG1D105 containing the reshaped version A of F19 heavy chain variable region. This vector contains a cDNA version of the human gamma-1 constant region (allotype G1m^{Non-a}).

The essential components of the construct are:

1 - 2501	= pBR322 based sequence including Ampicillin resistance gene and ColEI origin plus the SV40 origin and the crippled SV40 early promoter
2502 - 3226	= dhfr gene
3233 - 4073	= SV40 poly A sequence etc.
4080 - 4302	= SPA site plus C2 termination signal
4303 - 5867	= HCMVi promoter/enhancer

5879 - 5885 = unique HindIII restriction site for cloning of immunoglobulin variable genes
 5886 - 5894 = Kozak sequence
 5895 - 5951 = signal peptide
 5952 - 6323 = reshaped F19 heavy chain version A
 6323 - 6330 = splice donor site
 6331 - 6336 = unique BamHI restriction site for cloning of immunoglobulin variable genes
 6337 - 7388 = cDNA copy of human gamma-1 constant regions preceded by a 62 bp intron
 7389 - 7709 = Arnie termination sequence

The human gamma-1 constant region used in this construct has a G1m^{Non-a} allotype which is defined by a Glutamic acid (E) residue at position 356 (according to Eu numbering) and a Methionine (M) residue at position 358 (according to Eu numbering). These two residues are underlined in the sequence above.

Fig. 34. Heavy (panel A) and light (panel B) chains RNA splicing events taking place during antibody F19 expression in mammalian cells - schematic overview.

A. Heavy chain RNA splicing

B. Kappa light chain RNA splicing

Fig. 35. Concentration dependence of L_AH_C supernatant binding to CD8-FAP.

Fig. 36. Binding of biotinylated L_AH_C to human FAP.

Fig. 37. CD8-FAP carries the F19 epitope as detected with cF19.

Examples

Example 1: Construction of mouse - human chimeric genes

[0078] The chimeric F19 (cF19) antibody was designed to have the mouse F19 V_L and V_H regions linked to human kappa and gamma-1 constant regions, respectively. PCR primers were used to modify the 5'- and 3'- sequences flanking the cDNA sequences coding for the mouse F19 V_L and V_H regions (Table 1). PCR primers specific for F19 light chain V-region were designed. These adapted mouse F19 variable regions were then subcloned into mammalian cell expression vectors already containing the human kappa (pKN100 vector) or gamma-1 (pG1D105 vector) constant regions (Figure 23).

[0079] These vectors employ the human cytomegalovirus (HCMV) promoter/enhancer to efficiently transcribe the light and heavy chains. The vectors also contain the SV40 origin of replication to permit efficient DNA replication and subsequent protein expression in cos cells. The expression vectors were designed to have the variable regions inserted as HindIII-BamHI DNA fragments. PCR primers were designed to introduce these restrictions sites at the 5'- (HindIII) and 3'- (BamHI) ends of the cDNAs coding for the V-regions. In addition the PCR primers were designed to introduce the Kozak sequence (GCCGCCACC) at the 5'-ends of both the light and heavy chain cDNAs to allow efficient translation (Kozak M.: At least six nucleotides preceding the AUG initiator codon enhance translation in mammalian cells. *J. Mol. Biol.* (1987) 196: 947), and to introduce splice donor sites at the 3'-ends of both the light and heavy chain cDNAs for the variable regions to be spliced to the constant regions. The PCR primers used in the construction of the chimeric F19 light and heavy chains are shown in Table 1. The DNA and amino acid sequences of the mouse F19 V_L and V_H regions as adapted for use in the construction of chimeric F19 light and heavy chains are shown in Figures 24 and 25. The DNA sequences of mouse F19 light and heavy chains cloned into the eukaryotic expression vectors pKN100 and pG1D105, respectively, are shown in Figures 26 and 27.

TABLE 1: PCR primers for the construction of chimeric F19 antibody.**A. Light chain variable region**

1. Primer for the construction of the 5'-end (37mer)

5' CAGA AAGCTT GCCGCCACC ATG GAT TCA CAG GCC CAG 3'HindIII Kozak sequence M D S Q A Q

2. Primer for the construction of the 3'-end (35mer)

5' CCGA GGATCC ACTCACG TTT CAG CTC CAG CTT GGT 3'BamHI Splice donor site**B. Heavy chain variable region**

1. Primer for the construction of the 5'-end (37mer)

5' CAGA AAGCTT GCCGCCACC ATG GGA TGG AGC TGG GTC 3'HindIII Kozak sequence M G W S W V

2. Primer for the construction of the 3'-end (35mer)

5' CCGA GGATCC ACTCACG TGA GGA GAC GGT GAC TGA 3'BamHI Splice donor site**Example 2: Expression and binding activity of chimeric F19 antibody**

[0080] The two plasmid DNAs coding for the chimeric F19 light and heavy chains (see example 1) were co-transfected into cos cells to look for transient expression of chimeric F19 antibody as described below. After 72 h incubation, the medium was collected, centrifuged to remove cellular debris, and analysed by ELISA for the production of a human IgG1-like antibody. The cos cell supernatant containing the chimeric F19 antibody was analysed for its ability to bind to HT 1080 cells (see example 13) expressing the FAP antigen on their surface.

Transfection of cos cells using electroporation

[0081] The mammalian expression vectors pg1d105 and pKN100 containing the chimeric or reshaped human heavy and light chains versions, respectively, were tested in cos cells to look for transient expression of F19 antibodies. Cos

7 cells were passaged routinely in DMEM (Gibco BRL cat. #41966) containing penicillin (50 IU/ml), streptomycin (50 µg/ml), L-glutamine and 10% heat-inactivated gamma globulin-free foetal calf serum (FCS, Harlan Sera-Lab cat. # D0001). The DNA was introduced into the cos cells by electroporation using the Gene Pulsar apparatus (BioRad). DNA (10 µg of each vector) was added to a 0.8 ml aliquot of 1×10^7 cells/ml in Phosphate-buffered saline (PBS, Ca^{2+} and Mg^{2+} free). A pulse was delivered at 1,900 volts, 25 µF capacitance. After a 10 min recovery period at ambient temperature the electroporated cells were added to 8 ml of DMEM containing 5% FCS. After 72h incubation at 37°C, the medium was collected, centrifuged to remove cellular debris, and stored under sterile conditions at 4°C for short periods of time, or at -20°C for longer periods.

ELISA method for measuring assembled IgG1/kappa antibody concentrations in cos cell supernatants

[0082] Samples of antibodies produced in transfected cos cells were assayed by ELISA to determine how much reshaped human antibody had been produced. For the detection of human antibody, plates were coated with goat anti-human IgG (Fcγ fragment specific) antibody (Jackson ImmunoResearch Laboratories Inc., #109-005-098). The samples from cos cells were serially diluted and added to each well. After incubation for 1h at 37°C and washing, horseradish peroxidase conjugated goat anti-human kappa light chain (Sigma, A-7164) was added. After incubation for 30 mins at 37°C and washing, K-blue substrate (mixer of 3,3',5,5' tetramethylbenzidine and hydrogen peroxide, Bionostics Limited, #KB175) was added. After standing at room temperature for 30 mins, the reaction was stopped using Red Stop solution (Bionostics Limited, #RS20) and the optical density read on a microplate reader at 650 nm. Purified human IgG1/Kappa antibody (Sigma, I-3889) of known concentration was used as a standard.

[0083] The expression of chimeric F19 antibody in COS cells was poor (Table 2), between 10 and 60 ng/ml which is at least 10 fold less than most antibodies.

[0084] In an attempt to increase expression levels of the chimeric F19 antibody, the leader sequence of F19 V_L region was changed by substitution of Leucine to Proline at position -9. This single change in amino acid in the leader sequence resulted in at least doubling the amount of chimeric antibody produced in COS cells.

[0085] The test results show that chimeric F19 binds specifically and with the expected avidity to the FAP target.

TABLE 2

Chimeric F19 antibody concentrations in COS cell supernatants (These are the results of three independent transfections)		
Transfected Antibody components		Human γ1/K
Heavy chain	Kappa light chain	[in µg/ml]
cF19	cF19 (F19 leader sequence)	0.060
cF19	cF19 (mutated leader sequence)	0.212
cF19	cF19 (F19 leader sequence)	0.056
cF19	cF19 (mutated leader sequence)	0.108
cF19	cF19 (F19 leader sequence)	0.011
cF19	cF19 (mutated leader sequence)	0.087

Example 3: Construction of the reshaped human F19 light chain versions a to c (La-Lb)

[0086] The construction of the first version of reshaped human F19 V_L region (La) was carried out using overlapping PCR fragments in a method similar to that described by Daugherty B. L., DeMartino J. A., Law M. F., Kawka D. W., Singer I. I. and Mark G. E. (1991) Polymerase chain reaction facilitates the cloning, CDR-grafting, and rapid expression of a murine monoclonal antibody directed against the CD18 component of leukocyte integrins. *Nucl. Acids Res.* 19: 2471. Ten oligonucleotides were synthesised that consisted of five primer pairs, APCR1-vla1, vla2-vla3, vla4-vla5, vla6-vla7, and vla8-APCR4 (Table 3 and Figure 28). There was an overlapping sequence of at least 21 bases between adjacent pairs (Figure 28). APCR1 and APCR4 hybridised to the flanking pUC19 vector sequences. The mutagenic primers were designed such that their 5' end immediately followed the wobble position of a codon. This strategy was used to counteract the gratuitous addition of one nucleotide to the 3' end of the strand complementary to the mutagenic primer by the DNA polymerase during PCR (Sharrocks A. D. and Shaw P. E. (1992) Improved primer design for PCR-based, site-directed mutagenesis. *Nucl. Acids Res.* 20: 1147). The appropriate primer pairs (0.2 µM of each) were combined

with 10ng of version "b" of reshaped human L25V_L region cDNA, and 1 unit of AmpliTaq (Perkin Elmer Cetus) DNA polymerase in 50µl of PCR buffer containing 10mM Tris-HCl (pH8.3), 50mM KCl, 200µM dNTPs, and 1.5mM MgCl₂. This was overlaid with mineral oil and PCR was performed for 25 cycles, each cycle consisting of a denaturation step at 94°C for 1 min, a primer annealing step at 55°C for 1 min, and an extension step at 72°C for 2 mins. This was followed by a single cycle consisting of a further elongation step at 72°C for 10 mins followed by cooling to 4°C. The ramp time between the primer-annealing and extension steps was 2.5 mins. The PCR products of the five reactions (A, B, C, D and E) were then purified by gel electrophoresis followed by DNA elution using Wizard PCR preps (Promega). PCR products A, B, C, D, and E were assembled by their complementarity to one another. In the second set of PCR reactions, PCR products B and C, and D and E, (50ng of each) were added to 50µl PCR reactions (as described above) each containing 1 unit of AmpliTaq (Perkin Elmer Cetus) DNA polymerase. The reactions were cycled for 20 cycles as described above with the exception that the annealing temperature was raised to 60°C. In the third set of PCR reactions, PCR products F and G were PCR-amplified using 1 µl of each prior PCR reaction and the appropriate pair of PCR primers (via2-vla5 or vla6-APCR4). The PCR reactions contained 1 unit of AmpliTaq DNA polymerase in 50 µl PCR reaction (as described above) and were amplified for 25 cycles as in the first stage. In the fourth set of PCR reactions, the PCR product H was PCR-amplified using 1 µl of each prior PCR reaction and the vla2-APCR4 pair of PCR primers. Finally, PCR products A and H were assembled by their own complementarity in a two step-PCR reaction similar to that described above using RSP and UP as the terminal primers. The fully assembled fragment representing the entire reshaped human F19 V_L region including a leader sequence was digested with HindIII and BamHI and cloned into pUC19 for sequencing. A clone having the correct DNA sequence was designated reshF19La (Figure 29) and was then subcloned into the eukaryotic expression vector pKN100. The DNA sequence of reshF19La cloned into pKN100 is shown in Figure 30.

[0087] The second version of reshaped human F19 V_L region (Lb) was constructed using the same scheme as that described for La but where vla4 and vla7 primers were substituted by vlb4 and vlb7 respectively (Table 3). The DNA sequence of Lb is shown in Figure 29.

[0088] The third version of reshaped human F19 V_L region (Lc) was constructed using the QuikChange™ site-directed mutagenesis kit from Stratagene. The QuikChange site-directed mutagenesis method was performed according to the manufacturer's instructions, using reshF19La in pKN100 vector as double stranded DNA template. The mutagenic oligonucleotide primers F19Lc-sense and F19Lc-antisense (Table 3) for use in this protocol were designed according to the manufacturers instructions. Briefly, both the mutagenic primers contained the desired point mutation (codon TTT at Kabat residue position 49 (Phe) changed to TAT coding for Tyr) and annealed to the same sequence on opposite strands of La in pKN100 vector. The point mutation was verified by DNA sequencing the entire V_L region. The DNA sequence of Lc is shown in Figure 29. To eliminate the possibility that random mutations occurred in the pKN100 during the PCR reaction, the V_L region was cut out of the pKN100 vector as an HindIII/BamHI fragment and re-subcloned into an unmodified pKN100 vector cut with the same two restriction enzymes beforehand.

TABLE 3: PCR primers for the construction of reshaped human F19 light chain variable regions

1. Primers for the synthesis of version "a"

F19vla1 (36 mer):

5' GTCATCACAATGTCTCCGGAGGAACCTGGAACCCAG 3'

F19vla2 (29 mer):

5' CTCCGGAGACATTGTGATGACCCAATCTC 3'

F19vla3 (45 mer):

5' GAATATAAAAGGCTCTGACTGGACTTGCAGTTGATGGTGGCCCTC 3'

F19v1a4 (72 mer):

5' CAGTCAGAGCCTTTTATATTCTAGAAATCAAAAGAACTACTTGGCCTGGTAT
CAGCAGAAACCAGGACAGCC 3'

F19v1a5 (44 mer):

5' ACCCCAGATTCCCTAGTGCTAGCCCCAAAAGATGAGGAGTTTGGG 3'

F19v1a6 (67 mer):

5' TAGCACTAGGGAATCTGGGGTACCTGATAGGTTCACTGGCAGTGGGTTTG
GGACAGACTTCACCCTC 3'

F19v1a7 (53 mer):

5' GTCCCTTGTCGGAACGTGAGCGGATAGCTAAAATATTGCTGACAGTAA
TAAAC 3'

F19v1a8 (33 mer):

5' GCTCACGTTCGGACAAGGGACCAAGGTGGAAAT 3'

2. Primers for the synthesis of version "b"

F19v1b4 (72 mer):

5' CAGTCAGAGCCTTTTATATTCTAGAAATCAAAAGAACTACTTGGCCTGG
TTCCAGCAGAAACCAGGACAGCC 3'

F19v1b7 (57 mer):

5' GTCCCTTGTCGGAACGTGAGCGGATAGCTAAAATATTGCTGACAGTCATA
AACTGCC 3'

3. Primers for the synthesis of version "c"

F19Lc-sense (34 mer):

5' CCCAAACTCCTCATCTATTGGGCTAGCACTAGGG 3'

F19Lc-antisense (34 mer):

5' CCCTAGTGCTAGCCCAATAGATGAGGAGTTTGGG 3'

4. Primers hybridizing to the flanking PUC19 vector sequences

APCR1 (17 mer, sense primer): 5' TACGCAAACCGCCTCTC 3'

APCR4 (18 mer, anti-sense primer): 5' GAGTGCACCATATGCGGT 3'

RSP (-24) (16 mer, sense primer): 5' AACAGCTATGACCATG 3'

UP (-40) (17 mer, anti-sense primer): 5' GTTTTCCCAGTCACGAC 3'

Example 4: Construction of the reshaped human F19 heavy chain versions a to e (Ha-He)

[0089] Version "a" of reshaped human F19 V_H regions (Ha) was constructed using the same PCR methods as described for the construction of version "a" of reshaped human F19 V_L region (La) (Figure 31). The template DNA was version "a" of reshaped human 226 V_H (Léger O. J. P., Yednock T. A., Tanner L., Horner H. C., Hines D. K., Keen S., Saldanha J., Jones T., Fritz L. C. and Bendig M. M. (1997). Humanization of a mouse antibody against human alpha-4 integrin: a potential therapeutic for the treatment of multiple sclerosis. *Hum. Antibod.* 8: 3). Six PCR primers were designed and synthesized for the construction of version "a" of reshaped human F19 V_H region (Table 4). PCR products A, B, C, and D were obtained using APCR1-Vha1, Vha2-Vha3, Vha4-Vha5 and Vha6-APCR4 as PCR primer pairs, respectively. The PCR conditions were essentially as described for the construction of reshaped human F19 V_L region. A clone having the correct DNA sequence was designated reshF19Ha (Figure 32) and was then subcloned into the eukaryotic expression vector pG1D105. The DNA sequence of reshF19Ha cloned into pG1D105 is shown in Figure 33.

[0090] The third version of reshaped human F19 V_H region (Hc) was constructed using the same scheme as that described for Ha but where Vha4 primer was substituted by Vhc4 (Table 4). The DNA sequence of Hc is shown in Figure 32. The second (Hb) and fourth (Hd) version of reshaped human F19 V_H region were constructed based on the PCR-mutagenesis methods of Kamman et al. (Kamman M., Laufs J., Schell J. and Gronenborn B. (1989) Rapid insertional mutagenesis of DNA by polymerase chain reaction (PCR). *Nucl. Acids Res.* 17: 5404). For Hb and Hd, a mutagenic primer F19VHbd6 (Tyr-91 to Phe-91, Table 4) was used paired with APCR4 in PCR reactions with Ha and Hc as the template DNA, respectively. The PCR products VHb and VHD were restriction enzyme digested with PstI and BamHI and subcloned into reshF19Ha and reshF19Hc, respectively, previously digested with the same two restriction enzymes. The DNA sequences of Hb and Hd are shown in Figure 32.

[0091] Version e of reshaped human F19 V_H region (He) was constructed based on the PCR-mutagenesis methods of Kamman et al. (1989) already mentioned above:

[0092] For reshF19He mutagenic primer F19MscIHe (Table 5) was used paired with primer F19V_HHindIII (Table 5) in PCR reactions with Hc cloned in pg1d105 mammalian expression vector as the template DNA. The appropriate primer pairs (0.2 μM of each) were combined with 10ng of cDNA of version "a" of reshaped human 226 V_H region in 100 μl of PCR buffer containing 10mM KCl, 10mM (NH₄)₂SO₄, 20mM Tris-HCl (pH 8.8) 2mM MgSO₄, 0.1% Triton X-100 and 200 μM dNTPs. Reaction mixtures were overlaid with mineral oil and kept at 94°C for 5 mins. Then 1 unit of Deep Vent DNA polymerase (New England Biolabs) was added ("Hot Start" PCR; Chou Q., Russell M., Birch D., Raymond J. and Bloch W. (1992) Prevention of pre-PCR mis-priming and primer dimerization improves low-copy-number amplifications. *Nucl. Acids Res.* 20: 1717) and PCR was performed for 25 cycles on a TRIO-Thermoblock Thermal Cycler (Biometra, Göttingen, Germany). Each cycle consisting of a denaturation step at 94°C for 1 min, a primer annealing step at 70°C for 1 min, and an extension step at 72°C for 2 mins. This was followed by a single cycle consisting of a further elongation step at 72°C for 10 mins followed by cooling at 4°C. The PCR products were then extracted and purified from a TAE 1.4% standard agarose gel using a QIAquick™ gel extraction kit, following the protocol supplied by the manufacturer

(QIAGEN Ltd., UK). The PCR product V_He was then restriction enzyme digested with MscI and HindIII and ligated into reshF19Hc cloned in pg1d105 previously digested with the same two restriction enzymes. The MscI restriction recognition site is unique to all the reshaped human F19 V_H region versions and is not present in the pg1d105 expression vector. The HindIII restriction recognition site is a unique site in pg1d105 for clotting of V_H immunoglobulin genes.

- 5 [0093] Electroporation-competent XL-1 Blue E. coli cells were transformed with 1 µl of the ligated DNA and plated on agarose plates containing Ampicillin. Colonies were then screened for the presence and correct size of inserts by direct PCR on colonies (Güssow D. and Clackson T. (1989) Direct clone characterization from plaques and colonies by the polymerase chain reaction. *Nucl. Acids Res.* 17: 4000) with primers HCMi and Hucy1 hybridising to the flanking pg1d105 vector sequences (Table 5). DNA from positive colonies was prepared using a Plasmid Midi kit, following the
- 10 protocol supplied by the manufacturer (QIAGEN Ltd., UK). DNA sequencing was performed by the dideoxy chain termination method (Sanger F., Nicklen S. and Coulson A. (1977) DNA sequencing with chain-terminating inhibitors. *Proc. natn. Acad. Sci. U. S. A.* 74: 5463) directly from circular vector DNA using conventional heat denaturation (Andersen A., Pettersson A. and Kieldsen T. (1992) A fast and simple technique for sequencing plasmid DNA with sequenase using heat denaturation. *Biotechniques* 13: 678) and Sequenase 2.0 (USB, Cleveland, OH). The DNA sequences of
- 15 reshF19He is shown in Figure 32.

TABLE 4: PCR primers for the construction of reshaped human F19 heavy chain variable regions versions a to d.

1. Primers for the synthesis of version "a"

F19vha1 (47mer):

5' GTGTATTCAGTGAAGGTGTATCTACTAGTTTTACAGCTGACTTTTCAC 3'

F19vha2 (53 mer):

5' TAGTAGATACACCTTCACTGAATACACCATACTGGGTTAGACAGG
CCCCTG 3'

F19vha3 (71 mer):

5' CCCTTGAACCTTCTGGTTGTAGTTAGGAATACCATTGTTAGGATTAATACC
TCCTATCCACTCCAGCCTTTG 3'

F19vha4 (71 mer):

5' TAACTACAACCAGAAGTTCAAGGGCCGGGCCACCTTGACCGTAGGCAA
GTCTGCCAGCACCGCCTACATGG 3'

F19vha5 (63 mer):

5' GCATGGCCCTCGTCGTAACCATAGGCGATTCTTCTTCTGGCGCAGTAGT
AGACTGCAGTGTCC 3'

F19vha6 (48 mer):

5' CTATGGTTACGACGAGGGCCATGCTATGGACTACTGGGGTCAAGGAAC 3'

2. Primers for the synthesis of version "c"

F19vhc4 (71 mer):

5' TAACTACAACCAGAAGTTCAAGGGCCGGGTCAACCATCACCGTAGACA
CCTCTGCCAGCACCGCCTACATGG 3'

3. Primers for the synthesis of version "b" and "d"

F19vhbd6 (27 mer):

5' GGACACTGCAGTCTACTTCTGCGCCAG 3'

4. Primers hybridizing to the flanking PUC19 vector sequences

APCR1 (17 mer, sense primer): 5' TACGCAAACCGCCTCTC 3'

APCR4 (18 mer, anti-sense primer): 5' GAGTGCACCATATGCGGT 3'

TABLE 5: PCR primer for the construction of reshaped human F19 heavy chain variable regions version e

1. Primer for the synthesis of version "e"

F19MscIHe (65 mer, anti-sense):

5' CCTT TGGCCAGGGGCCTGTCTAACCCAGTGTATGGTGTATTCAGTGAAGGTG
MscI
TATCCACTAGTTTCCACTAGTTT 3'

2. Primers hybridizing to the flanking pg1d105 mammalian expression vector sequences

HCMi (28 mer, sense): 5' GTCACCGTCCTTGACACGCGTCTCGGGA 3'

Hucy1 (17 mer, anti-sense): 5' TTGGAGGAGGGTGCCAG 3'

Example 5: Reshaped human F19 antibody concentrations in COS cells supernatants

[0094] COS cells were transfected with one pair of a series of reshaped human F19 antibody constructs and the human antibody concentration was measured using the IgG1/Kappa ELISA as described in example 2.

TABLE 6

Reshaped human F19 antibody concentrations in COS cell supernatants		
Transfected Antibody components		Human γ 1/K
Heavy chain	Kappa light chain	concentration [μ g/ml]
Ha	La	2.50
Ha	Lb	0.18
Hb	La	1.25
Hb	Lb	0.10
Hd	La	1.15
Hd	Lb	0.18
Ha	La	1.50
Ha	Lc	1.56

TABLE 6 (continued)

Reshaped human F19 antibody concentrations in COS cell supernatants		
Transfected Antibody components		Human γ 1/K
Heavy chain	Kappa light chain	concentration [μ g/ml]
Hc	La	1.47
Hc	Lc	1.97
cF19	La	1.54
cF19	Lb	0.07
cF19	Lc	2.14

TABLE 7

Reshaped human F19 antibody concentrations in COS cell supernatants		
Transfected Antibody components		Human γ 1/K
Heavy chain	Kappa light chain	concentration [μ g/ml]
Ha	La	2.00
Ha	Lc	2.50
Hc	La	2.90
Hc	Lc	3.00
He	La	2.80
He	Lc	3.50

RNA splicing events required for the expression of immunoglobulin genes in mammalian cells

[0095] Both mammalian expression vectors pKN100 and pg1d105 have an intron between the variable and the constant regions which is removed during the process of gene expression to give rise to an messenger RNA. The splicing event which consists of a DNA recombination between the heavy or light chain splice donor sites and the immunoglobulin splice acceptor site is described in Figure 34.

Example 6: Flow cytometric analysis of the binding of cF19 and L_AH_C to FAP-expressing human cells

[0096] The ability of L_AH_C to bind to both recombinant and endogenously expressed FAP on cell surface was tested.

[0097] The example was conducted to determine the binding of L_AH_C to cellular FAP. Both naturally FAP expressing MF-SH human tumour cells and FAP-transfected human tumour cell lines were used as cellular targets. L_AH_C was studied in cytofluorometric assays evaluating direct binding to target cells as well as by the inhibitory effect on the binding of either murine F19 or chimeric cF19 anti-FAP antibodies.

[0098] Antibodies and cell lines used were F19 (murine monoclonal anti-human FAP antibody, IgG1 subclass), mIgG (murine immunoglobulin, IgG class), cF19 (chimeric monoclonal anti-human FAP antibody, IgG1 subclass), L_AH_C (reshaped monoclonal anti-human FAP antibody, IgG1 subclass), hIgG1 (human immunoglobulin, IgG1 subclass), MF-SH (human malignant fibrous histiocyte cell line), HT-1080 (human fibrosarcoma cell line), HT-1080FAP clone 33 (HT-1080 cell line transfected with cDNA encoding human FAP)

Direct binding of L_AH_C to FAP on the surface of human tumour cell lines

[0099] 5×10^5 cells of the tumour cell line under investigation were incubated with the indicated concentration of test or control antibody in a total volume of 0.2 ml phosphate-buffered saline (PBS) supplemented with 1% bovine serum albumin (BSA) for 30 min on ice.

[0100] Subsequently, cells were washed twice with 2 ml of PBS, resuspended in 0.2 ml of PBS supplemented with 1% BSA, the appropriate anti-Ig-antibody as secondary reagent (either a 1:20 dilution of goat anti-mouse Ig FITC-labeled [Dianova] or a 1:20 dilution of mouse anti-human IgG FITC-labeled [Dianova]) and incubated for another 30 min on ice.

[0101] Cells were again washed twice with 2 ml of PBS, resuspended in a total volume of 0.5 ml of PBS supplemented with 1% paraformaldehyde (PFA) and kept on ice. Single cell fluorescence was determined cytofluorometrically by analysing the cellular green fluorescence in the 488nm light of an EPICS XL (Coulter).

Inhibitory effect of L_AH_C on binding of biotinylated cF19 to FAP on the surface of human cell lines

[0102] 5×10^5 cells of the tumour cell line under investigation were incubated with the indicated concentration of the biotin-labelled antibody in a total volume of 0.2 ml PBS supplemented with 1% BSA and the simultaneously added unlabelled test or control antibody for 30 min on ice. Subsequently, cells were washed twice with 2 ml of PBS, resuspended in 0.2 ml of PBS supplemented with 1% BSA, 1:40 diluted streptavidin-FITC (Dianova) as secondary reagent and incubated for another 30 min on ice.

[0103] Alternatively, cells were incubated with the indicated concentrations of murine F19 and cell-bound antibody detected via 1:20 diluted goat anti-mouse Ig labelled with FITC by comparable incubation steps.

[0104] In each case, cells were finally washed twice with 2 ml of PBS, resuspended in a total volume of 0.5 ml PBS supplemented with 1% PFA and kept on ice. Single cell fluorescence was determined cytofluorometrically by analysing the cellular green fluorescence in the 488nm light of an EPICS XL (Coulter).

[0105] Both, cF19 and L_AH_C bind in a concentration dependent manner specifically to FAP-transfected HT-1080FAP clone33 human tumour cells (Table 8). No binding to FAP-negative HT-1080 cells was detectable (Table 9). Both cF19 and L_AH_C bound in a concentration dependent manner to human MF-SH cells endogenously expressing FAP (Table 10).

[0106] Biotinylated cF19 in a concentration dependent manner bound to human HT-1080FAP clone 33 (Table 11). No binding was detectable to FAP-negative HT-1080 cells (Table 12).

[0107] Binding of biotinylated cF19 to HT-1080FAP clone 33 cells was inhibited by both unlabelled cF19 and unlabelled L_AH_C (Table 13).

[0108] Chimeric anti-human FAP monoclonal antibody cF19 as well as reshaped human anti-human FAP monoclonal antibody L_AH_C (example 10) were shown to bind directly to FAP expressed on human cell lines either endogenously expressing this protein or transfected with cDNA encoding for it. This binding was shown to be concentration dependent. Binding of biotinylated cF19 could be inhibited by both unlabelled cF19 and unlabelled L_AH_C .

[0109] Using cytofluorometric technology, direct binding as well as inhibition of specifically binding reagents showed specificity of chimeric cF19 and reshaped L_AH_C human monoclonal antibodies to cell surface expressed FAP.

Table 8

Binding of anti-FAP antibodies to HT-1080FAP clone 33 cells			
Concentration of anti-body	Mean fluorescence intensity		
[ng/mL]	hlgG1	cF19	L_AH_C
500.0	0.12	6.65	2.76
100.0	0.12	1.63	0.66
20.0	0.12	0.43	0.22
4.0	0.12	0.17	0.15
0.8	0.12	0.14	0.13

Table 9

Binding of anti-FAP antibodies to non-transfected HT-1080 cells			
Concentration of anti-body	Mean fluorescence intensity		
[ng/mL]	hlgG1	cF19	L _A H _C
500.0	0.11	0.11	0.12
100.0	0.11	0.11	0.11
20.0	0.11	0.11	0.12
4.0	0.11	0.11	0.12
0.8	0.11	0.11	0.11

Table 10

Binding of anti-FAP antibodies to MF-SH cells			
Concentration of anti-body	Mean fluorescence intensity		
[ng/mL]	hlgG1	cF19	L _A H _C
4.0	0.6	3.6	2.8
2.0	n.d.	3.3	2.5
1.0	n.d.	2.4	1.9
0.5	n.d.	1.8	1.3

n.d.: not done

Table 11

Binding of biotinylated cF19 antibody to HT-1080FAP clone 33 cells		
Concentration of anti-body	Mean fluorescence intensity	
[ng/ml]	Biotinylated hlgG1	Biotinylated cF19
5,000.0	0.2	36.5
1,000.0	0.2	18.1
200.0	0.2	4.5
40.0	0.2	1.3
8.0	0.2	0.5
1.6	0.3	0.3

Table 12

Binding of biotinylated cF19 antibody to non-transfected HT-1080 cells		
Concentration of anti-body	Mean fluorescence intensity	
[ng/ml]	Biotinylated hlgG1	Biotinylated cF19
5,000.0	0.1	0.1
1,000.0	0.1	0.1
200.0	0.1	0.1
40.0	0.1	0.1
8.0	0.1	0.1
1.6	0.1	0.1

Table 13

Competition of anti-FAP antibodies with the binding of biotinylated cF19 to HT-1080FAP clone 33 cells		
	Concentration of competitor antibody	Mean fluorescence concentration
Competitor antibody	[μ g/mL]	
no	0.00	11.2
hlgG1	1.00	9.0
hlgG1	3.16	11.3
hlgG1	10.00	9.8
hlgG1	31.66	10.3
cF19	1.00	7.5
cF19	3.16	4.8
cF19	10.00	1.3
cF19	31.66	1.2
L _A H _C	1.00	8.0
L _A H _C	3.16	5.5
L _A H _C	10.00	2.9
L _A H _C	31.66	1.7
Biotinylated cF19 was used at a concentration of 1 μ g/mL in all tests shown in the table.		

Example 7: In vitro immune effector functions of monoclonal antibody L_AH_C

[0110] This experiment was conducted to determine the potential of the monoclonal antibody (mab) L_AH_C with specificity for fibroblast activation antigen (FAP) to lyse FAP-expressing targets in the presence of human complement or human mononuclear leukocytes, respectively.

[0111] In particular, the ability of L_AH_C to mediate cytotoxic effects against HT-1080FAP clone 33 cells, which expressed human FAP on the surface, was studied. Cytotoxicity was determined in vitro using the following approach: ^{51}Cr -labelled target cells were incubated in the presence of L_AH_C with human serum as source of complement or human MNC (peripheral blood mononuclear cells) as effectors. Release of ^{51}Cr was measured as measure of target-cell lysis.

[0112] Antibodies and cell lines used were L_AH_C (reshaped human anti-human FAP IgG1 antibody), hIgG1 (human IgG1 isotype control), 3S193 (murine monoclonal anti-Lewis^y IgG3 antibody), mIgG (murine IgG control), HT-1080 (human fibrosarcoma), HT-1080FAP clone 33, (HT1080 transfected with cDNA encoding human FAP), MCF-7 (human breast adenocarcinoma cell line).

Complement-mediated lysis of target cells by L_AH_C

[0113] Tumour cells were radiolabelled by incubation in RPMI1640 medium with 100 μ l ^{51}Cr (NEN) at 37° C for one hour. Subsequently, cells were washed twice in ^{51}Cr -free medium and resuspended at a concentration of 2×10^5 cells per mL.

[0114] Human serum as source of complement was freshly prepared from blood of different volunteers. Blood was taken by puncturing the arm vein, remained at room temperature for one hour to allow clotting to occur, and was kept at 4° C over night. Serum was separated by centrifugation and taken off from the sediment.

[0115] The antibody under study was diluted from the stock solution to the appropriate concentration in RPMI1640 cell culture medium.

[0116] 1×10^4 radiolabelled tumour cells of the indicated cell line were incubated in the presence of different concentrations of test or control antibody and 25% of the human serum used as source of complement for 2 h at 37° C in a 95% air and 5% CO_2 incubator. Incubation was performed in U-shaped 96-well plates in a total volume of 200 μ l RPMI1640 and done in triplicate. After the incubation period, plates were centrifugated, 100 μ l of the supernatant were taken off and radioactivity was determined in a gamma-counter. Total number of incorporated radioactivity was determined by measuring 10^4 target cells. Spontaneous release was defined as activity released from the target cells in the absence of both antibody and complement during the described incubation period.

[0117] Specific lysis was calculated as follows:

$$\text{specific lysis (in \%)} = \frac{[\text{activity sample}] - [\text{activity spontaneous release}]}{[\text{maximum activity}] - [\text{activity spontaneous release}]} \times 100$$

Antibody-dependent cellular cytotoxicity (ADCC) of L_AH_C

[0118] Tumour cells were radiolabelled by incubation in RPMI1640 medium with 100 μ l ^{51}Cr at 37°C for one hour. Subsequently, cells were washed twice in ^{51}Cr -free medium and resuspended at a concentration of 2×10^5 cells per mL.

[0119] MNC (peripheral blood mononuclear cells) were prepared from peripheral blood taken by puncturing the arm vein of different healthy human volunteers. Clotting was prevented by the addition of 20% citrate buffer. MNC from 4 mL of this blood preparation were purified by centrifugation (30 min at 400 G and room temperature) on 3 mL of lymphocyte preparation medium (Boehringer Mannheim, Germany). MNC (peripheral blood mononuclear cells) were taken off from the gradient, washed three times and diluted with RPMI1640 to the appropriate concentration. Lymphocyte activated killer (LAK) cells were derived from MNC (peripheral blood mononuclear cells) by incubation for 5 days at 37° C in a 95% air and 5% CO_2 incubator at an initial density of 1.3×10^6 cells per mL in the presence of 100U recombinant human Interleukin-2 (IL-2). The antibody under study was diluted from the stock solution to the appropriate concentration in RPMI1640 cell culture medium.

[0120] 1×10^4 radiolabelled tumour cells of the indicated cell line were incubated for 5 h at 37°C and 5% CO_2 in the presence of different concentrations of test or control antibody and MNC (peripheral blood mononuclear cells) in a number necessary to reach the indicated effector:target cell ratio. Incubation was performed in U-shaped 96-well plates in a total volume of 200 μ l RPMI1640 and done in duplicate.

[0121] After the incubation period, plates were centrifugated, 100 μ l of the supernatant were taken off and radioactivity was determined in a gamma-counter. Total number of incorporated radioactivity was determined by measuring 10^4

target cells. Spontaneous release was defined as activity released from the target cells in the absence of both antibody and effector cells during the described incubation period.

[0122] Specific lysis was calculated as follows:

$$\text{specific lysis (in \%)} = \frac{[\text{activity sample}] - [\text{activity spontaneous release}]}{[\text{maximum activity}] - [\text{activity spontaneous release}]} \times 100$$

Antibody mediated complement lysis of tumour cells

[0123] No complement mediated lysis above control was seen in HT-1080FAP clone 33 cells with L_AH_C up to a concentration of 50 $\mu\text{g/mL}$ (Table 14, Table 15a)

[0124] Lytic activity of human serum used as source of complement was shown by lysis of MCF-7 human breast carcinoma cells in the presence of 12.5 $\mu\text{g/mL}$ 3S193, a murine monoclonal anti-Lewis^y antibody with known complement activating ability (Table 15b)

Antibody mediated cellular lysis of tumour cells

[0125] In the presence of L_AH_C in a concentration of up to 10 $\mu\text{g/mL}$, no lysis of HT-1080FAP clone 33 above isotype control was detectable in ADCC mediated by human MNC (peripheral blood mononuclear cells, Table 16) or human LAK cells (lymphokine activated killer cell) (Table 17) at an effector:target ratio of 50:1:

[0126] In appropriate in vitro assays with either human complement or with human MNC (peripheral blood mononuclear cells) as effector mechanisms, human anti-FAP monoclonal antibody L_AH_C revealed no relevant cytotoxic effect above controls on FAP expressing tumor cell line HT-1080FAP clone 33.

[0127] In vitro, L_AH_C is unable to mediate cytotoxicity effected by human complement or human MNC (peripheral blood mononuclear cells) on a cell line positive for FAP, the antigen recognized by this antibody.

Table 14

Specific complement lysis (in %) of HT-1080FAP clone 33 tumor cell targets mediated by L_AH_C		
Source of human serum:	HT-1080 clone 33:	
concentration of anti-body	hlgG1 isotype control	L_AH_C
A 50 $\mu\text{g/mL}$	5	4
A 10 $\mu\text{g/mL}$	5	3
B 50 $\mu\text{g/mL}$	7	5
B 10 $\mu\text{g/mL}$	6	5
0 $\mu\text{g/mL}$	0	0
Incubation: 2 hours at 37°C, 25% serum from human volunteers A or B, respectively, as source of complement.		

Table 15a

Specific complement lysis (in %) of HT-1080FAP clone 33 tumor cell targets mediated by human anti-FAP monoclonal antibody L _A H _C		
Source of human serum:	HT1080clone 33:	
concentration of anti-body	hIgG1	L _A H _C
A 10.00 µg/ml	2	1
A 2.50 µg/ml	2	2
A 0.60 µg/ml	1	1
A 0.15 µg/ml	1	2
A 0.00 µg/ml	2	2
B 10.00 µg/ml	2	2
B 2.50 µg/ml	2	2
B 0.60 µg/ml	2	2
B 0.15 µg/ml	2	2
B 0.00 µg/ml	2	2
C 10.00 µg/ml	2	2
C 2.50 µg/ml	1	1
C 0.60 µg/ml	1	1
C 0.15 µg/ml	2	1
C 0.00 µg/ml	3	3
Incubation: 2 hours at 37°C, 25% serum from human volunteers A, B or C, respectively, as source of complement.		

Table 15b

Specific complement lysis (in %) of MCF-7 tumour cell targets mediated by murine anti-Lewis ^y monoclonal antibody 3S193		
Source of human serum:	MCF-7:	
concentration of anti-body	mIgG	3S193
A 10.00 µg/ml	0	21
A 2.50 µg/ml	1	21
A 0.60 µg/ml	0	21
A 0.15 µg/ml	1	18
A 0.00 µg/ml	0	0
B 10.00 µg/ml	1	13
B 2.50 µg/ml	0	17

Table 15b (continued)

Specific complement lysis (in %) of MCF-7 tumour cell targets mediated by murine anti-Lewis ^y monoclonal antibody 3S193		
Source of human serum:	MCF-7:	
concentration of anti-body	mIgG	3S193
B 0.60 µg/ml	1	18
B 0.15 µg/ml	1	15
B 0.00 µg/ml	0	0
C 10.00 µg/ml	1	22
C 2.50 µg/ml	0	23
C 0.60 µg/ml	1	26
C 0.15 µg/ml	1	20
C 0.00 µg/ml	1	1
Incubation: 2 hours at 37° C, 25% serum from human volunteers A, B or C, as source of complement.		

Table 16

ADCC (antibody-dependant cellular cytotoxicity) (specific lysis in %) of HT-1080FAP clone 33 target cells by human MNC (peripheral blood mononuclear cells) mediated by L _A H _C .		
HT-1080FAP clone 33:		
Concentration of anti-body:	HT-1080FAP clone 33:	
[in µg/mL]	hIgG1	L _A H _C
10.000	2	2
2.500	2	2
0.625	2	2
0.156	3	3
0.000	3	3
Incubation: 5 hours at 37°C, 10 ⁴ target cells and an effector:target cell ration of 50:1.		

Table 17

ADCC (antibody-dependent cellular cytotoxicity, specific lysis in %) of HT-1080FAP clone 33 target cells by LAK cells (lymphokine activated killer cells) mediated by L _A H _C .		
Concentration of anti-body:	HT-1080FAP clone 33:	
[in µg/mL]	hIgG1	L _A H _C
10.000	12	14
2.500	14	17
0.625	14	21
0.156	15	21
0.000	14	14
Incubation: 5 hours at 37°C, 10 ⁴ target cells and an effector:target cell ratio of 50:1.		

Example 8: Immunohistochemical analysis of monoclonal antibody L_AH_C binding to normal and neoplastic human tissues

[0128] This experiment was performed to determine the binding characteristics of the humanized mAb L_AH_C to normal and neoplastic human tissues.

[0129] The following antibodies were used: L_AH_C, cF19, and the negative control hu IgG1 were directly biotinylated according to methods of the state of the art and used at concentrations of 2.5 to 0.25 mg/ ml in 2% BSA/PBS (bovine serum albumin in phosphate-buffered saline). Murine mAb F19 was used as tissue culture supernatant of the F19 hybridoma, at dilutions of 1:5 to 1:10 in 2% BSA/PBS.

[0130] The following reagents were used for immunochemical assays: Streptavidin peroxidase complex (Vector Labs., Burlingame, CA, USA), Avidin-biotin peroxidase complex (Vector Labs.), Biotinylated horse anti-mouse (Vector Labs.), DAB (diaminobenzidine, Sigma Chemical Co. St. Louis, MO, USA), Harris' hematoxylin.

[0131] Fresh frozen tissue samples examined included the following: Normal colon, breast, lung, stomach, pancreas, skin, larynx, urinary bladder, smooth and skeletal muscle.

[0132] Among the tumors tested were carcinomas from breast, colon, lung, esophagus, uterus, ovary, pancreas, stomach, and head and neck.

[0133] An indirect immunoperoxidase method was carried out according to state of the art methods (Garin-Chesa P, Old LJ, Rettig WJ: Cell surface glycoprotein of reactive stromal fibroblasts as a potential antibody target in human epithelial cancers. Proc Natl Acad Sci USA 1990; 87:7235-7239) on five micrometer thickness fresh frozen sections.

[0134] DAB was used as a substrate for the final reaction product. The sections were counterstained with Harris' hematoxylin and examined for antigen expression.

L_AH_C expression in normal human tissues

[0135] The normal tissues tested were negative for L_AH_C expression, except for the normal pancreas in which a subset of positive endocrine cells in the islets of Langerhans (A cells) were identified with L_AH_C, cF19 and F19. (Table 18). No immunoreactivity was observed with the hu IgG1 (human immunoglobulin IgG1 subclass) used as a negative control.

L_AH_C expression in tumors

[0136] In the tumor samples, L_AH_C, cF19 and F19 showed an indistinguishable pattern of expression in the tumor stromal fibroblasts. A strong and homogeneous expression was found in the majority of the cases examined, especially in the cancer samples derived from breast, colon, lung, pancreas and in the squamous cell carcinomas (SQCC) of the head and neck tested (Table 19). No immunoreactivity was observed with the hu IgG1 used as negative control.

[0137] L_AH_C, cF19 and F19 showed immunoreactivity with the tumor stromal fibroblasts in the epithelial cancer samples tested. No L_AH_C or F19 immuno-reactivity was seen with either the fibrocytes of the normal organ mesenchyme or

the parenchymal cells of normal adult organs. The only exception was a subset of endocrine cells in the pancreatic islets, presumably glucagon-producing A cells, which react with the anti-FAP antibodies.

[0138] Immunohistochemical analysis of L_AH_C in normal human tissues and FAP-expressing human carcinomas showed indistinguishable patterns of binding for L_AH_C, cF19 and murine mAb F19.

Table 18

Immunoreactivity of mAbs L _A H _C , cF19 and F19 with normal human tissues				
Tissue type		L _A H _C	cF19	F19
Breast	-Duct epithelium	-	-	-
	-Myoepithelial cells	-	-	-
Colon	-Glandular epithelium	-	-	-
	-Smooth muscle	-	-	-
Lung	-Bronchial epithelium	-	-	-
	-Alveolar epithelium	-	-	-
Stomach	-Glandular epithelium	-	-	-
	-Smooth muscle	-	-	-
Urinary bladder	-Urothelium	-	-	-
	-Smooth muscle	-	-	-
Pancreas	-Exocrine acini	-	-	-
	-Endocrine islet cells	+ subset only	+subset only	+ subset only
Larynx -Squamous epithelium		-	-	-
Lymph node -Lymphocytes		-	-	-
Skeletal muscle-		-	-	-
Connective tissue		-	-	-
Skin	-Keratinocytes	-	-	-
	-Sweat glands	-	-	-

Table 19

Immunoreactivity of mAbs L _A H _C , cF19 and F19 with human tumor samples				
Tumor type	No.	L _A H _C	cF19	F19
Breast cancers (infiltrating ductal type)	7	7 Positive	7 Positive	7 Positive
Colon cancers (adenocarcinomas)	7	7 Positive	7 Positive	7 Positive
Lung carcinomas (adenocarcinoma (2) large cell type (2) squamous type (4)	8	7 Positive	7 Positive	7 Positive
		1 Negative	1 Negative	1 Negative
Esophageal cancers (squamous type)	1	1 Positive	1 Positive	1 Positive
Endometrial cancers (adenocarcinoma)	1	1 Negative	1 Negative	1 Negative
Gastric cancers (adenocarcinoma)	2	2 Negative	2 Negative	2 Negative
Ovarian cancers (serous denocarcinoma)	2	1 Positive	1 Positive	1 Positive
		1 Negative	1 Negative	1 Negative

Table 19 (continued)

Immunoreactivity of mAbs L _A H _C , cF19 and F19 with human tumor samples				
Tumor type	No.	L _A H _C	cF19	F19
Pancreatic cancers (adenocarcinomas)	2	2 Positive	2 Positive	2 Positive
Head and neck cancers (squamous cell type)	4	4 Positive	4 Positive	4 Positive
Abbreviations: No, number of cases from different patients studied; positive, number of cases showing antigen expression in the tumor stroma; negative, number of cases tested that lacked detectable antigen expression.				

Example 9: Species specificity of L_AH_C binding in tissue sections

[0139] This experiment was conducted to assess the reactivity of L_AH_C with tissues from mouse, rat, rabbit and cynomolgus monkeys by immunohistochemical methods.

[0140] Also used in these tests were cF19 and huIgG1 as negative controls. The reagents used for immunohistochemistry were Streptavidin peroxidase complex (Vector Labs., Burlingame, CA, USA), DAB (Sigma Chemical Co., St. Louis, MO, USA) and Harris' hematoxylin.

[0141] The following fresh frozen tissue samples from mouse, rat, rabbit and cynomolgus were tested: Brain, liver, lung, kidney, stomach, pancreas, intestine, thymus, skin, muscle, heart, spleen, ovary, uterus and testes. As positive control, sections from normal human pancreas and a breast carcinoma sample were included in every assay.

Immunohistochemistry

[0142] An indirect immunoperoxidase method was carried out as described in the state of the art (Garin-Chesa P, Old LJ, Rettig WJ: Cell surface glycoprotein of reactive stromal fibroblasts as a potential antibody target in human epithelial cancers. Proc Natl Acad Sci USA 1990; 87:7235-7239) on five micrometer thickness fresh frozen sections. The antibodies L_AH_C, cF19 and hu IgG1 (at 1 µg/ml) were biotinylated according to the state of the art and were detected with streptavidin peroxidase complex. DAB was used as a substrate for the final reaction product. The sections were counterstained with Harris' hematoxylin and examined for antigen expression.

[0143] The normal tissues tested did not react with either L_AH_C or cF19 in the experiments (Table 1).

[0144] The normal human pancreas used as positive control showed L_AH_C and cF19 binding in a subset of endocrine cells in the islets of Langerhans as previously described for F19. In addition, binding of L_AH_C and cF19 was seen in the tumor stromal fibroblasts in the breast carcinoma sample.

[0145] Immunohistochemical analysis of normal tissues from mouse, rat, rabbit and cynomolgus failed to detect any binding of either L_AH_C or cF19, in the experiments performed.

Table 20

Binding of L _A H _C to tissue sections of non-human species, as determined by immunohistochemistry.					
Organ / Tissue type		Mouse	Rat	Rabbit	Cynomolgus
Brain	-Cerebral cortex	-	-	-	
	-Cerebellum	-	-	-	-
Liver	-Hepatocytes	-	-	-	-
	-Portal triad	-	-	-	-
Lung	-Bronchi	-	-	-	-
	-Alveoli	-	-	-	-
Kidney	-Glomeruli	-	-	-	-
	-Tubular epithelium	-	-	-	-
Stomach	-Glandular epithelium				
	-Smooth muscle	-	-	-	-
Pancreas	-Exocrine acini	-	-	-	-
	-Endocrine islets	-	-	-	-
Intestine	-Glandular epithelium	-	-	-	-
	-Smooth muscle	-	-	-	-
Thymus -Lymphocytes		-	-	-	-
Skin	-Keratinocytes	-	-	-	-
	-Sweat glands	-	-	-	-
	-Hair follicles	-	-	-	-
Skeletal muscle		-	-	-	-
Heart		-	-	-	-
Spleen -Lymphocytes		-	-	-	-
Ovary	-Follicular epithelium	-	-	-	-
	-Stroma	-	-	-	-
Uterus	-Myometrium	-	-	-	-
	-Cervix uteri	-	-	-	-
Testis -Tubular epithelium		nt	nt	nt	-
Connective tissue		-	-	-	-

nt, not tested

Example 10: Construction of cell lines producing chimeric and reshaped anti-FAP monoclonal antibodies

[0146] The objective of this experiment was to demonstrate stable cell lines according to the invention expressing L_AH_C, L_AH_A, L_BH_B, L_BH_D, and cF19 in CHO DG44 cells. Stable cell lines transfected with humanized or chimeric F19 antibodies were produced and their identity was confirmed by PCR amplification of heavy and light variable regions using genomic DANN derived from each transfectant as template.

[0147] CHO DG44 cells maintained under serum-free conditions in SFM-II medium. Lipofectin and SFM-II serum-free medium were obtained from Gibco/BRL. Geneticin and all restriction enzymes were obtained from Boehringer Mannheim. Pfu polymerase was obtained from Stratagene.

[0148] DNA for transfections was purified from E. coli cells using QiaFilter Maxi Cartridges (Qiagen) as directed by the manufacturer. All DNA preparations were examined by restriction enzyme digestion. Sequences of L_AH_C variable regions in their respective vectors were confirmed using an ABI PRISM 310 Sequencer.

[0149] Further information regarding the vectors and DNA sequences employed is available in the prior examples.

Transfection of CHO DG44 cells

[0150] Cells in logarithmic growth were plated into 6 well plates containing 1 mL fresh SFM-II medium. Plasmids encoding heavy and light chains of humanized or chimeric F19 versions were cotransfected into CHO DG44 cells using liposomal transfection. Liposomes were prepared using 6 μ l Lipofectin reagent and 0.5 μ g of each vector (one for the desired heavy chain and one for the light) as described for LipofectAMINE transfections except that SFM-II medium was used to dilute all reagents. Twenty-four hours later, cells were diluted 1:10 into SFM-II medium containing 300 μ g/mL Geneticin. After the initial phase of cell killing was over (10-14 days), the concentration of Geneticin was reduced to 200 mg/mL and methotrexate was added to a final concentration of 5 nM. Methotrexate concentrations were increased after 10-14 days to a final concentration of 20 nM.

PCR Amplification of transfectant DNA

[0151] 10^7 CHO DG44 cells were centrifuged in an Eppendorf microcentrifuge briefly at full speed, washed once with PBS, and pelleted once again. Genomic DNA was prepared by ethanol precipitation after SDS lysis and Proteinase K treatment of the cell pellets.

[0152] A mixture containing one of the following primer pairs, dNTPs, buffer, and Pfu polymerase was used to amplify either the heavy or light chain variable region using genomic DNA as template. The resulting PCR products were digested with the appropriate restriction enzyme and analyzed by agarose gel electrophoresis to confirm their identity.

Light chain primer set:

[0153]

5'-GAG ACA TTG TGA CCC AAT CTC C - 3' PKN 1690

5'- GAC AGT CAT AAA CTG CCA CAT CTT C - 3' PKN.1930.R

Heavy chain primer set:

[0154]

5'-TTG ACA CGC GTC TCG GGA AGC TT - 3' PG 5863

5'- GGC GCA GAG GAT CCA CTC ACC T - 3' PG 6332.R

[0155] The undigested heavy chain PCR product has a predicted size of 469 bp while the light chain PCR product has a predicted size of 286 bp. Verification of identity was determined by restriction enzyme digest with BstEII (heavy chain) or NlaIV (light chain).

[0156] CHO cell lines were transfected with L_AH_C , L_AH_A , L_BH_B , L_BH_D , as well as cF19. Geneticin-resistant cells were obtained and these cells were further selected for resistance to methotrexate. PCR amplification of the light and heavy chain DNA produced the expected bands and confirmed the identity of L_AH_C , L_AH_A and L_BH_D transfectants. The L_AH_C full length heavy chain PCR product was subcloned and resequenced in its entirety.

[0157] The cells described were maintained under serum-free conditions at all times and were not treated with animal-derived products such as trypsin.

[0158] Producer cell lines transfected with expressing monoclonal L_AH_C , L_AH_A , L_BH_B , L_BH_D and cF19 antibodies were produced. Their identities were confirmed using PCR amplification of both their heavy and light chain variable regions. The DNA sequence of the heavy chain variable region PCR products for L_AH_C -transfected cells was confirmed.

Example 11: Expression of antibody proteins in Chinese hamster ovary DG 44 cells and their purification

[0159] The objective of this experiment was to express and purify of L_AH_C , L_AH_A , L_BH_B , and L_BH_D mAbs to enable their characterization. Other goals included the establishment of a quantitative ELISA to permit measurement of anti-

body concentrations in both crude media samples as well as purified Ig samples and determination of relative expression levels of various humanized F19 constructs using this assay.

[0160] Serum-free CHO DG44 cells and USP-grade methotrexate were obtained from the Biotechnical Production Unit of the Dr. Karl Thomae GmbH, Biberach, Germany; both products are also commercially available. Cells were maintained under serum-free conditions at all times. SFM-II serum-free medium was obtained from Gibco/BRL.

[0161] Protein A agarose was from Pierce Chemical (Indianapolis, IN, USA). Human IgG1 standards (Cat. No. I 3889), p-Nitrophenyl phosphate tablets (N 2640), bovine serum albumin (BSA) (A 7906), and goat anti-human kappa chain specific alkaline phosphatase-conjugated antibody (A 3813) were obtained from Sigma Chemical (St. Louis, MO, USA). Goat anti-human gamma-chain specific alkaline phosphatase-conjugated antibody was obtained from Jackson Immunoresearch Laboratories (through Stratech Scientific). Tris-buffered saline (TBS) consisted of 150 mM NaCl, 50 mM Tris, pH 7.5.

Cell culture conditions for antibody expression

[0162] Cells were cultured and L_AH_C -producing cells were maintained in T-175 flasks in SFM-II serum-free medium without agitation. The medium contained 200 μ g/mL Geneticin and 20 nM methotrexate without antibiotics. Cells were passaged by dilution, were not adherent, and grew in small clusters. When the cells reached stationary phase, the medium was collected and centrifuged to remove cells and frozen at -20°C until needed.

Purification of L_AH_C

[0163] All purification steps were carried out at 4° C. A C10/10 column (Pharmacia Fine Chemicals) was packed with Protein A agarose (3 mL bed volume). The column was washed with TBS and preeluted once with 0.1 M Na citrate, pH 3.0 to insure that no loosely bound material remained on the column. The column was then immediately reequilibrated with TBS and stored at 4°C. Spent culture supernatants were thawed and centrifuged at 10,000 xg for 30 minutes prior to Protein A chromatography to remove debris and diluted with an equal volume of TBS. This material was loaded onto the Protein A column at 0.5 mL/min using a P-1 peristaltic pump (Pharmacia) and washed with TBS until the absorbance at 280 nm was undetectable. Elution of the antibody was initiated with 0.1 M Na citrate pH 3.0 at approximately 0.2 mL/min. The elution was monitored at 280 nm and one mL fractions of the eluted material were collected into tubes containing sufficient Tris base pH 9 to neutralize the citrate buffer. Protein-containing fractions were pooled and concentrated using an Amicon filtration apparatus with a YM-30 filter and dialyzed against PBS. The column was immediately regenerated with TBS. Protein dye-binding assays were performed with the BioRad (Hercules, California) protein determination kit, according to the manufacturer's instructions, using bovine serum albumin as a standard.

Human IgG (gamma immunoglobulin) ELISA

[0164] ELISA plates were coated overnight with 100 μ L of goat anti-human gamma-chain specific alkaline phosphatase-conjugated antibody at 0.4 mg/mL in coating buffer at 4°C. Coating antibody was removed and plates were blocked with 2% BSA in PBS for 2 hours. All subsequent steps were performed at 37°C. Blocking buffer was replaced with antibody samples or human IgG1 standard diluted in dilution buffer, serially diluted in a 200mL volume, and incubated for one hour. Negative controls included dilution buffer and/or culture medium of nontransfected cells. Wells were washed and 100 μ L of goat anti-human kappa chain specific alkaline phosphatase-conjugated antibody diluted 1:5000 was added and incubated for one hour. Wells were washed and 100 μ L reaction buffer was added and incubated for 30 minutes. The reaction was stopped by addition of 1 M NaOH and absorbance read at 405 nm in an ELISA plate reader. Results were analyzed by four-parameter iterative curve fitting.

[0165] Amino acid analysis was performed according to methods available in the state of the art.

[0166] Monoclonal antibody L_AH_C was produced and purified to homogeneity using Protein A affinity chromatography. ELISA assays using human IgG1 as standard indicated L_AH_C recoveries exceeding 70%. The purity of the material was estimated to be >90% by SDS-polyacrylamide gel electrophoresis. Representative expression data and typical purification yields are shown in Table 21.

Table 21

Expression data and purification yields FAP antibody proteins in CHO cells			
Antibody	Expression levels in crude media samples (ELISA)	Purified antibody yields	Yield improvement [purified antibody]
H _C L _A	7 - 10 mg/L	~ 5 - 7 mg/L	500 - 700
H _A L _A	5 - 7 mg/mL	~ 3 - 4 mg/L	300 - 400
H _B L _B	0.5 - 1 mg/mL	~ 0.2 - 0.5 mg/L	20 - 50
H _D L _B	0.8 - 1.5 mg/mL	~ 0.3 - 0.8 mg/L	30 - 60
Chimeric F19	~ 0.02 mg/mL	< 0.01 mg/L	1
Representative expression data for each of the anti-FAP antibodies produced in this study are shown. Recoveries after Protein A agarose affinity chromatography were based on protein dye-binding measurements of the purified Ig using BSA as a standard.			

Example 12: Binding of monoclonal antibody L_AH_C to isolated recombinant human FAP

[0167] The objective of this study was to characterize binding of L_AH_C to isolated recombinant human FAP.

CD8-FAP ELISA

[0168] ELISA plates were coated overnight with 100 µL of mouse anti-rat antibody (Sigma Chemical R0761) at 1:2000 in coating buffer at 4 °C. Coating antibody was removed and plates were blocked with 2% BSA in PBS for one hour. All subsequent steps were performed at room temperature. Blocking buffer was replaced with 100 µL of 1 µg/mL rat anti-CD8 antibody (Pharmingen 01041D) and incubated for one hour. Plates were washed and 100 µL CD8-FAP culture supernatant (1:2 in PBS) was added and allowed to bind for one hour. Plates were washed and antibody samples were added (two-fold serial dilutions) in a 100 µL volume and incubated for one hour. Negative controls included human IgG and/or culture medium of nontransfected cells. Wells were washed and 100 µL of horse radish peroxidase (HRP) conjugated mouse anti-human IgG1 antibody (Zymed 05-3320) diluted 1:500 in dilution buffer were added and incubated for one hour. Wells were washed and 100 µL HRP substrate, (azino-bis (3-ethylbenzthiazoline 6-sulfonic) acid, Sigma Chemical A9941), were added and incubated for 60 minutes. The reaction was stopped by addition of 1 M NaOH and absorbance read at 405/490 nm in an ELISA plate reader. Results were analyzed by four parameter curve iterative curve fitting.

[0169] Alternatively, plates were coated directly with cF19. FAP (recombinant human FAP) was allowed to bind to these plates as above and biotinylated L_AH_C (~1 µg/mL) was then added. Antibody binding was detected with HRP-streptavidin conjugate as above.

Solubilization of membrane-bound human FAP

[0170] FAP-expressing 293FAP 1/2 cells or control 293 cells were washed with PBS and lysed with 1% Triton X-114 in Tris-buffered saline. Nuclei and debris were removed by centrifugation at 10,000 xg. The supernatant was phase-partitioned (Estreicher A, Wohlend A, Belin D, Scheuning WD, Vasalli JD. Characterization of the cellular binding site for the urokinase-type plasminogen activator. J Biol Chem 1989; 264:1180-1189) to enrich membrane proteins. The detergent phase was collected and diluted in buffer containing 1% Empigen BB (Calbiochem) to prevent reaggregation of the Triton X-114.

[0171] This material was subjected to Concanavalin A agarose chromatography (Rettig WJ, Garin-Chesa P, Healey JH, Su SL, Ozer HL, Schwab, M, Albino AP, Old LJ. Regulation and heteromeric structure of the fibroblast activation protein in normal and transformed cells of mesenchymal and neuroectodermal origin. Cancer Res 1993; 53:3327-3335).

Biotinylation of L_AH_C

[0172] L_AH_C (1-2 mg) was dialyzed against 50mM bicarbonate buffer and biotinylated with a ten-fold molar excess of

sulfosuccinimidyl-6-biotinamido hexanoate (NHS-LC biotin, Pierce Chemical, Rockford, Illinois, USA) for 2 hours at room temperature. Unreacted product was removed by repeated microdialysis in a microconcentrator.

Transient transfections

[0173] COS-7 cells (American Type Tissue Culture Collection, reference number CRL 1651) were cotransfected by electroporation with the heavy and light chain vectors encoding L_AH_C.

[0174] Anti-CD8 monoclonal antibody was immobilized onto microtiter plates. CD8-FAP from medium of insect cells infected with CD8-FAP baculovirus was allowed to bind to these plates. Spent medium from COS-7 cell cultures transiently transfected with two separate vectors encoding L_AH_C was serially diluted and added to the wells containing the immobilized CD8-FAP. L_AH_C bound to isolated immobilized CD8-FAP protein (Figure 35). Culture supernatants from mock-transfected COS-7 cells failed to demonstrate binding.

[0175] Recombinant membrane-bound FAP from detergent extracts of 293FAP 1/2 cells or control extracts was serially diluted and immobilized via chimeric F19 monoclonal antibody bound to microtiter plates. Biotinylated L_AH_C bound recombinant human FAP immobilized with cF19 (Figure 36) in a concentration-dependent manner.

[0176] L_AH_C recognized isolated immobilized recombinant human FAP carrying the epitope for murine F19. L_AH_C bound to both CD8-FAP produced in insect cells, as well as FAP protein produced in 293FAP 1/2 cells.

[0177] Culture supernatants from COS7 cells transfected with either heavy and light chain vectors encoding L_AH_C or without DNA (Control) were collected three days posttransfection. CD8-FAP was immobilized via an anti-CD8 antibody as described in the text. Serial dilutions of the COS7 supernatants were allowed to bind to the immobilized CD8-FAP and subsequently detected with an HRP-conjugated anti-human IgG1 antibody.

[0178] Detergent extracts of FAP-expressing 293FAP 1/2 cells or control 293 cells were serially diluted and added to cF19-coated microtiter plates. Biotinylated L_AH_C was added and binding of biotinylated L_AH_C was detected with HRP-conjugated streptavidin.

Example 13: Characterization of HT-1080 fibrosarcoma cells and 293 human embryonic kidney cells transfected with cDNA for human FAP

[0179] Fibroblast activation protein (FAP) is a cell-surface, membrane-bound protein which carries the F19 epitope and is expressed on tumor stromal fibroblasts. Cell lines expressing recombinant FAP protein and matched controls lacking FAP were generated for the characterization of anti-FAP monoclonal antibodies.

[0180] Cells used were HT-1080 cells (reference number CCL 121) and 293 human embryonic kidney cells (reference number CRL 1573) were obtained from the American Type Culture Collection (Maryland, USA). Transfectam was obtained from Promega. Geneticin and all restriction enzymes were obtained from Boehringer Mannheim. DNA for transfections was purified from E. coli cells using QiaFilter Maxi Cartridges (Qiagen) as directed by the manufacturer. All DNA preparations were examined by restriction enzyme digestion. Vector sequences were confirmed using an ABI PRISM 310 Sequencer.

[0181] Further information regarding the vectors and DNA sequences employed has been described in Scanlan MJ, Raj BK, Calvo B, Garin-Chesa P, Sanz-Moncasi MP, Healey JH, Old LJ, Rettig WJ. Molecular cloning of fibroblast activation protein alpha, a member of the serine protease family selectively expressed in stromal fibroblasts of epithelial cancers. *Proc Natl Acad Sci USA* 1992; 89:10832-10836. The FAP cDNA sequence has been deposited in Genbank (accession number HS09287).

Cell culture and immunoassays

[0182] HT-1080 cells were transfected with 1 mg DNA using Transfectam according to the manufacturer's instructions. Human embryonic kidney 293 cells were transfected by calcium phosphate transfection (Brann MR; Buckley NJ; Jones SVP; Bonner TI).

[0183] Expression of cloned muscarinic receptor in A9 L cells. *Mol Pharmacol* 1987; 32:450-455) with 10 mg DNA. Twenty-four hours later, cells were diluted 1:10 into fresh medium containing 200 mg/mL Geneticin. Colonies were picked and examined by immunofluorescence for FAP expression as described in Rettig WJ; Garin-Chesa P; Beresford HR; Oettgen HF; Melamed MR; Old LJ. Cell-surface glycoproteins of human sarcomas: differential expression in normal and malignant tissues and cultured cells. *Proc Natl Acad Sci USA* 1988; 85:3110-3114.

[0184] Immunoprecipitations with cF19 were performed with metabolically labelled cells as described in Rettig WJ, Garin-Chesa P, Healey JH, Su SL, Ozer HL, Schwab, M, Albino AP, Old LJ. Regulation and heteromeric structure of the fibroblast activation protein in normal and transformed cells of mesenchymal and neuroectodermal origin. *Cancer Res* 1993; 53:3327-3335.

[0185] HT-1080 and 293 cells were tested for FAP antigen expression in immunofluorescence assays with anti-FAP

antibodies and were found to be antigen-negative. Transfection of these cells with FAP:38 vector resulted in the generation of Geneticin-resistant colonies. Isolated colonies were picked and analyzed by immunofluorescence for FAP expression. Two cell clones were identified, designated HT-1080FAP clone 33 and 293FAP I/2, which express cell surface-bound FAP protein, as recognized by cF19 antibody. Staining of nonpermeabilized HT-1080FAP clone 33 cells and 293FAP I/2 with cF19 antibody confirmed the cell surface localization of the FAP protein.

[0186] Immunoprecipitation of radiolabelled FAP protein with cF19 from extracts of ³⁵S-methionine labelled HT-1080FAP clone 33 cells or 293FAP I/2 cells resulted in the appearance of a 93 kilodalton band after autoradiography. This band is absent in immunoprecipitates of parental HT-1080 or 293 cell extracts.

[0187] Two stably transfected cell lines, HT-1080FAP clone 33 and 293FAP I/2, express FAP on the cell surface as determined in immunological assays with anti-FAP mAbs. Neither parental HT-1080 cells nor parental 293 cells express detectable levels of FAP.

Example 14: Generation and characterization of CD8-FAP fusion protein

[0188] A soluble form of human FAP (fibroblast activation protein) in the form of a CD8-FAP fusion protein was produced in insect cells for the characterization of L_AH_C containing the binding site for anti-FAP mAbs. Murine CD8 was chosen to permit secretion of the protein and to provide an additional epitope tag.

[0189] The cDNA encoding the extracellular domain of CD8, consisting of the first 189 amino acids of murine CD8, was linked to that of the extracellular domain of FAP (amino acids 27 to 760), essentially as described by Lane, et al. (Lane P, Brocker T, Hubele S, Padovan E, Lazavacchia A, McConnell. Soluble CD40 ligand can replace the normal T cell-derived CD40 ligand signal to B cells in T cell-dependent activation. J Exp Med 1993, 177:1209-1213) using standard PCR protocols. The authenticity of all clones was verified by DNA sequencing. The resulting DNA was inserted into the pVL1393 vector (Invitrogen) and transfection of Sf9 cells (Invitrogen) with this vector and amplification of the resulting recombinant baculovirus were performed as described (Baculovirus Expression Vectors. A Laboratory Manual. O'Reilly DR, Miller LK, Luckow VA, (Eds.), Oxford University Press: New York, 1994). The spent medium of High Five™ cells (Invitrogen) infected with recombinant CD8-FAP baculovirus for four days was collected and cleared by ultracentrifugation.

[0190] The CD8-FAP ELISA (enzyme-linked immunosorbent assay) has been described above (Example 12).

[0191] Insect cell cultures infected with CD8-FAP virus secreted a fusion protein into the medium which carries the F19 epitope and is recognized by an anti-FAP antibody (Figure 1). Neither the cell culture medium alone nor medium from insect cells infected with CD8-CD40L fusion protein bound anti-FAP antibody.

[0192] Soluble CD8-FAP protein carrying the F19 epitope was secreted into the medium of infected insected cell cultures. Culture supernatant from cells infected with a control construct did not contain antigen bearing the F19 epitope.

[0193] A soluble form of FAP, CD8-FAP, was produced in insect cells and CD8-FAP was shown to carry the epitope recognized by cF19.

[0194] Supernatants from insect cells infected with recombinant baculovirus encoding either CD8-FAP or CD8-CD40L fusion protein were collected four days postinfection. Cell culture medium without cells was used as an additional control (medium). Serial dilutions of these materials were added to anti-CD8 antibody-coated microtiter plates and allowed to bind. cF19 (1 mg/mL) was subsequently added and allowed to bind.

[0195] Bound cF19 was detected with horseradish peroxidase-conjugated anti-human antibody.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Boehringer Ingelheim International GmbH
 (B) STREET: Rheinstrasse
 (C) CITY: Ingelheim am Rhein
 (E) COUNTRY: Germany
 (F) POSTAL CODE (ZIP): 55216
 (G) TELEPHONE: ++49-6132-772770
 (H) TELEFAX: ++49-6132-774377

(ii) TITLE OF INVENTION: FAP alpha-specific antibody with improved producibility

(iii) NUMBER OF SEQUENCES: 101

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 339 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GACATTGTGA TGACCCAATC TCCAGACTCT TTGGCTGTGT CTCTAGGGGA GAGGGCCACC	60
ATCAACTGCA AGTCCAGTCA GAGCCTTTTA TATTCTAGAA ATCAAAAGAA CTA CTGAGGCC	120
TGGTATCAGC AGAAACCAGG ACAGCCACCC AAACCTCTCA TCTTTTGGGC TAGCACTAGG	180
GAATCTGGGG TACCTGATAG GTTCAGTGGC AGTGGGTTTG GGACAGACTT CACCCTCACC	240
ATTAGCAGCC TGCAGGCTGA AGATGTGGCA GTTTATTACT GTCAGCAATA TTTTAGCTAT	300
CCGCTCACGT TCGGACAAGG GACCAAGGTG GAAATAAAA	339

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu	Gly
1				5					10					15	

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 5 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 10 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 15 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 Lys

(2) INFORMATION FOR SEQ ID NO: 3:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 339 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GACATTGTGA TGACCCAATC TCCAGACTCT TTGGCTGTGT CTCTAGGGGA GAGGGCCACC 60
 30 ATCAACTGCA AGTCCAGTCA GAGCCTTTTA TATTCTAGAA ATCAAAAGAA CTACTTGGCC 120
 TGGTTCCAGC AGAAACCAGG ACAGCCACCC AAACCTCTCA TCTTTTGGGC TAGCACTAGG 180
 GAATCTGGGG TACCTGATAG GTTCAGTGGC AGTGGGTTTG GGACAGACTT CACCCTCACC 240
 35 ATTAGCAGCC TGCAGGCTGA AGATGTGGCA GTTTATGACT GTCAACAATA TTTTAGCTAT 300
 CCGCTCACGT TCGGACAAGG GACCAAGGTG GAAATAAAA 339

(2) INFORMATION FOR SEQ ID NO: 4:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 45 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

50 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Arg Asn Gln Lys Asn Tyr Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu Ser Gly Val
 5 50 55 60
 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Asp Cys Gln Gln
 85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 Lys

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 339 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GACATTGTGA TGACCCAATC TCCAGACTCT TTGGCTGTGT CTCTAGGGGA GAGGGCCACC 60
 ATCAACTGCA AGTCCAGTCA GAGCCTTTTA TATTCTAGAA ATCAAAAGAA CTACTTGGCC 120
 TGGTATCAGC AGAAACCAGG ACAGCCACCC AAACCTCCTCA TCTATTGGGC TAGCACTAGG 180
 GAATCTGGGG TACCTGATAG GTTCAGTGGC AGTGGGTTTG GGACAGACTT CACCCTCACC 240
 ATTAGCAGCC TGCAGGCTGA AGATGTGGCA GTTTATTACT GTCAGCAATA TTTTAGCTAT 300
 CCGCTCACGT TCGGACAAGG GACCAAGGTG GAAATAAAA 339

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val

50 55 60
 5 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 10 Lys

(2) INFORMATION FOR SEQ ID NO: 7:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 372 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CAGGTGCAAC TAGTGCAATC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
 25 AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
 CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCCTA ACAATGGTAT TCCTAACTAC 180
 AACCAGAAGT TCAAGGGCCG GGCCACCTTG ACCGTAGGCA AGTCTGCCAG CACCGCCTAC 240
 30 ATGGAAGTGT CCAGCCTGCG CTCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA 300
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCCCTGTC 360
 ACCGTCTCCT CA 372

(2) INFORMATION FOR SEQ ID NO: 8:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60

55

Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 5 85 90 95
 Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 372 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
 AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
 CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCCTA ACAATGGTAT TCCTAACTAC 180
 AACCAGAAGT TCAAGGGCCG GGCCACCTTG ACCGTAGGCA AGTCTGCCAG CACCGCCTAC 240
 ATGGAAGTGT CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTTCTGCGC CAGAAGAAGA 300
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCCCTTGT 360
 ACCGTCTCCT CA 372

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 124 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95

Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
100 105 110

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 372 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

CAGGTGCAAC TAGTGCAATC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
AGCTGTAAAA CTAAGTACATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCTTA ACAATGGTAT TCCTAACTAC 180
AACCAGAAGT TCAAGGGCCG GGTCAACATC ACCGTAGACA CCTCTGCCAG CACCGCCTAC 240
ATGGAAGTGT CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA 300
ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCCCTGTC 360
ACCGTCTCCT CA 372

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 124 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
20 25 30
Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
35 40 45
Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
50 55 60
Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CAGGTGCAAC TAGTGCAATC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
 AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
 CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCCTA ACAATGGTAT TCCTAACTAC 180
 AACCAGAAGT TCAAGGGCCG GGTCAACATC ACCGTAGACA CCTCTGCCAG CACCGCCTAC 240
 ATGGAAGTGT CCAGCCTGCG CTCGAGGAC ACTGCAGTCT ACTTCTGCGC CAGAAGAAGA 300
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCTTGTTC 360
 ACCGTCTCTCT CA 372

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala 15
 1 5 10
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr 30
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile 45
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe 60
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr 80
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Phe Cys 95
 85 90 95

Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
 AGCTGTAAAA CTAGTGGATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
 CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCCTA ACAATGGTAT TCCTAACTAC 180
 AACCAGAAGT TCAAGGGCCG GGTCAACATC ACCGTAGACA CCTCTGCCAG CACCGCCTAC 240
 ATGGAAGTGT CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA 300
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCCCTGTC 360
 ACCGTCTCCT CA 372

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 220 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15
 Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30
 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln
35 40 45
 Ser Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60
 Pro Asp Arg Phe Thr Gly Ser Gly Phe Gly Thr Asp Phe Asn Leu Thr
65 70 75 80
 Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Asp Cys Gln Gln
85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
100 105 110
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
115 120 125
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
130 135 140
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
145 150 155 160
 Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
165 170 175
 Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
180 185 190
 Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
195 200 205
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215 220

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 453 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

5 Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser
 1 5 10 15
 Val Lys Met Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr Thr
 20 25 30
 Ile His Trp Val Arg Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly
 35 40 45
 10 Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe Lys
 50 55 60
 Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ser Ser Thr Ala Tyr Met
 65 70 75 80
 15 Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala
 85 90 95
 Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp Tyr
 100 105 110
 20 Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Ser Thr Lys Gly
 115 120 125
 Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly
 130 135 140
 25 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
 145 150 155 160
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 165 170 175
 30 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
 180 185 190
 Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val
 195 200 205
 35 Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys
 210 215 220
 Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu
 225 230 235 240
 Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 245 250 255
 40 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 260 265 270
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 275 280 285
 45 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 290 295 300
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 305 310 315 320
 50 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 325 330 335
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro

55

340 345 350
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln
 355 360 365
 5 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 370 375 380
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 385 390 395 400
 10 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 405 410 415
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 420 425 430
 15 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 435 440 445
 Leu Ser Pro Gly Lys
 450

(2) INFORMATION FOR SEQ ID NO: 19:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 321 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

30 CGTACTGTGG CTGCACCATC TGTCTTCATC TTCCCGCCAT CTGATGAGCA GTTGAAATCT 60
 GGAATGCCT CTGTTGTGTG CCTGCTGAAT AACTTCTATC CCAGAGAGGC CAAAGTACAG 120
 TGGAAGGTGG ATAACGCCCT CCAATCGGGT AACTCCCAGG AGAGTGTAC AGAGCAGGAC 180
 AGCAAGGACA GCACCTACAG CCTCAGCAGC ACCCTGACGC TGAGCAAAGC AGACTACGAG 240
 35 AAACACAAAG TCTACGCTG CGAAGTCACC CATCAGGGCC TGAGCTCGCC CGTCACAAAG 300
 AGCTTCAACA GGGGAGAGTG T 321

(2) INFORMATION FOR SEQ ID NO: 20:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 107 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 45 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

50 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 1 5 10 15
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 20 25 30

55

EP 0 953 639 A1

5	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser
	50						55					60				
	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu
	65					70					75					80
10	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser
					85					90					95	
	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys					
				100					105							

(2) INFORMATION FOR SEO ID NO: 21:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 990 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

25	GCCTCCACCA	AGGGCCCCATC	GGTCTTCCCC	CTGGCACCCT	CCTCCAAGAG	CACCTCTGGG	60
	GGCACAGCGG	CCCTGGGCTG	CCTGGTCAAG	GACTACTTCC	CCGAACCGGT	GACGGTGTCTG	120
	TGGAACTCAG	GCGCCCTGAC	CAGCGGCGTG	CACACCTTCC	CGGCTGTCTT	ACAGTCCTCA	180
	GGACTCTACT	CCCTCAGCAG	CGTGGTGACC	GTGCCCTCCA	GCAGCTTGGG	CACCCAGACC	240
30	TACATCTGCA	ACGTGAATCA	CAAGCCCAGC	AACACCAAGG	TGGACAAGAA	AGTTGAGCCC	300
	AAATCTTG TG	ACAAAATCTA	CACATGCCCA	CCGTGCCCAG	CACCTGAACT	CCTGGGGGGA	360
	CCGTCAGTCT	TCCTCTTCCC	CCCAAAACCC	AAGGACACCC	TCATGATCTC	CCGGACCCCT	420
	GAGGTCACAT	GCGTGGTGGT	GGACGTGAGC	CACGAAGACC	CTGAGGTCAA	GTTCAACTGG	480
35	TACGTGGACG	GCGTGGAGGT	GCATAATGCC	AAGACAAAGC	CGCGGGAGGA	GCAGTACAAC	540
	AGCACGTACC	GGGTGGTCAG	CGTCTTCACC	GTCTTCGACC	AGGACTGGCT	GAATGGCAAG	600
	GAGTACAAGT	GCAAGGTCTC	CAACAAAGCC	CTCCAGCCCC	CCATCGAGAA	AACCATCTCC	660
40	AAAGCCAAAG	GGCAGCCCCG	AGAACCACAG	GTGTACACCC	TGCCCCCATC	CCGGGAGGAG	720
	ATGACCAAGA	ACCAGGTCAG	CCTGACCTGC	CTGGTCAAAG	GCTTCTATCC	CAGCGACATC	780
	GCCGTGGAGT	GGGAGAGCAA	TGGGCAGCCG	GAGAACAAC T	ACAAGACCAC	GCCTCCCGTG	840
45	CTGGAICTCG	ACGGCTCCTT	CTTCTCTTAC	AGCAAGCTCA	CCGTGGACAA	GAGCAGGTGG	900
	CAGCAGGGGA	ACGTCTTCTC	ATGTCCCGTG	ATGCATGAGG	CTCTGCACAA	CCACTACACG	960
	CAGAAGAGCC	TCTCCCTGTC	TCCGGGTAAA				990

(2) INFORMATION FOR SEO ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 330 amino acids
(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

10 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 15 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 20 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 25 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140
 30 Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175
 Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190
 35 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220
 40 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 225 230 235 240
 Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255
 45 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300
 50 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
325 330

(2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 427 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

AAGCTTGCCG CCACCATGGA TTCACAGGCC CAGGTTCTTA TGTTACTGCC GCTATGGGTA 60
TCTGGTACCT GTGGGGACAT TGTGATGTCA CAGTCTCCAT CCTCCCTAGC TGTGTCAGTT 120
GGAGAGAAGG TTACTATGAG CTGCAAGTCC AGTCAGAGCC TTTTATATAG TCGTAATCAA 180
AAGAACTACT TGGCCTGGTT CCAGCAGAAG CCAGGGCAGT CTCCTAAACT GCTGATTTTC 240
TGGGCATCCA CTAGGGAATC TGGGGTCCCT GATCGCTTCA CAGGCAGTGG ATTTGGGACG 300
GATTTCAATC TCACCATCAG CAGTGTGCAG GCTGAGGACC TGGCAGTTTA TGA CTGTCAG 360
CAATATTTTA GCTATCCGCT CACGTTCCGT GCTGGGACCA AGCTGGAGCT GAAACGTGAG 420
TGGATCC 427

(2) INFORMATION FOR SEQ ID NO: 24:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 133 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Met Asp Ser Gln Ala Gln Val Leu Met Leu Leu Pro Leu Trp Val Ser
1 5 10 15
Gly Thr Cys Gly Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala
20 25 30
Val Ser Val Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser
35 40 45
Leu Leu Tyr Ser Arg Asn Gln Lys Asn Tyr Leu Ala Trp Phe Gln Gln
50 55 60
Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg
65 70 75 80
Glu Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Phe Gly Thr Asp
85 90 95
Phe Asn Leu Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr
100 105 110

Asp Cys Gln Gln Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr
 115 120 125

Lys Leu Glu Leu Lys
 130

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 457 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

AAGCTTGCCG CCACCATGGG ATGGAGCTGG GTCTTTCTCT TTCTCCTGTC AGGAACTGCA 60
 GGTGTCCTCT CTGAGGTCCA GCTGCAACAG TCTGGACCTG AGCTGGTGAA GCCTGGGGCT 120
 TCAGTAAAGA TGTCTGCAA GACTTCTAGA TACACATTCA CTGAATACAC CATACTGG 180
 GTGAGACAGA GCCATGGAAA GAGCCTTGAG TGGATTGGAG GTATTAATCC TAACAATGGT 240
 ATTCCTAACT ACAACCAGAA GTTCAAGGGC AGGGCCACAT TGA CTGTAGG CAAGTCCTCC 300
 AGCACCGCCT ACATGGAGCT CCGCAGCCTG ACATCTGAGG ATTCTGCGGT CTATTTCTGT 360
 GCAAGAAGAA GAATCGCCTA TGGTTACGAC GAGGGCCATG CTATGGACTA CTGGGGTCAA 420
 GGAACCTCAG TCACCGTCTC CTCAGGTGAG TGGATCC 457

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 143 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met Gly Trp Ser Trp Val Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
 1 5 10 15
 Val Leu Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
 20 25 30
 Pro Gly Ala Ser Val Lys Met Ser Cys Lys Thr Ser Arg Tyr Thr Phe
 35 40 45
 Thr Glu Tyr Thr Ile His Trp Val Arg Gln Ser His Gly Lys Ser Leu
 50 55 60
 Glu Trp Ile Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn
 65 70 75 80
 Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ser Ser
 85 90 95

Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val
100 105 110

Tyr Phe Cys Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His
115 120 125

Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser
130 135 140

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8068 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

20	GAATTCACAGC AACTGGCGG CCGTTACTAG TTATTAATAG TAATCAATTA CGGGGTCATT	60
	AGTTCATAGC CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCGCCCTGG	120
	CTGACCGCCC AACGACCCCC GCCCATGAC GTCAATAATG ACGTATGTTT CCATAGTAAC	180
	GCCAAATAGG ACTTTCCATT GACGTCAATG GGTGGAGTAT TTACGGTAAA CTGCCCACTT	240
25	GGCAGTACAT CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA ATGACGGTAA	300
	ATGGCCCGCC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCTTA CTGGCAGTA	360
	CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACATCAATGG	420
30	GCGTGGATAG CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG	480
	GAGTTTGTIT TGGCACCAA ATCAACGGGA CTTTCCAAA TGTCGTAACA ACTCCGCCCC	540
	ATTGACGCAA ATGGGCGGTA GCGGTGTACG GTGGGAGGTC TATATAAGCA GAGTCGTTT	600
	AGTGAACCGT CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA	660
35	CCGGGACCGA TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCCGTGC	720
	CAAGAGTGAC GTAAGTACCG CCTATAGAGT CTATAGGCCC ACCCCCTTGG CTTCTTATGC	780
	ATGCTATACT GTTTTGGCT TGGGGTCTAT ACACCCCGC TTCCTCATGT TATAGGTGAT	840
40	GGTATAGCTT AGCCTATAGG TGTGGGTTAT TGACCATTAT TGACCACTCC CCTATTGGTG	900
	ACGATACTTT CCATTACTAA TCCATAACAT GGCTCTTTGC CACAACCTCT TTTATTGGCT	960
	ATATGCCAAT ACACGTCTCT TCAGAGACTG ACACGGACTC TGTATTTTTA CAGGATGGGG	1020
45	TCTCATTTAT TATTTACAAA TTCACATATA CAACACCACC GTCCCCAGTG CCCGCAGTTT	1080
	TTATTAAACA TAACGTGGGA TCTCCACGCG AATCTCGGGT ACGTGTCCG GACATGGGCT	1140
	CTTCTCCGGT AGCGGCGGAG CTTCTACATC CGAGCCCTGC TCCCATGCCT CCAGCGACTC	1200
	ATGGTCGCTC GGCAGCTCCT TGCTCCTAAC AGTGGAGGCC AGACTTAGGC ACAGCACGAT	1260
50	GCCCCACCACC ACCAGTGTGC CGCACAAGGC CGTGGCGGTA GGGTATGTGT CTGAAAATGA	1320
	GCTCGGGGAG CGGGCTTGCA CCGCTGACGC ATTTGGAAGA CTTAAGGCAG CGGCAGAAGA	1380

AGATGCAGGC AGCTGAGTTG TTGTGTTCTG ATAAGAGTCA GAGGTAACTC CCGTTGCGGT 1440
 GCTGTTAACG GTGGAGGGCA GTGTAGTCTG AGCAGTACTC GTTGCTGCCG CGCGCGCCAC 1500
 5 CAGACATAAT AGCTGACAGA CTAACAGACT GTTCCTTTCC ATGGGTCTTT TCTGCAGTCA 1560
 CCGTCCTTGA CACGCGTCTC GGAAGCTTG CCGCCACCAT GGATTCACAG GCCCAGGTTT 1620
 TTATGTTACT GCCGCTATGG GTATCTGGTA CCTGTGGGA CATTGTGATG TCACAGTCTC 1680
 10 CATCCTCCCT AGCTGTGTCA GTTGGAGAGA AGGTTACTAT GAGCTGCAAG TCCAGTCAGA 1740
 GCCTTTTATA TTCTAGAAAT CAAAAGAACT ACTTGGCCTG GTTCCAGCAG AAGCCAGGGC 1800
 AGTCTCCTAA ACTGCTGATT TTCTGGGCAT CCACTAGGGA ATCTGGGGTC CCTGATCGCT 1860
 TCACAGGCAG TGGATTTGGG ACGGATTTCA ATCTCACCAT CAGCAGTGTG CAGGCTGAGG 1920
 15 ACCTGGCAGT TTATGACTGT CAGCAATATT TTAGCTATCC GCTCACGTTT GGTGCTGGGA 1980
 CCAAGCTGGA GCTGAAACGT GAGTGGATCC ATCTGGGATA AGCATGCTGT TTCTGTCTG 2040
 TCCCTAACAT GCCCTGTGAT TATGCGCAAA CAACACACCC AAGGGCAGAA CTTTGTACT 2100
 20 TAAACACCAT CCTGTTTGCT TCTTCTCTCA GGAAGTGTG CTGCACCATC TGTCTTCATC 2160
 TTCCCGCCAT CTGATGAGCA GTTGAAATCT GGAAGTGCCT CTGTTGTGTG CCTGCTGAAT 2220
 AACTTCTATC CCAGAGAGGC CAAAGTACAG TGGAAGGTGG ATAACGCCCT CCAATCGGGT 2280
 AACTCCCAGG AGAGTGTAC AGAGCAGGAC AGCAAGGACA GCACCTACAG CCTCAGCAGC 2340
 25 ACCCTGACGC TGAGCAAAGC AGACTACGAG AAACACAAAG TCTACGCCCTG CGAAGTCACC 2400
 CATCAGGGCC TGAGCTCGCC CGTCACAAAG AGCTTCAACA GGGGAGAGTG TTAGAGGGAG 2460
 AAGTGCCCCC ACCTGCTCCT CAGTTCACG CTGACCCCTT CCCATCCTTT GGCCTCTGAC 2520
 30 CCTTTTCCA CAGGGGACCT ACCCTATTG CGGTCTCCA GCTCATCTTT CACCTCACCC 2580
 CCCTCCTCCT CCTTGGCTTT AATTATGCTA ATGTTGGAGG AGAATGAATA AATAAAGTGA 2640
 ATCTTTGCAC CTGTGTTGGA TCTAATAAAA GATATTTAT TTATTAGAT ATGTGTGTG 2700
 35 GTTTTTGTG TGCACTGCCT CTATCTGGAG GCCAGGTAGG GCTGGCCTTG GGGGAGGGG 2760
 AGGCCAGAAT GACTCCAAGA GCTACAGGAA GGCAGGTCAG AGACCCCACT GGACAAACAG 2820
 TGGCTGGACT CTGCACCATA ACACACAATC AACAGGGGAG TGAGCTGGAA ATTTGCTAGC 2880
 GAATTCTTGA AGACGAAAGG GCCTCGTGAT ACGCCTATTT TTATAGGTTA ATGTCATGAT 2940
 40 AATAATGGTT TCTTAGACGT CAGGTGGCAC TTTTCGGGA AATGTGCGCG GAACCCCTAT 3000
 TTGTTTATTT TTCTAAATAC ATTCAAATAT GTATCCGCTC ATGAGACAAT AACCTGATA 3060
 AATGCTTCAA TAATATTGAA AAAGGAAGAG TATGAGTATT CAACATTTCC GTGTCGCCCT 3120
 45 TATTCCTTTT TTGCGGCAT TTGCGCTTC TGTTTTGTG CACCCAGAAA CGCTGGTGAA 3180
 AGTAAAAGAT GCTGAAGATC AGTTGGGTGC ACGAGTGGGT TACATCGAAC TGGATCTCAA 3240
 CAGCGGTAAG ATCCTTGAGA GTTTTCGCC CGAAGAACGT TTTCCAATGA TGAGCACTTT 3300
 50 TAAAGTTCTG CTATGTGGCG CGGTATTATC CCGTGTGAC GCCGGGCAAG AGCAACTCGG 3360
 TCGCCGCATA CACTATTCTC AGAATGACTT GGTGAGTAC TCACCAGTCA CAGAAAAGCA 3420
 TCTTACGGAT GGCATGACAG TAAGAGAATT ATGCAGTGCT GCCATAACCA TGAGTGATAA 3480

	CACTGCGGCC	AACITACTTC	TGACAACGAT	CGGAGGACCG	AAGGAGCTAA	CCGCTTTTTT	3540
	GCACAACATG	GGGGATCATG	TAACTCGCCT	TGATCGTTGG	GAACCGGAGC	TGAATGAAGC	3600
5	CATACCAAAC	GACGAGCGTG	ACACCACGAT	GCCTGCAGCA	ATGGCAACAA	CGTTGCGCAA	3660
	ACTATTAACT	GGCGAACTAC	TTACTCTAGC	TTCCCGGCAA	CAATTAATAG	ACTGGATGGA	3720
	GGCGGATAAA	GTTGCAGGAC	CACTTCTGCG	CTCGGCCCTT	CCGGCTGGCT	GGTTTATTGC	3780
10	TGATAAATCT	GGAGCCGGTG	AGCGTGGGTC	TCGCGGTATC	ATTGCAGCAC	TGGGGCCAGA	3840
	TGGTAAGCCC	TCCCGTATCG	TAGTTATCTA	CACGACGGGG	AGTCAGGCAA	CTATGGATGA	3900
	ACGAAATAGA	CAGATCGCTG	AGATAGGTGC	CTCACTGATT	AAGCATTGGT	AACTGTCAGA	3960
	CCAAGTTTAC	TCATATATAC	TTTAGATTGA	TTTAAACTT	CATTTTTAAT	TTAAAAGGAT	4020
15	CTAGGTGAAG	ATCCTTTTTG	ATAATCTCAT	GACCAAAATC	CCTTAACGTG	AGTTTTCGTT	4080
	CCACTGAGCG	TCAGACCCCG	TAGAAAAGAT	CAAAGGATCT	TCTTGAGATC	CTTTTTTCT	4140
	GCGCGTAATC	TGCTGCTTGC	AAACAAAAAA	ACCACCGCTA	CCAGCGGTGG	TTTGTTTGCC	4200
20	GGATCAAGAG	CTACCAACTC	TTTTTCCGAA	GGTAACTGGC	TTCAGCAGAG	CGCAGATACC	4260
	AAATACTGTC	CTTCTAGTGT	AGCCGTAGTT	AGGCCACCAC	TTCAAGAACT	CTGTAGCACC	4320
	GCCTACATAC	CTCGCTCTGC	TAATCCTGTT	ACCACTGGCT	GCTGCCAGTG	GCGATAAGTC	4380
	GTGTCTTACC	GGGTGGACT	CAAGACGATA	GTTACCGGAT	AAGGCGCAGC	GGTCGGGCTG	4440
25	AACGGGGGGT	TCGTGCACAC	AGCCCAGCTT	GGAGCGAACG	ACCTACACCG	AACTGAGATA	4500
	CCTACAGCGT	GAGCTATGAG	AAAGCGCCAC	GCTTCCCGAA	GGGAGAAAGG	CGGACAGGTA	4560
	TCCGGTAAGC	GGCAGGGTCG	GAACAGGAGA	GCGCACGAGG	GAGCTTCCAG	GGGGAAACGC	4620
30	CTGGTATCTT	TATAGTCTTG	TCGGGTTTCG	CCACCTCTGA	CTTGAGCGTC	GATTTTTGTG	4680
	ATGCTCGTCA	GGGGGGCGGA	GCCTATGGAA	AAACGCCAGC	AACGCGGCCT	TTTTACGGTT	4740
	CCTGGCCTTT	TGCTGGCCTT	TTGCTCACAT	GTTCTTTCCT	GCGTTATCCC	CTGATTCTGT	4800
35	GGATAACCGT	ATTACCGCCT	TTGAGTGAGC	TGATACCGCT	CGCCGCAGCC	GAACGACCGA	4860
	GCGCAGCGAG	TCAGTGAGCG	AGGAAGCGGA	AGAGCGCCTG	ATGCGGTATT	TTCTCCTTAC	4920
	GCATCTGTGC	GGTATTTTAC	ACCGCATATG	GTGCACTCTC	AGTACAATCT	GCTCTGATGC	4980
	CGCATAGTTA	AGCCAGTATA	CACTCCGCTA	TCGCTACGTG	ACTGGGTCAT	GGCTGCGCCC	5040
40	CGACACCCGC	CAACACCCGC	TGACGCGCCC	TGACGGGCTT	GTCTGCTCCC	GGCATCCGCT	5100
	TACAGACAAG	CTGTGACCGT	CTCCGGGAGC	TGCATGTGTC	AGAGGTTTTT	ACCGTCATCA	5160
	CCGAAACGCG	CGAGGCAGCT	GTGGAATGTG	TGTCAGTTAG	GGTGTGGAAA	GTCCCCAGGC	5220
45	TCCCCAGCAG	GCAGAAGTAT	GCAAAGCATG	CATCTCAATT	AGTCAGCAAC	CAGGCTCCCC	5280
	AGCAGGCAGA	AGTATGCAAA	GCATGCATCT	CAATTAGTCA	GCAACCATAG	TCCCGCCCTT	5340
	AACTCCGCCC	ATCCCGCCCC	TAACTCCGCC	CAGTTCCGCC	CATTCTCCGC	CCCATGGCTG	5400
50	ACTAATTTTT	TTTATTTTATG	CAGAGGCCGA	GGCCGCCTCG	GCCTCTGAGC	TATTCCAGAA	5460
	GTAGTGAGGA	GGCTTTTTTG	GAGGCCTAGG	CTTTTGCAAA	AAGCTAGCTT	CACGCTGCCG	5520
	CAAGCACTCA	GGGCGCAAGG	GCTGTAAAG	GAAGCGGAAC	ACGTAGAAAG	CCAGTCCGCA	5580
55							

5 GAAACGGTGC TGACCCCGGA TGAATGTCAG CTA CTGGGCT ATCTGGACAA GGGAAAACGC 5640
 AAGCGCAAAG AGAAAGCAGG TAGCTTGACAG TGGGCTTACA TGGCGATAGC TAGACTGGGC 5700
 5 GGTTTTATGG ACAGCAAGCG AACCGGAATT GCCAGCTGGG GCGCCCTCTG GTAAGGTTGG 5760
 GAAGCCCTGC AAAGTAACT GGATGGCTTT CTGCGGCCA AGGATCTGAT GCGCAGGGG 5820
 ATCAAGATCT GATCAAGAGA CAGGATGAGG ATCGTTTCGC ATGATTGAAC AAGATGGATT 5880
 10 GCACGCAGGT TCTCCGGCCG CTGGGGTGA GAGGCTATTG GGCTATGACT GGGCACAACA 5940
 GACAATCGGC TGCTCTGATG CCGCGGTGTT CCGGCTGTCA GCGCAGGGG GCGCGGTTCT 6000
 TTTTGTCAAG ACCGACCTGT COGGTGCCCT GAATGAACTG CAGGACGAGG CAGCGCGGCT 6060
 ATCGTGGCTG GCCACGACGG GCGTTCCTTG CCGAGCTGTG CTCGACGTTG TCACTGAAGC 6120
 15 GGGGAAGGGAC TGCTGCTAT TGGGCGAAGT GCCGGGGCAG GATCTCCTGT CATCTCACCT 6180
 TGCTCCTGCC GAGAAAGTAT CCATCATGGC TGATGCAATG CGGCGGCTGC ATACGCTTGA 6240
 TCCGGCTACC TGCCCATTCG ACCACCAAGC GAAACATCGC ATCGAGCGAG CACGTACTCG 6300
 20 GATGGAAGCC GGTCTTGTG ATCAGGATGA TCTGGACGAA GAGCATCAGG GGCTCGCGCC 6360
 AGCCGAAC TGCGCCAGGC TCAAGGCGCG CATGCCCGAC GCGGAGGATC TCGTCGTGAC 6420
 CCATGGCGAT GCCTGCTTGC CGAATATCAT GGTGGAAAAT GGCCGCTTTT CTGGATTTCAT 6480
 25 CGACTGTGGC CGGCTGGGTG TGGCGGACCG CTATCAGGAC ATAGCGTTGG CTACCCGTGA 6540
 TATTGCTGAA GAGCTTGCG GCGAATGGCG TGACCGCTTC CTCGTGCTTT ACGGTATCGC 6600
 CGCTCCCGAT TCGCAGCGCA TCGCCTTCTA TCGCCTTCTT GACGAGTTCT TCTGAGCGGG 6660
 ACTCTGGGT TCGAAATGAC CGACCAAGCG ACGCCCAACC TGCCATCAG AGATTTTCGAT 6720
 30 TCCACCGCCG CTTCTATGA AAGGTTGGGC TTCCGAATCG TTTTCCGGGA CGCCGGCTGG 6780
 ATGATCCTCC AGCGCGGGGA TCTCATGCTG GAGTTCTTCG CCCACCCCGG GCTCGATCCC 6840
 CTCGCGAGTT GGTTCAGCTG CTGCCTGAGG CTGGACGACC TCGCGGAGTT CTACCGGCAG 6900
 35 TGCAATCCG TCGGCATCCA GGAACCAGC AGCGGCTATC CGCGCATCCA TGCCCCGAA 6960
 CTGCAGGAGT GGGGAGGCAC GATGGCCGCT TTGGTCCCGG ATCTTTGTGA AGGAACCTTA 7020
 CTTCTGTGGT GTGACATAAT TGGACAACT ACCTACAGAG ATTTAAAGCT CTAAGGTAAA 7080
 TATAAAATTT TTAAGTGTAT AATGTGTTAA ACTACTGATT CTAATTGTTT GTGTATTTTA 7140
 40 GATTCCAACC TATGGAAC TGATGATGAG GCTACTGCTG ACTCTCAACA 7260
 CCTGTTTTC TCAGAAGAAA TGCCATCTAG TGATGATGAG GCTACTGCTG ACTCTCAACA 7260
 TTCTACTCT CCAAAAAGA AGAGAAAGGT AGAAGACCCC AAGGACTTTC CTTCAGAATT 7320
 45 GCTAAGTTTT TTGAGTCATG CTGTGTTTGA TAATAGAACT CTTGCTTGCT TTGCTATTTA 7380
 CACCACAAAG GAAAAAGCTG CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC 7440
 CTTTATAAGT AGGCATAACA GTTATAATCA TAACATACTG TTTTTCCTTA CTCCACACAG 7500
 50 GCATAGAGTG TCTGCTATTA ATAACATATG TCAAAAATTG TGTACCTTTA GCTTTTAAAT 7560
 TTGTAAAGGG GTTAATAAGG AATATTTGAT GTATAGTGCC TTGACTAGAG ATCATAATCA 7620
 GCCATACCAC ATTTGTAGAG GTTTTACTTG CTTTAAAAAA CCTCCACAC CTCCCCCTGA 7680

ACCTGAACA TAAATGAAT GCAATTGTTG TTGTTAACTT GTTTATTGCA GCTTATAATG 7740
 GTTACAAATA AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTITTT TCACTGCATT 7800
 5 CTAGTTGTGG TTTGTCCAAA CTCATCAATG TATCTTATCA TGTCTGGATC TAATAAAGA 7860
 TATTTATTTT CATTAGATAT GTGTGTGGT TTTTGTGTG CAGTGCCTCT ATCTGGAGGC 7920
 CAGGTAGGGC TGGCCTTGGG GGAGGGGGAG GCCAGAATGA CTCCAAGAGC TACAGGAAGG 7980
 10 CAGGTCAGAG ACCCCACTGG ACAACAGTG GCTGGACTCT GCACCATAAC ACACAATCAA 8040
 CAGGGGAGTG AGCTGGAAAT TTGCTAGC 8068

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 239 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Asp Ser Gln Ala Gln Val Leu Met Leu Leu Pro Leu Trp Val Ser Gly
 1 5 10 15
 25 Thr Cys Gly Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val
 20 25 30
 Ser Val Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu
 35 40 45
 30 Leu Tyr Ser Arg Asn Gln Lys Asn Tyr Leu Ala Trp Phe Gln Gln Lys
 50 55 60
 Pro Gly Gln Ser Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu
 65 70 75 80
 35 Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Phe Gly Thr Asp Phe
 85 90 95
 Asn Leu Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Asp
 100 105 110
 40 Cys Gln Gln Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys
 115 120 125
 Leu Glu Leu Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro
 130 135 140
 Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
 145 150 155 160
 45 Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
 165 170 175
 Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp
 180 185 190
 50 Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys
 195 200 205
 Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln
 210 215 220

EP 0 953 639 A1

Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
225 230 235

(2) INFORMATION FOR SEQ ID NO: 29:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7731 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

15	TTGAAGACGA AAGGGCCTCG TGATACGCCT ATTTTATAG GTTAATGTCA TGATAATAAT	60
	GGTTTCTTAG ACGTCAGGTG GCACTTTTCG GGGAAATGTG CGCGGAACCC CTATTTGTTT	120
	ATTTTCTTAA ATACATTCAA ATATGTATCC GCTCATGAGA CAATAACCCCT GATAAATGCT	180
20	TCAATAATAT TGAAAAGGA AGAGTATGAG TATTCAACAT TTCCGTGTCG CCCTTATTC	240
	CTTTTTCGCG GCATTTTGCC TTCCTGTTTT TGCTCACCCA GAAACGCTGG TGAAAGTAAA	300
	AGATGCTGAA GATCAGTTGG GTGCACGAGT GGGTTACATC GAACTGGATC TCAACAGCGG	360
	TAAGATCCTT GAGAGTTTTC GCCCCGAAGA ACGTTTCCA ATGATGAGCA CTTTTAAAGT	420
25	TCTGCTATGT GCGCGGTAT TATCCCGTGT TGACGCCGGG CAAGAGCAAC TCGGTCGCCG	480
	CATACACTAT TCTCAGAATG ACTTGTTGA GTACTACCA GTCACAGAAA AGCATCTTAC	540
	GGATGGCATG ACAGTAAGAG AATTATGCAG TGCTGCCATA ACCATGAGTG ATAACACTGC	600
30	GGCCAACCTA CTCTGACAA CGATCGGAGG ACCGAAGGAG CTAACCGCTT TTTGACAA	660
	CATGGGGGAT CATGTAATC GCCTTGATCG TTGGGAACCG GAGCTGAATG AAGCCATACC	720
	AAACGACGAG CGTGACACCA CGATGCCTGC AGCAATGGCA ACAACGTTGC GCAAACTATT	780
35	AACTGGCGAA CTACTTACTC TAGCTTCCCG GCAACAATTA ATAGACTGGA TGGAGGCGGA	840
	TAAAGTTGCA GGACCACTTC TCGCTCGGC CCTTCCGGCT GGCTGGTTTA TTGCTGATAA	900
	ATCTGGAGCC GGTGAGCGTG GGTCTCGCGG TATCATTGCA GCACTGGGGC CAGATGGTAA	960
	GCCCTCCCGT ATCGTAGTTA TCTACACGAC GGGGAGTCAG GCAACTATGG ATGAACGAAA	1020
40	TAGACAGATC GCTGAGATAG GTGCCTCACT GATTAAGCAT TGGTAACTGT CAGACCAAGT	1080
	TTACTCATAT ATACTTTAGA TTGATTTAAA ACTTCATTTT TAATTTAAAA GGATCTAGGT	1140
	GAAGATCCTT TTTGATAATC TCATGACCAA AATCCCTTAA CGTGAGTTTT CGTTCCACTG	1200
45	AGCGTCAGAC CCCGTAGAAA AGATCAAAGG ATCTTCTTGA GATCCTTTTT TTCTGCGCGT	1260
	AATCTGCTGC TTGCAAACAA AAAAACCACC GCTACCAGCG GTGGTTTGTT TGCCGGATCA	1320
	AGAGCTACCA ACTCTTTTTT CGAAGGTAAC TGGCTTCAGC AGAGCGCAGA TACCAAATAC	1380
50	TGTCCTTCTA GTGTAGCCGT AGTTAGGCCA CCACTTCAAG AACTCTGTAG CACCGCCTAC	1440
	ATACCTCGCT CTGCTAATCC TGTTACCAAGT GGCTGCTGCC AGTGCGGATA AGTCGTGTCT	1500
	TACCGGGTTG GACTCAAGAC GATAGTTACC GGATAAGGCG CAGCGGTCCG GCTGAACGGG	1560

EP 0 953 639 A1

	GGTTCGTGC ACACAGCCCA GCTTGGAGCG AACGACCTAC ACCGAACTGA GATACCTACA	1620
	GCGTGAGCTA TGAGAAAGCG CCACGCTTCC CGAAGGGAGA AAGGCGGACA GGTATCCGGT	1680
5	AAGCGGCAGG GTCGGAACAG GAGAGCGCAC GAGGGAGCTT CCAGGGGGAA ACGCCTGGTA	1740
	TCTTTATAGT CCTGTCGGGT TTCGCCACCT CTGACTTGAG CGTCGATTTC TGTGATGCTC	1800
	GTCAGGGGGG CGGAGCCTAT GGAAAAACGC CAGCAACGCG GCCTTTTTTAC GGTTCCTGGC	1860
10	CTTTTGCTGG CCTTTTGCTC ACATGTTCCT TCCTGCGTTA TCCCCTGATT CTGTGGATAA	1920
	CCGTATTACC GCCTTTGAGT GAGCTGATAC CGCTCGCCGC AGCCGAACGA CCGAGCGCAG	1980
	CGAGTCAGTG AGCGAGGAAG CGGAAGAGCG CCTGATGCGG TATTTTCTCC TTACGCATCT	2040
	GTGCGGTATT TCACACCGCA TATGGTGAC TCTCAGTACA ATCTGCTCTG ATGCCGCATA	2100
15	GTTAAGCCAG TATACACTCC GCTATCGCTA CGTGACTGGG TCATGGCTGC GCCCCGACAC	2160
	CCGCCAACAC CCGCTGACGC GCCCTGACGG GCTTGTCTGC TCCCGGCATC CGCTTACAGA	2220
	CAAGCTGTGA CCGTCTCCGG GAGCTGCATG TGTGAGAGT TTTCAACGTC ATCACCAGAA	2280
20	CGCGCGAGGC AGCATGCATC TCAATTAGTC AGCAACCATA GTCCCGCCCC TAACTCCGCC	2340
	CATCCCGCCC CTAACTCCGC CCAGTTCGCG CCAATCTCCG CCCCATGGCT GACTAATTTT	2400
	TTTTATTTAT GCAGAGGCCG AGGCCGCCCTC GGCCTCTGAG CTATTCCAGA AGTAGTGAGG	2460
25	AGGCTTTTTT GGAGGCCTAG GCTTTTGCAA AAAGCTAGCT TACAGCTCAG GGCTGCGATT	2520
	TGCGGCCAAA CTTGACGGCA ATCCTAGCGT GAAGGCTGGT AGGATTTTAT CCCCCTGCC	2580
	ATCATGGTTC GACCATTGAA CTGCATCGTC GCCGTGTCCC AAAATATGGG GATTGGCAAG	2640
	AACGGAGACC TACCCTGGCC TCCGCTCAGG AACGAGTTCA AGTACTTCCA AAGAATGACC	2700
30	ACAACCTCTT CAGTGAAGG TAAACAGAAT CTGGTGATTA TGGGTAGGAA AACCTGGTTC	2760
	TCCATTCTCG AGAAGAATCG ACCTTTAAAG GACAGAATTA ATATAGTTCT CAGTAGAGAA	2820
	CTCAAAGAAC CACCACGAGG AGCTCATTTT CTTGCCAAAA GTTTGGATGA TGCCTTAAGA	2880
35	CTTATTGAAC AACCGGAATT GGCAAGTAAA GTAGACATGG TTTGGATAGT CGGAGGCAGT	2940
	TCTGTTTACC AGGAAGCCAT GAATCAACCA GGCCACCTCA GACTCTTTGT GACAAGGATC	3000
	ATGCAGGAAT TTGAAAGTGA CACGTTTTTC CCAGAAATG ATTTGGGGAA ATATAAACTT	3060
	CTCCAGAAT ACCCAGGCGT CCTCTCTGAG GTCCAGGAGG AAAAAGGCAT CAAGTATAAG	3120
40	TTTGAAGTCT ACGAGAAGAA AGACTAACAG GAAGATGCTT TCAAGTTCTC TGCTCCCTC	3180
	CTAAAGCTAT GCATTTTAT AAGACCATGG GACTTTTGCT GGCTTTAGAT CTTGTGAAG	3240
	GAACCTTACT TCTGTGGTGT GACATAATTG GACAACTAC CTACAGAGAT TTAAAGCTCT	3300
45	AAGGTAAATA TAAAATTTT AAGTGATATAA TGTGTTAAAC TACTGATTCT AATTGTTTGT	3360
	GTATTTTAGA TTCCAACCTA TGGAACGAT GAATGGGAGC AGTGGTGGAA TGCCTTTAAT	3420
	GAGGAAAACC TGTTTTGCTC AGAAGAAATG CCATCTAGTG ATGATGAGGC TACTGCTGAC	3480
50	TCTCAACATT CTAATCTCC AAAAAGAAG AGAAAGGTAG AAGACCCCAA GGACTTTCCT	3540
	TCAGAATTGC TAAGTTTTT GAGTCATGCT GTGTTTAGTA ATAGAACTCT TGCTTGCTTT	3600
	GCTATTTACA CCACAAAGGA AAAAGCTGCA CTGCTATACA AGAAAATTAT GGAAAAATAT	3660

55

EP 0 953 639 A1

	TCTGTAACCT TTATAAGTAG GCATAACAGT TATAATCATA ACATACTGTT TTTTCTTACT	3720
	CCACACAGGC ATAGAGTGTG TGCTATTAAT AACTATGCTC AAAAATGTG TACCTTTAGC	3780
5	TTTTTAATTT GTAAAGGGGT TAATAAGGAA TATTTGATGT ATAGTGCCTT GACTAGAGAT	3840
	CATAATCAGC CATACCACAT TTGTAGAGGT TTTACTTGCT TTAAAAAACC TCCCACACCT	3900
	CCCCCTGAAC CTGAAACATA AAATGAATGC AATTGTTGTT GTTAACTTGT TTATTGCAGC	3960
10	TTATAATGGT TACAAATAAA GCAATAGCAT CACAAATTTC ACAAATAAAG CATTTTTTTC	4020
	ACTGCATTCT AGTTGTGGTT TGTCCAAACT CATCAATGTA TCTTATCATG TCTGGATCTA	4080
	ATAAAAGATA TTIATTTTCA TTAGATATGT GTGTTGGTTT TTTGTGTGCA GTGCCTCTAT	4140
	CTGGAGGCCA GGTAGGGCTG GCCTTGGGGG AGGGGGAGGC CAGAATGACT CCAAGAGCTA	4200
15	CAGGAAGGCA GGTGAGAGAC CCCACTGGAC AAACAGTGGC TGGACTCTGC ACCATAACAC	4260
	ACAATCAACA GGGGAGTGAG CTGGAAATTT GCTAGCGAAT TCCAGCACAC TGGCGGCCGT	4320
	TACTAGTTAT TAATAGTAAT CAATTACGGG GTCATTAGTT CATAGCCCAT ATATGGAGTT	4380
20	CCGCGTTTACA TAACCTACGG TAAATGGCCC GCCTGGCTGA CCGCCCAACG ACCCCCGCCC	4440
	ATTGACGTCA ATAATGACGT ATGTTCCCAT AGTAACGCCA ATAGGGACTT TCCATTGACG	4500
	TCAATGGGTG GAGTATTTAC GGTAAACTGC CCACCTGGCA GTACATCAAG TGTATCATAT	4560
	GCCAAGTACG CCCCTATTG ACGTCAATGA CCGTAAATGG CCCGCCTGGC ATTATGCCCA	4620
25	GTACATGACC TTATGGGACT TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT	4680
	TACCATGGTG ATGCGGTTTT GGCAGTACAT CAATGGGCGT GGATAGCGGT TTGACTCACG	4740
	GGGATTTCCA AGTCTCCACC CCATTGACGT CAATGGGAGT TTGTTTTGGC ACCAAAATCA	4800
30	ACGGGACTTT CCAAAATGTC GTAACAACCT CGCCCCATTG ACGCAATAG GCGGTAGGCG	4860
	TGTACGGTGG GAGGTCTATA TAAGCAGAGC TCGTTTAGTG AACCGTCAGA TCGCCTGGAG	4920
	ACGCCATCCA CGCTGTTTTG ACCTCCATAG AAGACACCGG GACCGATCCA GCCTCGCGG	4980
	CCGGGAACGG TGCAATTGAA CGCGGATTCC CCGTGCCAAG AGTGACGTAA GTACCGCTA	5040
35	TAGAGTCTAT AGGCCACCC CCTTGGCTTC TTATGCATGC TATACTGTTT TTGGCTTGGG	5100
	GTCTATACAC CCCCGCTTCC TCATGTTATA GGTGATGGTA TAGCTTAGCC TATAGGTGTG	5160
	GGTTATTGAC CATATTGAC CACTCCCCTA TTGGTGACGA TACTTTCCAT TACTAATCCA	5220
40	TAACATGGCT CTTTGCCACA ACTCTCTTTA TTGGCTATAT GCCAATACAC TGTCTTCAG	5280
	AGACTGACAC GGACTCTGTA TTTTACAGG ATGGGGTCTC ATTTATTATT TACAAATTCA	5340
	CATATACAAC ACCACCGTCC CCAAGTCCCG CAGTTTTTAT TAAACATAAC GTGGGATCTC	5400
45	CACGCGAATC TCGGGTACGT GTTCCGGACA TGGGCTCTTC TCCGGTAGCG GCGGAGCTTC	5460
	TACATCCGAG CCCTGCTCCC ATGCCTCCAG CGACTCATGG TCGCTCGGCA GCTCCTTGCT	5520
	CCTAACAGTG GAGGCCAGAC TTAGGCACAG CACGATGCCC ACCACCACCA GTGTGCCGCA	5580
	CAAGGCCGTG GCGGTAGGGT ATGTGTCTGA AAATGAGCTC GGGGAGCGGG CTGACACGCG	5640
50	TGACGCATTT GGAAGACTTA AGGCAGCGGC AGAAGAAGAT GCAGGCAGCT GAGTTGTTGT	5700
	GTTCTGATAA GAGTCAGAGG TAACTCCCGT TGCGGTGCTG TTAACGGTGG AGGGCAGTGT	5760

55

AGTCTGAGCA GTACTCGTTG CTGCCGCGCG CGCCACCAGA CATAATAGCT GACAGACTAA 5820
 CAGACTGTTC CTTTCCATGG GTCTTTTCTG CAGTCACCGT CCTTGACACG CGTCTCGGGA 5880
 5 AGCTTGCCGC CACCATGGGA TGGAGCTGGG TCTTTCTCTT TCTCCTGTCA GGAAGTCAG 5940
 GTGTCTCTC TGAGGTCCAG CTGCAACAGT CTGGACCTGA GCTGGTGAAG CCTGGGGCTT 6000
 CAGTAAAGAT GTCCTGCAAG ACTTCTAGAT ACACATTAC TGAATACACC ATACACTGGG 6060
 10 TGAGACAGAG CCATGGAAAG AGCCTTGAGT GGATTGGAGG TATTAATCCT AACATGGTA 6120
 TTCCTAACTA CAACCAGAAG TTCAAGGGCA GGGCCACATT GACTGTAGGC AAGTCTCTCA 6180
 GCACCGCTA CATGGAGCTC CGCAGCCTGA CATCTGAGGA TTCTGCGGTC TATTTCTGTG 6240
 CAAGAAGAAG AATCGCCTAT GGTACGACG AGGGCCATGC TATGGACTAC TGGGGTCAAG 6300
 15 GAACCTCAGT CACCGTCTCC TCAGGTGAGT GGATCCTCTG CGCCTGGGCC CAGCTCTGTC 6360
 CCACACCGCG GTCACATGGC ACCACCTCTC TTGCAGCCTC CACCAAGGGC CCATCGGTCT 6420
 TCCCCCTGGC ACCCTCCTCC AAGAGCACCT CTGGGGGCAC AGCGGCCCTG GGCTGCCTGG 6480
 20 TCAAGGACTA CTTCCCCGAA CCGTGACGG TGTCTGTGAA CTCAGGCGCC CTGACCAGCG 6540
 GCGTGACAC CTTCCCGGCT GTCCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG 6600
 TGACCGTGCC CTCCAGCAGC TTGGGCACCC AGACCTACAT CTGCAACGTG AATCACAAGC 6660
 CCAGCAACAC CAAGGTGGAC AAGAAAGTTG AGCCCAAATC TTGTGACAAA ACTCACACAT 6720
 25 GCCCACCGTG CCCAGCACCT GAATCCTGG GGGGACCGTC AGTCTTCCTC TTCCCCCAA 6780
 AATCCAAGGA CACCTCATG ATCTCCCGGA CCCCTGAGGT CACATGCGTG GTGGTGGACG 6840
 TGAGCCACGA AGACCCTGAG GTCAAGTTCA ACTGGTACGT GGACGGCGTG GAGGTGCATA 6900
 30 ATGCCAAGAC AAAGCCGCGG GAGGAGCAGT ACAACAGCAC GTACCGGGTG GTCAGCGTCC 6960
 TCACCGTCCT GCACAGGAC TGGCTGAATG GCAAGGAGTA CAAGTGCAAG GTCTCCAACA 7020
 AAGCCCTCCC AGCCCCATC GAGAAAACCA TCTCAAAGC CAAAGGGCAG CCCCAGAAAC 7080
 35 CACAGGTGTA CACCTGCCC CCATCCCGG AGGAGATGAC CAAGAACCAG GTCAGCCTGA 7140
 CCTGCCTGGT CAAAGGCTTC TATCCAGCG ACATCGCCGT GGAGTGGGAG AGCAATGGGC 7200
 AGCCGGAGAA CAACTACAAG ACCACGCCTC CCGTGCTGGA CTCCGACGGC TCCTTCTTCC 7260
 40 TCTACAGCAA GCTCACCGTG GACAAGAGCA GGTGGCAGCA GGGGAACGTC TTCTCATGCT 7320
 CCGTGATGCA TGAGGCTCTG CACAACCACT ACACGCAGAA GAGCCTCTCC CTGTCTCCGG 7380
 GTAAATGAGT GCGACGGCCG GCAAGCCCCG CTCCCCGGC TCTCGCGGTC GCACGAGGAT 7440
 GCTTGGCAGC TACCCCTGT ACATACTTCC CGGGCGCCCA GCATGGAAAT AAAGCACCGG 7500
 45 ATCTAATAAA AGATATTTAT TTTATTAGA TATGTGTGTT GGTTTTTTGT GTGCAGTGCC 7560
 TCTATCTGGA GGCCAGGTAG GGCTGGCCTT GGGGGAGGGG GAGGCCAGAA TGAATCCAAG 7620
 AGCTACAGGA AGGCAGGTCA GAGACCCAC TGGACAAACA GTGGCTGGAC TCTGCACCAT 7680
 50 AACACACAAT CAACAGGGGA GTGAGCTGGA AATTTGCTAG CGAATTAATT C 7731

(2) INFORMATION FOR SEQ ID NO: 30:

(1) SEQUENCE CHARACTERISTICS:

55

(A) LENGTH: 472 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

```

Met Gly Trp Ser Trp Val Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
1      5      10      15
Val Leu Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
20      25      30
Pro Gly Ala Ser Val Lys Met Ser Cys Lys Thr Ser Arg Tyr Thr Phe
35      40      45
Thr Glu Tyr Thr Ile His Trp Val Arg Gln Ser His Gly Lys Ser Leu
50      55      60
Glu Trp Ile Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn
65      70      75      80
Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ser Ser
85      90      95
Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val
100     105     110
Tyr Phe Cys Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His
115     120     125
Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ser
130     135     140
Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr
145     150     155     160
Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro
165     170     175
Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val
180     185     190
His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser
195     200     205
Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile
210     215     220
Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val
225     230     235     240
Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
245     250     255
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
260     265     270
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
275     280     285
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
290     295     300
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln

```

	305					310					315				320
5	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	Gln
					325					330				335	
	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Ala
				340					345					350	
	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Pro
			355					360					365		
10	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Thr
		370					375					380			
	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Ser
	385					390					395				400
15	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Tyr
					405					410				415	
	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Tyr
				420					425					430	
20	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Phe
			435					440					445		
	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Lys
		450					455					460			
	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys							
	465					470									

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

GACATTGTGA	TGACCCAATC	TCCAGACTCT	TTGGCTGTGT	CTCTAGGGGA	GAGGGCCACC	60
ATCAACTGCA	AGTCCAGTCA	GAGCCTTTTA	TATTCTAGAA	ATCAAAAGAA	CTACTTGGCC	120
TGGTATCAGC	AGAAACCAGG	ACAGCCACCC	AAACTCCTCA	TCTTTTGGGC	TAGCACTAGG	180
GAATCTGGGG	TACCTGATAG	GTTCA GTGGC	AGTGGGTTTG	GGACAGACTT	CACCCTCACC	240
ATTAGCAGCC	TGCAGGCTGA	AGATGTGGCA	GTTTATTACT	GTCAGCAATA	TTTTAGCTAT	300
CCGCTCACGT	TCGGACAAGG	GACCAAGGTG	GAAATAAAA			339

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 Lys

(2) INFORMATION FOR SEQ ID NO: 33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Asp Cys Gln Gln
 85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 Lys

(2) INFORMATION FOR SEQ ID NO: 34:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 100 105 110
 Lys

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8068 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

GAATTCACGC AACTGGCGG CGTTACTAG TTATTAATAG TAATCAATTA CGGGGTCATT 60
 AGTTCATAGC CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCCCGCCTGG 120
 CTGACCGCCC AACGACCCCC GCCCATTGAC GTCAATAATG ACGTATGTTC CCATAGTAAC 180
 GCCAATAGGG ACTTTCATT GACGTCAATG GGTGGAGTAT TTACGGTAAA CTGCCCACCTT 240
 GGCAGTACAT CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA ATGACGGTAA 300
 ATGGCCCCGC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA CTTGGCAGTA 360
 CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACATCAATGG 420
 GCGTGGATAG CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG 480
 GAGTTTGTTC TGGCACCAAA ATCAACGGGA CTTTCCAAA TGTCGTAACA ACTCCGCCCC 540
 ATTGACGCAA ATGGGCGGTA GCGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT 600

AGTGAACCGT CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA 660
 CCGGGACCGA TCCAGCCTCC GCGGCCGGA ACGGTGCATT GGAACGCGGA TTCCCCGTGC 720
 5 CAAGAGTGAC GTAAGTACCG CCTATAGAGT CTATAGGCCC ACCCCCTTGG CTTCTTATGC 780
 ATGCTATACT GTTTTTGGCT TGGGGTCTAT ACACCCCCGC TTCTCATGT TATAGGTGAT 840
 GGTATAGCTT AGCCTATAGG TGTGGGTAT TGACCATTAT TGACCACTCC CCTATTGGTG 900
 10 ACGATACTTT CCATTACTAA TCCATAACAT GGCTCTTTGC CACAACCTCTC TTTATTGGCT 960
 ATATGCCAAT AACTGTCTCT TCAGAGACTG ACACGGACTC TGTATTTTTA CAGGATGGGG 1020
 TCTCATTTAT TATTTACAAA TTCACATATA CAACACCACC GTCCCCAGTG CCCGCAGTTT 1080
 TTATTAAACA TAACGTGGGA TCTCCACGCG AATCTCGGGT ACGTGTTCG GACATGGGCT 1140
 15 CTCTCCGGT AGCGCGGAG CTCTACATC CGAGCCCTGC TCCCATGCCT CCAGCGACTC 1200
 ATGGTCGCTC GGCAGCTCCT TGCTCCTAAC AGTGGAGGCC AGACTTAGGC ACAGCACGAT 1260
 GCCCACCACC ACCAGTGTGC CGCACAAGGC CGTGGCGGTA GGGTATGTGT CTGAAAATGA 1320
 20 GCTCGGGGAG CGGGCTTGCA CCGCTGACGC ATTTGGAAGA CTTAAGGCAG CGGCAGAAGA 1380
 AGATGCAGGC AGCTGAGTTG TTGTGTTCTG ATAAGAGTCA GAGGTAATC CCGTTGCGGT 1440
 GCTGTTAACG GTGGAGGGCA GTGTAGTCTG AGCAGTACTC GTTGCTGCCG CGCGCGCCAC 1500
 CAGACATAAT AGCTGACAGA CTAACAGACT GTTCCTTTCC ATGGGTCTTT TCTGCAGTCA 1560
 25 CCGTCCTTGA CACGCGTCTC GGAAGCTTG CCGCCACCAT GGAGACAGAC AACTCCTGC 1620
 TATGGGTGCT GCTGCTCTGG GTTCCAGTTT CCTCCGAGA CATTGTGATG ACCCAATCTC 1680
 CAGACTCTTT GGCTGTGTCT CTAGGGGAGA GGGCCACCAT CAACTGCAAG TCCAGTCAGA 1740
 30 GCCTTTTATA TTCTAGAAAT CAAAAGAACT ACTTGGCCTG GTATCAGCAG AAACCAGGAC 1800
 AGCCACCCAA ACTCCTCATC TTTTGGGCTA GCACTAGGGA ATCTGGGGTA CCGATAGGT 1860
 TCAAGTGCAG TGGGTTTGGG ACAGACTTCA CCCTCACCAT TAGCAGCCTG CAGGCTGAAG 1920
 ATGTGGCAGT TTATTACTGT CAGCAATAIT TTAGCTATCC GCTCACGTTT GGACAAGGGA 1980
 35 CCAAGGTGGA AATAAAACGT GAGTGGATCC ATCTGGGATA AGCATGCTGT TTTCTGTCTG 2040
 TCCCTAACAT GCCCTGTGAT TATGCGCAA CAACACACC AAGGGCAGAA CTTTGTACT 2100
 TAAACACCAT CCGTTTGTCT TCTTCTCTCA GGAAGTGTG CTGCACCATC TGTCTTCATC 2160
 40 TTCCCGCCAT CTGATGAGCA GTTGAAATCT GGAAGTGCCT CTGTTGTGTG CTGCTGAAT 2220
 AACTTCTATC CCAGAGAGGC CAAAGTACAG TGGAAGGTGG ATAACGCCCT CCAATCGGGT 2280
 AACTCCAGG AGAGTGTAC AGAGCAGGAC AGCAAGGACA GCACCTACAG CCTCAGCAGC 2340
 45 ACCCTGACGC TGAGCAAAGC AGACTACGAG AAACACAAAG TCTACGCCTG CGAAGTCACC 2400
 CATCAGGGCC TGAGCTCGCC CGTCACAAAG AGCTTCAACA GGGGAGAGTG TTAGAGGGAG 2460
 AAGTGCCCCC ACCTGCTCCT CAGTTCCAGC CTGACCCCT CCCATCCTTT GGCCTCTGAC 2520
 CCTTTTCCA CAGGGGACCT ACCCCTATTG CGGTCTCCA GCTCATCTTT CACCTCACC 2580
 50 CCTCTCTCCT CCTTGGCTTT AATTATGCTA ATGTTGGAGG AGAATGAATA AATAAAGTGA 2640
 ATCTTTGCAC CTGTGGTGA TCTAATAAAA GATATTIATT TTCATTAGAT ATGTGTGTTG 2700

	GT TTTT TGTG TGCAGTGCCT CTATCTGGAG GCCAGGTAGG GCTGGCCTTG GGGGAGGGGG	2760
	AGGCCAGAAT GACTCCAAGA GCTACAGGAA GGCAGGTCAG AGACCCCACT GGACAAACAG	2820
5	TGGCTGGACT CTGCACCATA ACACACAATC AACAGGGGAG TGAGCTGGAA ATTTGCTAGC	2880
	GAATTCCTGA AGACGAAAGG GCCTCGTGAT ACGCCTATTT TTATAGGTTA ATGTCATGAT	2940
	AATAATGGTT TCTTAGACGT CAGGTGGCAC TTTTCGGGGA AATGTGCGCG GAACCCCTAT	3000
10	TTGTTTATTT TTCTAAATAC ATTCAAATAT GTATCCGCTC ATGAGACAAT AACCCTGATA	3060
	AATGCTTCAA TAATATTGAA AAAGGAAGAG TATGAGTATT CAACATTTCC GTGTCGCCCT	3120
	TATTCCTTTT TTTGCGGCAT TTTGCCCTCC TGT TTTTGCT CACCCAGAAA CGCTGGTGAA	3180
	AGTAAAGAT GCTGAAGATC AGTTGGGTGC ACGAGTGGGT TACATCGAAC TGGATCTCAA	3240
15	CAGCGTAAG ATCCTTGAGA GTTTTCGCCC CGAAGAACGT TTTCCAATGA TGAGCACTTT	3300
	TAAAGTTCTG CTATGTGGCG CGGTATTATC CCGTGTGAC GCCGGGCAAG AGCAACTCGG	3360
	TCGCCGCATA CACTATTCTC AGAATGACTT GGTGAGTAC TCACCAGTCA CAGAAAAGCA	3420
20	TCTTACGGAT GGCATGACAG TAAGAGAATT ATGCAGTGCT GCCATAACCA TGAGTGATAA	3480
	CACTGCGGCC AACTTACTTC TGACAACGAT CGGAGGACCG AAGGAGCTAA CCGCTTTTTT	3540
	GCACAACATG GGGGATCATG TAACTCGCCT TGATCGTTGG GAACCGGAGC TGAATGAAGC	3600
	CATACCAAAC GACGAGCGTG ACACCACGAT GCCTGCAGCA ATGGCAACAA CGTTGCGCAA	3660
25	ACTATTAAC TGGCGAACTAC TTA CTCTAGC TTCCCGGCAA CAATTAATAG ACTGGATGGA	3720
	GGCGGATAAA GTTGCAAGAC CACTTCTGCG CTCGGCCCTT CCGGCTGGCT GGT TTTATTGC	3780
	TGATAAATCT GGAGCCGGTG AGCGTGGGTC TCGCGGTATC ATTGCAGCAC TGGGGCCAGA	3840
30	TGGTAAGCCC TCCCGTATCG TAGTTATCTA CACGACGGGG AGTCAGGCAA CTATGGATGA	3900
	ACGAAATAGA CAGATCGCTG AGATAGGTGC CTCACTGATT AAGCATTGGT AACTGTCAGA	3960
	CCAAGTTTAC TCATATATAC TTTAGATTGA TTTAAACTT CATTTTAAAT TTTAAAGGAT	4020
35	CTAGGTGAAG ATCCTTTTGG ATAATCTCAT GACCAAAATC CCTTAACGTG AGTTTTCGTT	4080
	CCACTGAGCG TCAGACCCCG TAGAAAAGAT CAAAGGATCT TCTTGAGATC CTT TTTTCT	4140
	GCGCGTAATC TGCTGCTTGC AAACAAAAA ACCACCGCTA CCAGCGGTGG TTTGTTTGCC	4200
	GGATCAAGAG CTACCAACTC TTTTTCGAA GGTAACTGGC TTCAGCAGAG CGCAGATACC	4260
40	AAATACTGTC CTTCTAGTGT AGCCGTAGTT AGGCCACCAC TTCAAGAACT CTGTAGCACC	4320
	GCCTACATAC CTCGCTCTGC TAATCCTGTT ACCAGTGGCT GCTGCCAGTG GCGATAAGTC	4380
	GTGTCTTACC GGGTTGGACT CAAGACGATA GTTACCGGAT AAGGCGCAGC GGTGGGGCTG	4440
45	AACGGGGGGT TCGTGACAC AGCCCACTT GGAGCGAAGC ACCTACACCG AACTGAGATA	4500
	CCTACAGCGT GAGCTATGAG AAAGCGCCAC GCTTCCCGAA GGGAGAAAGG CGGACAGGTA	4560
	TCCGGTAAGC GGCAGGGTCG GAACAGGAGA GCGCAGGAG GAGCTTCCAG GGGGAAACGC	4620
	CTGGTATCTT TATAGTCTCG TCGGGTTTCG CCACCTCTGA CTTGAGCGTC GATT TTTTGTG	4680
50	ATGCTCGTCA GGGGGGCGGA GCCTATGGAA AAACGCCAGC AACGCGGCCT TTTTACGGTT	4740
	CCTGGCCTTT TGCTGGCCTT TTGCTCACAT GTTCTTTCCT GCGTTATCCC CTGATTCTGT	4800
55		

GGATAACCGT ATTACCGCT TTGAGTGAGC TGATACCGCT CGCCGCAGCC GAACGACCGA 4860
 GCGCAGCGAG TCAGTGAGCG AGGAAGCGGA AGAGCGCCTG ATGCGGTATT TTCTCCTTAC 4920
 5 GCATCTGTGC GGTATTTTCAC ACCGCATATG GTGCACTCTC AGTACAATCT GCTCTGATGC 4980
 CGCATAGTTA AGCCAGTATA CACTCCGCTA TCGCTACGTG ACTGGGTCAT GGCTGCGCCC 5040
 CGACACCCGC CAACACCCGC TGACGCGCCC TGACGGGCTT GTCTGCTCCC GGCATCCGCT 5100
 10 TACAGACAAG CTGTGACCGT CTCGGGAGC TGCATGTGTC AGAGGTTTTC ACCGTCAATCA 5160
 CCGAAACGCG CGAGGCAGCT GTGGAATGTG TGTCACTTAG GGTGTGGAAA GTCCCCAGGC 5220
 TCCCCAGCAG GCAGAAGTAT GCAAAGCATG CATCTCAATT AGTCAGCAAC CAGGCTCCCC 5280
 AGCAGGCAGA AGTATGCAAA GCATGCATCT CAATTAGTCA GCAACCATAG TCCCGCCCCT 5340
 15 AACTCCGCCC ATCCCGCCCC TAACTCGCC CAGTTCCGCC CATTCTCCGC CCCATGGCTG 5400
 ACTAATTTT TTTATTTATG CAGAGGCCGA GCGCCGCTCG GCCTCTGAGC TATTCCAGAA 5460
 GTAGTGAGGA GGCTTTTTTG GAGGCCTAGG CTTTTGCAAA AAGCTAGCTT CACGCTGCCG 5520
 20 CAAGCACTCA GGGCGCAAGG GCTGCTAAG GAAGCGGAAC ACGTAGAAAG CCAGTCCGCA 5580
 GAAACGGTGC TGACCCCGGA TGAATGTCAG CTACTGGGCT ATCTGGACAA GGGAAAACGC 5640
 AAGCGCAAAG AGAAAGCAGG TAGCTTGAGG TGGGCTTACA TGGCGATAGC TAGACTGGGC 5700
 GGTTTTATGG ACAGCAAGCG AACCGGAATT GCCAGCTGGG GCGCCCTCTG GTAAGGTTGG 5760
 25 GAAGCCCTGC AAAGTAACT GGATGGCTTT CTGCGGCCA AGGATCTGAT GCGCGAGGGG 5820
 ATCAAGATCT GATCAAGAGA CAGGATGAGG ATCGTTTCGC ATGATTGAAC AAGATGGATT 5880
 GCACGCAGGT TCTCCGGCCG CTTGGGTGGA GAGGCTATTC GGCTATGACT GGGCACAACA 5940
 30 GACAATCGGC TGCTCTGATG CCGCGTGT TCGGCTGTCA GCGCAGGGG GCGCGGTTCT 6000
 TTTTGTCAAG ACCGACCTGT CCGGTGCCCT GAATGAACTG CAGGACGAGG CAGCGCGGCT 6060
 ATCGTGGCTG GCCACGACGG GCGTTCCTTG CGCAGCTGTG CTCGACGTTG TCACTGAAGC 6120
 35 GGGAGGGGAC TGGCTGCTAT TGGGCGAAGT GCCGGGGCAG GATCTCCTGT CATCTCACCT 6180
 TGCTCCTGCC GAGAAAGTAT CCATCATGGC TGATGCAATG CGGCGGCTGC ATACGCTTGA 6240
 TCCGGCTACC TGCCCATTCG ACCACCAAGC GAAACATCGC ATCGAGCGAG CACGTACTCG 6300
 GATGGAAGCC GGTCTGTGCG ATCAGGATGA TCTGGACGAA GAGCATCAGG GGCTCGCGCC 6360
 40 AGCCGAACTG TTCGCCAGGC TCAAGGCGCG CATGCCCGAC GCGGAGGATC TCGTCGTGAC 6420
 CCATGGCGAT GCCTGCTTGC CGAATATCAT GGTGGAAAAT GCGCGCTTTT CTGGATTAT 6480
 CGACTGTGGC CGGCTGGGTG TGGCGGACCG CTATCAGGAC ATAGCGTTGG CTACCCGTGA 6540
 45 TATTGCTGAA GAGCTTGGCG GCGAATGGGC TGACCGCTTC CTCGTGCTTT ACGGTATCGC 6600
 CGCTCCCGAT TCGCAGCGCA TCGCCTTCTA TCGCCTTCTT GACGAGTTCT TCTGAGCGGG 6660
 ACTCTGGGGT TCGAAATGAC CGACCAAGCG ACGCCAACC TGCCATCACG AGATTTTCGAT 6720
 50 TCCACCGCCG CTTTCTATGA AAGGTTGGGC TTCGGAATCG TTTTCCGGGA GCGCGGCTGG 6780
 ATGATCCTCC AGCGCGGGGA TCTCATGCTG GAGTTCTTCG CCCACCCCGG GCTCGATCCC 6840
 CTCGCGAGTT GGTTCAGCTG CTGCCTGAGG CTGGACGACC TCGCGGAGTT CTACCGGCAG 6900

TGCAAATCCG TCGGCATCCA GGAACCAGC AGCGGCTATC CGCGCATCCA TGCCCCCGAA 6960
 CTGCAGGAGT GGGGAGGCAC GATGGCCGCT TTGGTCCCGG ATCTTTGTGA AGGAACCTTA 7020
 5 CTTCTGTGGT GTGACATAAT TGGACAACT ACCTACAGAG ATTTAAAGCT CTAAGGTAAA 7080
 TATAAAATTT TTAAGTGTAT AATGTGTTAA ACTACTGATT CTAATTGTTT GTGTATTTTA 7140
 GATTCCAACC TATGGAAC TGAAATGGGA GCAGTGGTGG AATGCCTTTA ATGAGGAAAA 7200
 10 CCTGTTTTGC TCAGAAGAAA TGCCATCTAG TGATGATGAG GCTACTGCTG ACTCTCAACA 7260
 TTCTACTCCT CCAAAAAAGA AGAGAAAGGT AGAAGACCCC AAGGACTTTC CTTCAGAATT 7320
 GCTAAGTTTT TTGAGTCATG CTGTGTTTAG TAATAGAACT CTTGCTTGCT TTGCTATTTA 7380
 CACCACAAAG GAAAAAGCTG CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC 7440
 15 CTTTATAAGT AGGCATAACA GTTATAATCA TAACATACTG TTTTTTCTTA CTCCACACAG 7500
 GCATAGAGTG TCTGCTATTA ATAACTATGC TCAAAAATTG TGTACCTTTA GCTTTTTAAT 7560
 TTGTAAAGGG GTTAATAAGG AATATTTGAT GTATAGTGCC TTGACTAGAG ATCATAATCA 7620
 20 GCCATACCAC ATTTGTAGAG GTTTTACTTG CTTTAAAAAA CCTCCACAC CTCCCCCTGA 7680
 ACCTGAAACA TAAATGAAT GCAATTGTTG TTGTTAACTT GTTTATTGCA GCTTATAATG 7740
 GTTACAAATA AAGCAATAGC ATCACAATT TCACAAATAA AGCATTTTTTT TCACTGCATT 7800
 CTAGTTGTGG TTTGTCCAAA CTCATCAATG TATCTTATCA TGCTGGATC TAATAAAGA 7860
 25 TATTTATTTT CATTAGATAT GTGTGTGGT TTTTGTGTG CAGTGCCTCT ATCTGGAGGC 7920
 CAGGTAGGGC TGGCCTTGGG GGAGGGGGAG GCCAGAATGA CTCCAAGAGC TACAGGAAGG 7980
 CAGGTCAGAG ACCCCACTGG ACAACAGTG GCTGGACTCT GCACCATAAC ACACAATCAA 8040
 30 CAGGGGAGTG AGCTGGAAT TTGCTAGC 8068

(2) INFORMATION FOR SEQ ID NO: 36:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 234 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1 5 10 15
 45 Gly Ser Ser Gly Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala
 20 25 30
 Val Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser
 35 40 45
 50 Leu Leu Tyr Ser Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln
 50 55 60
 Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile Phe Trp Ala Ser Thr Arg
 65 70 75 80

Glu Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Phe Gly Thr Asp
 85 90 95
 Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr
 100 105 110
 Tyr Cys Gln Gln Tyr Phe Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr
 115 120 125
 Lys Val Glu Ile Lys Arg Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140
 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160
 Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175
 Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190
 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205
 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 210 215 220
 Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230

(2) INFORMATION FOR SEQ ID NO: 37:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 372 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

CAGGTGCAAC TAGTGCAATC CGCGCCGAA GTGAAGAAAC CCGGTGCTTC CGTGAAAGTC 60
 AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA TACACTGGGT TAGACAGGCC 120
 CCTGGCCAAA GGCTGGAGTG GATAGGAGGT ATTAATCCTA ACAATGGTAT TCCTAACTAC 180
 AACCAGAAAT TCAAGGGCCG GGCCACCTTG ACCGTAGGCA AGTCTGCCAG CACCGCCTAC 240
 ATGGAAGTGT CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA 300
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG AACCCCTTGTC 360
 ACCGTCTCCT CA 372

(2) INFORMATION FOR SEQ ID NO: 38:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 124 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

5 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 10 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 15 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110
 20 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

35 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 40 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 45 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Phe Cys
 85 90 95
 Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110
 50 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 124 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Glu Tyr
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His Ala Met Asp
 100 105 110

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7731 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

TTGAAGACGA AAGGGCCTCG TGATACGCCT ATTTTATAG GTTAATGTCA TGATAATAAT	60
GGTTTCTTAG ACGTCAGGTG GCACTTTTCG GGGAAATGTG CGCGGAACCC CTATTTGTTT	120
ATTTTCTTAA ATACATTCAA ATATGTATCC GCTCATGAGA CAATAACCCCT GATAAATGCT	180
TCAATAATAT TGAAAAAGGA AGAGTATGAG TATTCAACAT TTCCGTGTG CCGTTATTCC	240
CTTTTTCGCG GCATTTTGCC TTCTGTGTTT TGCTCACCCA GAAACGCTGG TGAAAGTAAA	300
AGATGCTGAA GATCAGTTGG GTGCACGAGT GGGTTACATC GAACTGGATC TCAACAGCGG	360
TAAGATCCTT GAGAGTTTC GCCCGAAGA ACGTTTCCA ATGATGAGCA CTTTAAAGT	420
TCTGCTATGT GCGCGGTAT TATCCCGTGT TGACGCGGG CAAGAGCAAC TCGGTCGCCG	480
CATACACTAT TCTCAGAAATG ACTTGGTTGA GTACTCACCA GTCACAGAAA AGCATCTTAC	540
GGATGGCATG ACAGTAAGAG AATTATGCAG TGCTGCCATA ACCATGAGTG ATAACACTGC	600
GGCCAACTTA CTTCTGACAA CGATCGGAGG ACCGAAGGAG CTAACGCTT TTTGCACAA	660
CATGGGGGAT CATGTAATC GCCTTGATCG TTGGGAACCG GAGCTGAATG AAGCCATACC	720
AAACGACGAG CGTGACACCA CGATGCCTGC AGCAATGGCA ACAACGTTGC GCAAATATT	780
AACTGGCGAA CTACTTACTC TAGCTTCCCG GCAACAATTA ATAGACTGGA TGGAGGCGGA	840
TAAAGTTGCA GGACCACTTC TCGCTCGGC CCTCCGGCT GGCTGGTTTA TTGCTGATAA	900
ATCTGGAGCC GGTGAGCGTG GGTCTCGCG TATCATTGCA GCACTGGGGC CAGATGGTAA	960
GCCCTCCCGT ATCGTAGTTA TCTACACGAC GGGGAGTCAG GCAACTATGG ATGAACGAAA	1020
TAGACAGATC GCTGAGATAG GTGCCCTCACT GATTAAGCAT TGGTAACTGT CAGACCAAGT	1080
TTACTCATAT ATACTTTAGA TTGATTAAA ACTTCATTTT TAATTTAAA GGATCTAGGT	1140
GAAGATCCTT TTTGATAATC TCATGACCAA AATCCCTTAA CGTGAGTTT CGTTCCACTG	1200
AGCGTCAGAC CCCGTAGAAA AGATCAAAGG ATCTTCTTGA GATCCTTTT TTCTGCGCGT	1260
AATCTGCTGC TTGCAACAA AAAAACCACC GCTACCAGCG GTGGTTTGTG TGCCGGATCA	1320
AGAGCTACCA ACTCTTTTC CGAAGGTAAC TGGCTTCAGC AGAGCGCAGA TACCAAATAC	1380
TGCTCTTCTA GTGTAGCCGT AGTTAGGCCA CCACTTCAAG AACTCTGTAG CACCGCCTAC	1440
ATACCTCGCT CTGCTAATCC TGTTACCAGT GGCTGCTGCC AGTGGCGATA AGTCGTGTCT	1500

	TACCGGGTTG GACTCAAGAC GATAGTTACC GGATAAGGCG CAGCGGTCGG GCTGAACGGG	1560
	GGGTTTCGTGC ACACAGCCCA GCTTGGAGCG AACGACCTAC ACCGAACTGA GATACCTACA	1620
5	GCGTGAGCTA TGAGAAAGCG CCACGCTTCC CGAAGGGAGA AAGGCGGACA GGTATCCGGT	1680
	AAGCGGCAGG GTCGGAACAG GAGAGCGCAC GAGGGAGCTT CCAGGGGGAA ACGCCTGGTA	1740
	TCTTTATAGT CCTGTCCGGT TTCGCCACCT CTGACTTGAG CGTCGATTTT TGTGATGCTC	1800
10	GTCAGGGGGG CGGAGCCTAT GGAAAAACGC CAGCAACGCG GCCTTTTTTAC GGTTCCTGGC	1860
	CTTTTGCTGG CCTTTTGCTC ACATGTTCCT TCCTGCGTTA TCCCCTGATT CTGTGGATAA	1920
	CCGTATTACC GCCTTTGAGT GAGCTGATAC CGCTCGCCGC AGCCGAACGA CCGAGCGCAG	1980
	CGAGTCAGTG AGCGAGGAAG CGGAAGAGCG CCTGATGCGG TATTTTCTCC TTACGCATCT	2040
15	GTGCGGTATT TCACACCGCA TATGGTGAC TCTCAGTACA ATCTGCTCTG ATGCCGCATA	2100
	GTTAAGCCAG TATACACTCC GCTATCGCTA CGTGACTGGG TCATGGCTGC GCCCCGACAC	2160
	CCGCCAACAC CCGCTGACGC GCCCTGACGG GCTTGTCTGC TCCC GG CATC CGCTTACAGA	2220
20	CAAGCTGTGA CCGTCTCCGG GAGCTGCATG TGTCAGAGGT TTTCAACGTC ATCACC GAAA	2280
	CGCGCGAGGC AGCATGCATC TCAATTAGTC AGCAACCATA GTCCCGCCCC TAACTCCGCC	2340
	CATCCCGCCC CTAATCCGC CCAGTTCCGC CCATTCTCCG CCCCATGGCT GACTAATTTT	2400
	TTTTATTTAT GCAGAGGCCG AGGCCGCTC GGCCTCTGAG CTATTCCAGA AGTAGTGAGG	2460
25	AGGCTTTTTT GGAGGCCTAG GCTTTTGCAA AAAGCTAGCT TACAGCTCAG GGCTGCGATT	2520
	TCGCGCCAAA CTTGACGGCA ATCCTAGCGT GAAGGCTGGT AGGATTTTAT CCCCCTGCC	2580
	ATCATGGTTC GACCATTGAA CTGCATCGTC GCCGTGTCCT AAAATATGGG GATTGGCAAG	2640
30	AACGGAGACC TACCCTGGCC TCCGCTCAGG AACGAGTTCA AGTACTTCCA AAGARTGACC	2700
	ACAACCTCTT CAGTGAAGG TAAACAGAA CTGGTGATTA TGGGTAGGAA AACCTGGTTC	2760
	TCCATTCTCT AGAAGAATCG ACCTTTAAAG GACAGAATTA ATATAGTTCT CAGTAGAGAA	2820
35	CTCAAAGAAC CACCACGAGG AGCTCATTTT CTGCCCCAAA GTTTGGATGA TGCCTTAAGA	2880
	CTTATTGAAC AACCGGAAT GGCAAGTAAA GTAGACATGG TTTGGATAGT CGGAGGCAGT	2940
	TCTGTTTACC AGGAAGCCAT GAATCAACCA GGCCACCTCA GACTCTTTGT GACAAGGATC	3000
	ATGCAGGAAT TTGAAAGTGA CACGTTTTTC CCAGAAATG ATTTGGGGAA ATATAAACTT	3060
40	CTCCCAGAAT ACCCAGGCGT CCTCTCTGAG GTCCAGGAGG AAAAAGGCAT CAAGTATAAG	3120
	TTTGAAGTCT ACGAGAAGAA AGACTAACAG GAAGATGCTT TCAAGTTCTC TGCTCCCCCTC	3180
	CTAAAGCTAT GCATTTTAT AAGACCATGG GACTTTTGCT GGCTTTAGAT CTTTGTGAAG	3240
45	GAACCTTACT TCTGTGGTGT GACATAATTG GACAACTAC CTACAGAGAT TTAAAGCTCT	3300
	AAGGTAAATA TAAATTTTTT AAGTGATATA TGTGTTAAAC TACTGATTCT AATTGTTTGT	3360
	GTATTTTAGA TTCCAACCTA TGGAAGTAT GAATGGGAGC AGTGGTGGAA TGCCTTTAAT	3420
	GAGGAAAACC TGTTTTGCTC AGAAGAAATG CCATCTAGTG ATGATGAGGC TACTGCTGAC	3480
50	TCTCAACATT CTACTCTCC AAAAAGAAG AGAAGGTAG AAGACCCCAA GGACTTTCCT	3540
	TCAGAATTGC TAAGTTTTT GAGTCATGCT GTGTTAGTA ATAGA ACTCT TGCTTGCTTT	3600

	GCTATTTACA CCACAAAGGA AAAAGCTGCA CTGCTATACA AGAAAATTAT GGAAAAATAT	3660
	TCTGTAACCT TTATAAGTAG GCATAACAGT TATAATCATA ACATACTGTT TTTTCTTACT	3720
5	CCACACAGGC ATAGAGTGTC TGCTATTAAT AACTATGCTC AAAAATTGTG TACCTTTAGC	3780
	TTTTTAATTT GTAAAGGGGT TAATAAGGAA TATTTGATGT ATAGTGCCTT GACTAGAGAT	3840
	CATAATCAGC CATACCACAT TTGTAGAGGT TTTACTTGCT TTAAAAAACC TCCACACCT	3900
10	CCCCCTGAAC CTGAAACATA AAATGAATGC AATTGTTGTT GTTAAC TTGTTG CAGC	3960
	TTATAATGGT TACAAATAAA GCAATAGCAT CACAAATTTC ACAAATAAAG CATTTTTTTC	4020
	ACTGCATTCT AGTGTGGT TGTCCAACT CATCAATGTA TCTTATCATG TCTGGATCTA	4080
	ATAAAGATA TTTATTTTCA TTAGATATGT GTGTTGGTTT TTTGTGTGCA GTGCCTCTAT	4140
15	CTGGAGGCCA GGTAGGGCTG GCCTTGGGGG AGGGGGAGGC CAGAATGACT CCAAGAGCTA	4200
	CAGGAAGGCA GGTACAGAGC CCCACTGGAC AAACAGTGGC TGGACTCTGC ACCATAACAC	4260
	ACAATCAACA GGGGAGTGAG CTGGAATTT GCTAGCGAAT TCCAGCACAC TGGCGGCCGT	4320
20	TACTAGTTAT TAATAGTAAT CAATTACGGG GTCATTAGTT CATAGCCCAT ATATGGAGTT	4380
	CCGCGTTACA TAACTTACGG TAAATGGCCC GCCTGGCTGA CCGCCCAACG ACCCCCGCCC	4440
	ATTGACGTCA ATAATGACGT ATGTTCCCAT AGTAACGCCA ATAGGGACTT TCCATTGACG	4500
	TCAATGGGTG GAGTATTTAC GGTAACTGC CCACTTGGCA GTACATCAAG TGTATCATAT	4560
25	GCCAAGTACG CCCCTATTG ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCCA	4620
	GTACATGACC TTATGGGACT TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT	4680
	TACCATGGTG ATGCGGTTTT GGCAGTACAT CAATGGGCGT GGATAGCGT TTGACTCAGC	4740
30	GGGATTTCCA AGTCTCCACC CCATTGACGT CAATGGGAGT TTGTTTTGGC ACCAAAATCA	4800
	ACGGGACTTT CCAAATGTC GTAACAATC CGCCCCATTG ACGCAAATGG GCGGTAGGCG	4860
	TGTACGGTGG GAGGTCTATA TAAGCAGAGC TCGTTTAGTG AACCGTCAGA TCGCCTGGAG	4920
35	ACGCCATCCA CGCTGTTTTG ACCTCCATAG AAGACACCGG GACCGATCCA GCCTCCGCGG	4980
	CCGGAACGG TGCAATGGAA CGCGGATTCC CCGTGCCAAG AGTGACGTAA GTACCGCCTA	5040
	TAGAGTCTAT AGGCCACCC CTTGGCTTC TTATGCATGC TATACTGTTT TTGGCTTGGG	5100
	GTCTATACAC CCCCGCTTCC TCATGTATA GGTGATGGTA TAGCTTAGCC TATAGGTGTG	5160
40	GGTTATTGAC CATATTGAC CACTCCCCTA TTGGTGACGA TACTTTCCAT TACTAATCCA	5220
	TAACATGGCT CTTTGCCACA ACTCTCTTTA TTGGCTATAT GCCAATACAC TGTCTTCAG	5280
	AGACTGACAC GGACTCTGTA TTTTACAGG ATGGGGTCTC ATTTATTATT TACAAATTCA	5340
45	CATATACAAC ACCACCGTCC CCAGTGCCCG CAGTTTTTAT TAAACATAAC GTGGGATCTC	5400
	CACGCGAATC TCGGTACGT GTTCCGGACA TGGGCTCTTC TCCGGTAGCG GCGGAGCTTC	5460
	TACATCCGAG CCCTGCTCCC ATGCCTCCAG CGACTCATGG TCGCTCGGCA GCTCCTTGCT	5520
50	CCTAACAGTG GAGGCCAGAC TTAGGCACAG CACGATGCCC ACCACCACCA GTGTGCCGCA	5580
	CAAGGCCGTG GCGGTAGGGT ATGTGTCTGA AAATGAGCTC GGGGAGCGGG CTGACCCGC	5640
	TGACGCATTT GGAAGACTTA AGGCAGCGGC AGAAGAAGAT GCAGGCAGCT GAGTTGTTGT	5700
55		

	GTTCTGATAA GAGTCAGAGG TAACTCCCGT TCGGGTGCTG TTAACGGTGG AGGGCAGTGT	5760
	AGTCTGAGCA GTACTCGTTG CTGCCGCGCG CGCCACCAGA CATAATAGCT GACAGACTAA	5820
5	CAGACTGTTC CTTTCCATGG GTCTTTTCTG CAGTCACCGT CTTGACACG CGTCTCGGGA	5880
	AGCTTGCCGC CACCATGGAC TGGACCTGGC GCGTGTITTTG CCTGCTCGCC GTGGCTCCTG	5940
	GGGCCCCACAG CCAGGTGCAA CTGGTGCACT CCGGCGCCGA AGTGAAGAAA CCCGGTGCTT	6000
10	CCGTGAAAGT CAGCTGTAAA ACTAGTAGAT ACACCTTCAC TGAATACACC ATACACTGGG	6060
	TTAGACAGGC CCCTGGCCAA AGGCTGGAGT GGATAGGAGG TATTAATCCT AACAAATGGTA	6120
	TTCCTAACTA CAACCAGAAG TTCRAAGGCC GGGCCACCTT GACCGTAGGC AAGTCTGCCA	6180
	GCACCGCCTA CATGGAACTG TCCAGCCTGC GCTCCGAGGA CACTGCAGTC TACTACTGCG	6240
15	CCAGAAGAAG AATCGCCTAT GGTACGACG AGGGCCATGC TATGGACTAC TGGGGTCAAG	6300
	GAACCTTGT CACCGTCTCC TCAGGTGAGT GGATCCTCTG CGCCTGGGCC CAGCTCTGTC	6360
	CCACACCGCG GTCACATGGC ACCACCTCTC TTGCAGCCTC CACCAAGGGC CCATCGGTCT	6420
20	TCCCCCTGGC ACCCTCCTCC AAGAGCACCT CTGGGGGCAC AGCGGCCCTG GGCTGCCTGG	6480
	TCAAGGACTA CTTCCCCGAA CCGGTGACGG TGTCGTGGAA CTCAGGCGCC CTGACCAGCG	6540
	GCGTGACAC CTTCCCGGCT GTCCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG	6600
25	TGACCGTGCC CTCCAGCAGC TTGGGCACCC AGACCTACAT CTGCAACGTG AATCACAAGC	6660
	CCAGCAACAC CAAGGTGGAC AAGAAAGTTG AGCCCCAATC TTGTGACAAA ACTCACACAT	6720
	GCCCACCGTG CCCAGCACCT GAACTCCTGG GGGGACCGTC AGTCTTCCTC TCCCCCCAA	6780
	AACCCAAGGA CACCTCATG ATCTCCCGGA CCCCTGAGGT CACATGCGTG GTGGTGGACG	6840
30	TGAGCCACGA AGACCCTGAG GTCAAGTTCA ACTGGTACGT GGACGGCGTG GAGGTGCATA	6900
	ATGCCAAGAC AAAGCCGCGG GAGGAGCAGT ACAACAGCAC GTACCGGGTG GTCAGCGTCC	6960
	TCACCGTCCT GCACCAGGAC TGGCTGAATG GCAAGGAGTA CAAGTGCAAG GTCTCCAACA	7020
35	AAGCCCTCCC AGCCCCATC GAGAAAACCA TCTCCAAAGC CAAAGGGCAG CCCCAGAAC	7080
	CACAGGTGTA CACCTGCCC CCATCCCGGG AGGAGATGAC CAAGAACCAG GTCAGCCTGA	7140
	CCTGCCTGGT CAAAGGCTTC TATCCCAGCG ACATCGCCGT GGAGTGGGAG AGCAATGGGC	7200
	AGCCGGAGAA CAACTACAAG ACCACGCCCTC CCGTGTGGA CTCGACGGC TCCTTCTTCC	7260
40	TCTACAGCAA GCTCACCGTG GACAAGAGCA GGTGGCAGCA GGGGAACGTC TTCTCATGCT	7320
	CCGTGATGCA TGAGGCTCTG CACAACCACT ACACGCAGAA GAGCCTCTCC CTGTCTCCGG	7380
	GTAATGAGT GCGACGGCCG GCAAGCCCCG CTCGCCGGC TCTCGCGGTC GCACGAGGAT	7440
45	GCTTGGCAGG TACCCCTGT ACATACTTCC CGGGCGCCCA GCATGGAAAT AAAGCACCGG	7500
	ATCTAATAAA AGATATTTAT TTTCATTAGA TATGTGTGTT GGTITTTTGT GTGCAGTGCC	7560
	TCTATCTGGA GGCCAGGTAG GGCTGGCCTT GGGGGAGGGG GAGGCCAGAA TGAATCCAAG	7620
	AGCTACAGGA AGGCAGGTCA GAGACCCAC TGGACAAACA GTGGCTGGAC TCTGCACCAT	7680
50	AACACACAAT CAACAGGGGA GTGAGCTGGA AATTGCTAG CGAATTAATT C	7731

(2) INFORMATION FOR SEQ ID NO: 43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 472 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

Met Asp Trp Thr Trp Arg Val Phe Cys Leu Leu Ala Val Ala Pro Gly
 1 5 10 15
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
 20 25 30
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Thr Ser Arg Tyr Thr Phe
 35 40 45
 Thr Glu Tyr Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu
 50 55 60
 Glu Trp Ile Gly Gly Ile Asn Pro Asn Asn Gly Ile Pro Asn Tyr Asn
 65 70 75 80
 Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Val Gly Lys Ser Ala Ser
 85 90 95
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Arg Arg Ile Ala Tyr Gly Tyr Asp Glu Gly His
 115 120 125
 Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ser
 130 135 140
 Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr
 145 150 155 160
 Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro
 165 170 175
 Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val
 180 185 190
 His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser
 195 200 205
 Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile
 210 215 220
 Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val
 225 230 235 240
 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
 245 250 255
 Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 260 265 270
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
 275 280 285
 Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
 290 295 300

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
305 310 315 320

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
325 330 335

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
340 345 350

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
355 360 365

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
370 375 380

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
385 390 395 400

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
405 410 415

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
420 425 430

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
435 440 445

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
450 455 460

Ser Leu Ser Leu Ser Pro Gly Lys
465 470

(2) INFORMATION FOR SEQ ID NO: 44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

ACCGTCTCCT CAGGTGAGTG GATCC

(2) INFORMATION FOR SEQ ID NO: 45:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

CCTCTCTTGC AGCC

(2) INFORMATION FOR SEQ ID NO: 46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

CCTCTCTTGC AGCC

(2) INFORMATION FOR SEQ ID NO: 47:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

Thr Val Ser Ser
1

(2) INFORMATION FOR SEQ ID NO: 48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

Ser Thr Lys Gly
1

(2) INFORMATION FOR SEQ ID NO: 49:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

ACCGTCTCCT CAGCCTCCAC CAAGGGC

(2) INFORMATION FOR SEQ ID NO: 50:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 8 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

Thr Val Ser Ser Ser Thr Lys Gly
 1 5

(2) INFORMATION FOR SEQ ID NO: 51:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

ACCGTCTCCT CAGCCTCCAC CAAGGGC

(2) INFORMATION FOR SEQ ID NO: 52:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

Thr Val Ser Ser Ala Ser Thr Lys Gly
 1 5

(2) INFORMATION FOR SEQ ID NO: 53:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

GAAATAAAAC GTGAGTGGAT CC

(2) INFORMATION FOR SEQ ID NO: 54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:
 CTTCTTTTCCT CAGGAACGTG GGCTGCA

(2) INFORMATION FOR SEQ ID NO: 55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

Thr Val Ala Ala
 1

(2) INFORMATION FOR SEQ ID NO: 56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:
 GAAATAAAAC GAACTGTGGC TGCA

(2) INFORMATION FOR SEQ ID NO: 57:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

Glu Ile Lys Thr Val Ala Ala
 1 5

(2) INFORMATION FOR SEQ ID NO: 58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

GAAATAAAAC GAACTGTGGC TGCA

(2) INFORMATION FOR SEQ ID NO: 59:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

Glu Ile Lys Arg Thr Val Ala Ala
1 5

(2) INFORMATION FOR SEQ ID NO: 60:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

Met Asp Ser Gln Ala Gln Val Leu Met Leu Leu Leu Trp Val Ser
1 5 10 15
Gly Thr Cys Gly
20

(2) INFORMATION FOR SEQ ID NO: 61:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

Met Gly Trp Ser Trp Val Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly

1 5 10 15
Val Leu Ser

5

(2) INFORMATION FOR SEQ ID NO: 62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

GCCGCCACC

9

(2) INFORMATION FOR SEQ ID NO: 63:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PRIMER"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

CAGAAAGCTT GCCGCCACCA TGGATTCACA GGCCCAG

30

37

(2) INFORMATION FOR SEQ ID NO: 64:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: peptide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

Met Asp Ser Gln Ala Gln
1 5

45

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PRIMER"

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

CCGAGGATCC ACTCACGTTT CAGCTCCAGC TTGGT

35

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

CAGAAAGCTT GCCGCCACCA TGGGATGGAG CTGGGTC

37

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

Met Gly Trp Ser Trp Val
 1 5

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

CCGAGGATCC ACTCACCTGA GGAGACGGTG ACTGA

35

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

5 GTCATCACAA TGTCTCCGA GGAACCTGGA ACCCAG 36

(2) INFORMATION FOR SEQ ID NO: 70:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PRIMER"

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

CTCCGGAGAC ATTGTGATGA CCCAATCTC 29

(2) INFORMATION FOR SEQ ID NO: 71:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PRIMER"

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

CTCCGGAGAC ATTGTGATGA CCCAATCTC 29

(2) INFORMATION FOR SEQ ID NO: 72:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

45 CAGTCAGAGC CTTTATATT CTAGAAATCA AAAGAACTAC TTGGCCTGGT ATCAGCAGAA 60

ACCAGGACAG CC 72

(2) INFORMATION FOR SEQ ID NO: 73:

50 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

ACCCAGATT CCCTAGTGCT AGCCCAAAG ATGAGGAGTT TGGG

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 67 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

TAGCACTAGG GAATCTGGGG TACCTGATAG GTTCAGTGGC AGTGGGTTTG GGACAGACTT
CACCTC

(2) INFORMATION FOR SEQ ID NO: 75:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 53 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

GTCCCTTGTC CGAACGTGAG CGGATAGCTA AAATATTGCT GACAGTAATA AAC

(2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

GCTCACGTTT GGACAAGGGA CCAAGGTGGA AAT

(2) INFORMATION FOR SEQ ID NO: 77:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

CAGTCAGAGC CTTTATATT CTAGAAATCA AAAGAACTAC TTGGCCTGGT TCCAGCAGAA 60
ACCAGGACAG CC 72

- (2) INFORMATION FOR SEQ ID NO: 78:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

GTCCCTTGTC CGAACGTGAG CGGATAGCTA AATATTGCT GACAGTCATA AACTGCC 57

- (2) INFORMATION FOR SEQ ID NO: 79:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

CCCAAACTCC TCATCTATTG GGCTAGCACT AGGG 34

- (2) INFORMATION FOR SEQ ID NO: 80:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

CCCTAGTGCT AGCCCAATAG ATGAGGAGTT TGGG

34

(2) INFORMATION FOR SEQ ID NO: 81:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

TACGCAAACC GCCTCTC

17

(2) INFORMATION FOR SEQ ID NO: 82:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

GAGTGCACCA TATGCGGT

18

(2) INFORMATION FOR SEQ ID NO: 83:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

AACAGCTATG ACCATG

16

(2) INFORMATION FOR SEQ ID NO: 84:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

GTTTTCCCAG TCACGAC

17

(2) INFORMATION FOR SEQ ID NO: 85:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

GTGTATTTCAG TGAAGGTGTA TCTACTAGTT TTACAGCTGA CTTTCAC

47

(2) INFORMATION FOR SEQ ID NO: 86:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 53 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

TAGTAGATAC ACCTTCACTG AATACACCAT ACACTGGGTT AGACAGGCCCTG

53

(2) INFORMATION FOR SEQ ID NO: 87:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

CCCTTGAAGT TCTGGTTGTA GTTAGGAATA CCATTGTTAG GATTAATACC TCCTATCCAC
 TCCAGCCTTT G

60

71

(2) INFORMATION FOR SEQ ID NO: 88:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

5 TAACTACAAC CAGAAGTTCA AGGGCCGGGC CACCTTGACC GTAGGCAAGT CTGCCAGCAC 60
CGCCTACATG G 71

(2) INFORMATION FOR SEQ ID NO: 89:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 63 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 15 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

20 GCATGGCCCT CGTCGTAACC ATAGGCGATT CTTCTTCTGG CGCAGTAGTA GACTGCAGTG 60
TCC 63

(2) INFORMATION FOR SEQ ID NO: 90:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 30 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

35 CTATGGTTAC GACGAGGGCC ATGCTATGGA CTACTGGGGT CAAGGAAC 48

(2) INFORMATION FOR SEQ ID NO: 91:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 45 (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

50 TAACTACAAC CAGAAGTTCA AGGGCCGGGT CACCATCACC GTAGACACCT CTGCCAGCAC 60
CGCCTACATG G 71

(2) INFORMATION FOR SEQ ID NO: 92:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

GGACACTGCA GTCTACTTCT GCGCCAG

- (2) INFORMATION FOR SEQ ID NO: 93:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

TACGCAAACC GCCTCTC

- (2) INFORMATION FOR SEQ ID NO: 94:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

GAGTGCACCA TATGCGGT

- (2) INFORMATION FOR SEQ ID NO: 95:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 76 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

CCTTTGGCCA GGGGCCTGTC TAACCCAGTG TATGGTGTAT TCAGTGAAGG TGCTATCCAC

TAGTTTCCAC TAGTTT

76

(2) INFORMATION FOR SEQ ID NO: 96:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

GTCACCGTCC TTGACACGCG TCTCGGGA

28

(2) INFORMATION FOR SEQ ID NO: 97:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

TTGGAGGAGG GTGCCAG

17

(2) INFORMATION FOR SEQ ID NO: 98:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

GAGACATTGT GACCCAATCT CC

22

(2) INFORMATION FOR SEQ ID NO: 99:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

GACAGTCATA AACTGCCACA TCTTC

25

(2) INFORMATION FOR SEQ ID NO: 100:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

TTGACACGCG TCTCGGGAAG CTT

23

(2) INFORMATION FOR SEQ ID NO: 101:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PRIMER"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

GGCGCAGAGG ATCCAATCAC CT

22

40 Claims

1. An antibody protein having the complementary determining regions of the monoclonal antibody F19 (ATCC Accession No. HB 8269), said antibody protein specifically binding to fibroblast activation protein, characterized in that it has framework modifications resulting in the improved producibility in host cells as compared to a chimeric antibody having the variable regions of F19 and foreign constant regions.

2. An antibody protein characterised in that it has a variable light chain region and a variable heavy chain region according to claim 1, each joined to a human constant region.

3. The antibody protein of claim 2, wherein said human constant region of the light chain is a human kappa constant region.

4. The antibody protein of claim 2, wherein said human constant region of the heavy chain is a human gamma-1 constant region.

5. An antibody protein according to any one of claims 1 to 4, characterised in that its expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 10.

6. An antibody protein according to any one of claims 1 to 4, characterised in that its expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 20.

7. An antibody protein according to any one of claims 1 to 4, characterised in that its expression levels in crude media samples as determined by ELISA and/or purified antibody yields exceed the expression levels and/or purification yields of the chimeric antibodies without framework modifications by at least a factor of 100.

8. An antibody protein according to any one of claims 1 to 7, characterised in that it displays improved producibility in eucaryotic cells.

9. The antibody protein according to claim 8 wherein said eucaryotic cell is a chinese hamster ovary cell (CHO cell).

10. An antibody protein according to any one of claims 1 to 9, wherein the amino acid in Kabat position 87 of the light chain region is not asparagine.

11. The antibody protein of claim 10, wherein the amino acid in Kabat position 87 of the light chain region is selected from aromatic or aliphatic amino acids.

12. The antibody protein of claim 11, wherein said aromatic amino acid in Kabat position 87 of the light chain region is a tyrosine or phenylalanine.

13. The antibody protein according to any one of claims 1 to 12, wherein the amino acid in Kabat position 36 of the light chain region is selected from aromatic amino acids.

14. An antibody protein according to any one of claims 1 to 13 that contains the variable region of the light chain as set forth in SEQ ID NO: 2.

15. An antibody protein of claim 14 characterised in that the variable region of the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 1.

16. An antibody protein according to any one of claims 1 to 13 that contains the variable region of the light chain as set forth in SEQ ID NO: 6.

17. An antibody protein of claim 16 characterised in that the variable region of the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 5.

18. An antibody protein according to any one of claims 1 to 17 containing a variable region of the heavy chain as set forth in any one of SEQ ID NOs: 8, 10, 12, 14.

19. An antibody protein according to claim 18 characterised in that the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in SEQ ID NOs: 7, 9, 11, 13.

20. An antibody protein according to any one of claims 1 to 14 containing the variable region of the light chain as set forth in SEQ ID NO: 2 and the variable region of the heavy chain as set forth in SEQ ID NOs: 12.

21. The antibody protein of claim 20 characterised in that the variable region of the the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 1 and the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 11.

23. An antibody protein according to any one of claims 1 to 13 containing the variable region of the light chain as set forth in SEQ ID NO: 2 and the variable region of the heavy chain as set forth in SEQ ID NOs: 8.

24. The antibody protein of claim 23 characterised in that the variable region of the the light chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 1 and the variable region of the heavy chain is encoded by a nucleotide sequence as set forth in SEQ ID NO: 7.

25. A nucleotide sequence encoding an antibody protein according to any one of claims 1 to 24.

26. A recombinant DNA vector that contains a nucleotide sequence of claim 25.

27. The recombinant DNA vector of claim 26, said vector being an expression vector.

28. A host cell carrying a vector according to claims 26 or 27.

29. The host cell of claim 28, wherein said host cell is a eucaryotic cell.

30. The host cell of claim 29, wherein said eucaryotic host cell is a mammalian cell.

31. The host cell of claim 30, wherein said host cell is a CHO or a COS cell.

32. A method of producing antibody proteins according to any one of claims 1 to 24, said method comprising the steps of:

- (a) cultivating a host cell according to any one of claims 23 to 26 under conditions where said antibody protein is expressed by said host cell, and
- (b) isolating said antibody protein.

33. The method of claim 32, wherein said host cell is a mammalian cell, preferably a CHO or COS cell.

34. The method of claim 32 or 33, wherein said host cell is cotransfected with two plasmids carrying the expression units for light and heavy chains respectively.

35. An antibody protein according to any one of claims 1 to 24, wherein said antibody protein is conjugated to a therapeutic agent.

36. The antibody protein of claim 35, wherein said therapeutic agent is a therapeutic agent selected from the group consisting of radioisotopes, toxins, toxoids, inflammatory agents and chemotherapeutic agents.

37. The antibody protein of claim 36, wherein said radioisotopes are β -emitting radioisotopes.

38. The antibody protein of claim 37, wherein said radioisotopes are selected from the group consisting of ^{186}Re , ^{188}Re , ^{131}I and ^{90}Y .

39. An antibody protein according to any one of claims 1 to 24, characterised in that it is labeled.

40. The antibody protein of claim 39, wherein said label is a detectable marker.

41. The antibody protein of claim 40, wherein the detectable marker is a detectable marker selected from the group consisting of enzymes, dyes, radioisotopes, and biotin.

42. An antibody protein according to any one of claims 1 to 24 conjugated to an imageable agent.

43. The antibody protein of claim 42, wherein the imageable agent is a radioisotope.

44. The antibody protein of claim 43, wherein said radioisotopes are gamma-emitting radioisotopes??.

45. The antibody protein of claim 44, wherein said radioisotopes is ^{125}I .

46. A pharmaceutical composition containing an antibody protein according to any one of claims 1 to 24 and a pharmaceutically acceptable carrier useful for treating tumors, wherein said tumors are associated with activated stromal fibroblasts.

47. A pharmaceutical composition containing an antibody protein according to any one of claims 35 to 38 and a pharmaceutically acceptable carrier useful for treating tumors, wherein said tumors are associated with activated stromal fibroblasts.

48. A pharmaceutical composition containing an antibody protein according to any one of claims 42 to 45 and a pharmaceutically acceptable carrier useful for imaging the presence of activated stromal fibroblasts in a healing wound, inflamed skin or a tumor, in a human patient.

49. The pharmaceutical composition of claims 46 to 48, wherein said tumors are tumors selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

50. Use of an antibody protein according to anyone of claims 1 to 24 for the treatment of cancer.

51. Use of an antibody protein according to anyone of claims 35 to 38 for the treatment of cancer.

52. Use of an antibody protein according to anyone of claims 42 to 45 for imaging activated activated stromal fibroblasts.

53. Use of an antibody protein according to anyone of claims 39 to 41 for detecting the presence of activated stromal fibroblasts in a sample.

54. A method of treating tumors, wherein the tumor is associated with activated stromal fibroblasts capable of specifically forming a complex with antibody proteins according to any one of claims 1 to 24 or 35 to 38, which comprises contacting the tumor with an amount of said antibody proteins effective to treat the tumor.

55. The method of claim 54, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

56. The method of claim 54, wherein the contacting is effected in vitro.

57. The method of claim 54, wherein the contacting is effected in vivo.

58. A method of detecting the presence of activated stromal fibroblasts in wound healing, inflammation or a tumor, characterised in that

(a) a sample, possibly containing activated stromal fibroblasts, is contacted with an antibody protein according to any one of claims 1 to 24 or 39 to 41 under conditions suitable for the formation of a complex between said antibody and antigen,

(b) detecting the presence of said complex, thereby detecting the presence of activated stromal fibroblasts in wound healing, inflammation or a tumor.

59. The method of claim 58, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

60. The method of claim 58 or 59, wherein the antibody protein is a protein according to any one of claims 39 to 41.

61. A method of imaging the presence of activated stromal fibroblasts in a healing wound, inflamed skin or a tumor, in a human patient, characterised in that

(a) an antibody protein according to any one of claims 1 to 24 conjugated to an imageable agent is administered to a human patient under conditions suitable for the formation of an antibody-antigen complex,

(b) imaging any complex formed in this manner,

(c) thereby imaging the presence of activated stromal fibroblasts in a human patient.

62. The method of claim 61, wherein the tumor is a tumor having cancer cells selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

63. A method of detecting tumor-stroma, characterised in that

- (a) a suitable sample is contacted with an antibody protein according to any one of claims 1 to 24, under conditions suitable for the formation of an antibody-antigen complex,
- (b) detecting the presence of any complex so formed,
- (c) relating the presence of said complex to the presence of tumor-stroma.

5

64. The method of claim 62, wherein said antibody is labelled with a detectable marker.

65. A method of imaging tumor-stroma in a human patient, which comprises

10

- (a) administering to the patient an antibody protein according to any one of claims 42 to 45, under conditions suitable for the formation of an antibody-antigen complex,
- (b) imaging any complex so formed, and thereby imaging the presence of tumor-stroma in a human patient.

15

20

25

30

35

40

45

50

55

Fig. 1

1	11	21	31	41
GACATTGTGA	TGACCCAATC	TCCAGACTCT	TTGGCTGTGT	CTCTAGGGGA
51	61	71	81	91
GAGGGCCACC	ATCAACTGCA	AGTCCAGTCA	GAGCCTTTTA	TATTCTAGAA
101	111	121	131	141
ATCAAAAGAA	CTACTTGGCC	TGGTATCAGC	AGAAACCAGG	ACAGCCACCC
151	161	171	181	191
AAACTCCTCA	TCTTTTGGGC	TAGCACTAGG	GAATCTGGGG	TACCTGATAG
201	211	221	231	241
GTTTCAGTGGC	AGTGGGTTTG	GGACAGACTT	CACCCTCACC	ATTAGCAGCC
251	261	271	281	291
TGCAGGCTGA	AGATGTGGCA	GTTTATTACT	GTCAGCAATA	TTTTAGCTAT
301	311	321	331	339
CCGCTCACGT	TCGGACAAGG	GACCAAGGTG	GAAATAAAA	

Fig. 2

1	11	21	31	41
DIVMTQSPDS	LAVSLGERAT	INCKSSQSLL	YSRNQKNYLA	WYQQKPGQPP
51	61	71	81	91
KLLIFWASTR	ESGVPDRFSG	SGFGTDFTLT	ISSLQAEDVA	VYYCQQYFSY
101	111			
PLTFGQGTKV	EIK			

Fig. 3

1	11	21	31	41
GACATTGTGA	TGACCCAATC	TCCAGACTCT	TTGGCTGTGT	CTCTAGGGGA
51	61	71	81	91
GAGGGCCACC	ATCAACTGCA	AGTCCAGTCA	GAGCCTTTTA	TATTCTAGAA
101	111	121	131	141
ATCAAAAGAA	CTACTTGGCC	TGGT <u>T</u> CAGC	AGAAACCAGG	ACAGCCACCC
151	161	171	181	191
AAACTCCTCA	TCTTTTGGGC	TAGCACTAGG	GAATCTGGGG	TACCTGATAG
201	211	221	231	241
GTTTCAGTGGC	AGTGGGTTTG	GGACAGACTT	CACCCTCACC	ATTAGCAGCC
251	261	271	281	291
TGCAGGCTGA	AGATGTGGCA	GTTTAT <u>G</u> ACT	GTCA <u>A</u> CAATA	TTTTAGCTAT
301	311	321	331	339
CCGCTCACGT	TCGGACAAGG	GACCAAGGTG	GAAATAAAA	

Fig. 4

1	11	21	31	41
DIVMTQSPDS	LAVSLGERAT	INCKSSQSLL	YSRNQKNYLA	W <u>F</u> QOKPGOPP
51	61	71	81	91
KLLIFWASTR	ESGVPDRFSG	SGFGTDFTLT	ISSLQAEDVA	VY <u>D</u> CQYFSY
101	111			
PLTFGQGTKV	EIK			

Fig. 5

1	11	21	31	41
GACATTGTGA	TGACCCAATC	TCCAGACTCT	TTGGCTGTGT	CTCTAGGGGA
51	61	71	81	91
GAGGGCCACC	ATCAACTGCA	AGTCCAGTCA	GAGCCTTTTA	TATTCTAGAA
101	111	121	131	141
ATCAAAAGAA	CTACTTGGCC	TGGTATCAGC	AGAAACCAGG	ACAGCCACCC
151	161	171	181	191
AAACTCCTCA	TCT <u>A</u> TTGGGC	TAGCACTAGG	GAATCTGGGG	TACCTGATAG
201	211	221	231	241
GTTCAGTGGC	AGTGGGTTTG	GGACAGACTT	CACCCTCACC	ATTAGCAGCC
251	261	271	281	291
TGCAGGCTGA	AGATGTGGCA	GTTTATTACT	GTCAGCAATA	TTTTAGCTAT
301	311	321	331	339
CCGCTCACGT	TCGGACAAGG	GACCAAGGTG	GAAATAAAA	

Fig. 6

1	11	21	31	41
DIVMTQSPDS	LAVSLGERAT	INCKSSQSLL	YSRNQKNYLA	WYQOKPGOPP
51	61	71	81	91
KLLI <u>Y</u> WASTR	ESGVPDRFSG	SGFGTDFTLT	ISSLQAEDVA	VYYCQYFSY
101	111			
PLTFGQGTKV	EIK			

Fig. 7

1
 CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC
 51
 CGTGAAAGTC AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA
 101
 TACACTGGGT TAGACAGGCC CCTGGCCAAA GGCTGGAGTG GATAGGAGGT
 151
 ATTAATCCTA ACAATGGTAT TCCTAACTAC AACCAGAAGT TCAAGGGCCG
 201
 GGCCACCTTG ACCGTAGGCA AGTCTGCCAG CACCGCCTAC ATGGAAGTGT
 251
 CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA
 301
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG
 351 372
 AACCTTGTC ACCGTCTCCT CA

Fig. 8

1	11	21	31	41
QVQLVQSGAE	VKKPGASVKV	SCKTSRYTFT	EYTIHWVRQA	PGQRLEWIGG
51	61	71	81	91
INPNNGIPNY	NQKFKGRATL	TVGKSASTAY	MELSSLRSED	TAVYYCARRR
101	111	121-124		
IAYGYDEGHA	MDYWGQGLV	TVSS		

Fig. 9

1
 CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC
 51
 CGTGAAAGTC AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA
 101
 TACACTGGGT TAGACAGGCC CCTGGCCAAA GGCTGGAGTG GATAGGAGGT
 151
 ATTAATCCTA ACAATGGTAT TCCTAACTAC AACCAGAAGT TCAAGGGCCG
 201
 GGCCACCTTG ACCGTAGGCA AGTCTGCCAG CACCGCCTAC ATGGAAGTGT
 251
 CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTTCTGCGC CAGAAGAAGA
 301
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG
 351 372
 AACCTTGTC ACCGTCTCCT CA

Fig. 10

1	11	21	31	41
QVQLVQSGAE	VKKPGASVKV	SCKTSRYTFT	EYTIHWVRQA	PGQRLEWIGG
51	61	71	81	91
INPNNGIPNY	NQKFKGRATL	TVGKSASTAY	MELSSLRSED	TAVYFCARRR
101	111	121-124		
IAYGYDEGHA	MDYWGQGTLV	TVSS		

Fig. 11

1				
CAGGTGCAAC	TAGTGCAGTC	CGGCGCCGAA	GTGAAGAAAC	CCGGTGCTTC
51				
CGTGAAAGTC	AGCTGTAAAA	CTAGTAGATA	CACCTTCACT	GAATACACCA
101				
TAACTGGGT	TAGACAGGCC	CCTGGCCAAA	GGCTGGAGTG	GATAGGAGGT
151				
ATTAATCCTA	ACAATGGTAT	TCCTAACTAC	AACCAGAAAGT	TCAAGGGCCG
201				
GGTCACCATC	ACCGTAGACA	CCTCTGCCAG	CACCGCCTAC	ATGGAAGTGT
251				
CCAGCCTGCG	CTCCGAGGAC	ACTGCAGTCT	ACTACTGCGC	CAGAAGAAGA
301				
ATCGCCTATG	GTTACGACGA	GGGCCATGCT	ATGGACTACT	GGGGTCAAGG
351		372		
AACCCTTGTC	ACCGTCTCCT	CA		

Fig. 12

1	11	21	31	41
QVQLVQSGAE	VKKPGASVKV	SCKTSRYTFT	EYTIHWVRQA	PGQRLEWIGG
51	61	71	81	91
INPNNGIPNY	NQKFKGRVTI	TVDTASASTAY	MELSSLRSED	TAVYYCARRR
101	111	121-124		
IAYGYDEGHA	MDYWGQGTLV	TVSS		

Fig. 13

1
 CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC
 51
 CGTGAAAGTC AGCTGTAAAA CTAGTAGATA CACCTTCACT GAATACACCA
 101
 TACACTGGGT TAGACAGGCC CCTGGCCAAA GGCTGGAGTG GATAGGAGGT
 151
 ATTAATCCTA ACAATGGTAT TCCTAACTAC AACCAGAAGT TCAAGGGCCG
 201
 GGTCACCAATC ACCGTAGACA CCTCTGCCAG CACCGCCTAC ATGGAAGTGT
 251
 CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTTCTGCGC CAGAAGAAGA
 301
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG
 351 372
 AACCCCTGTC ACCGTCTCCT CA

Fig. 14

1	11	21	31	41
QVQLVQSGAE	VKKPGASVKV	SCKTSRYTFT	EYTIHWVRQA	PGORLEWIGG
51	61	71	81	91
INPNNGIPNY	NQKFKGR <u>VTI</u>	TV <u>D</u> TSASTAY	MELSSLRSED	TAVY <u>F</u> CARRR
101	111	121-124		
IAYGYDEGHA	MDYWGQGLV	TVSS		

Fig. 15

1
 CAGGTGCAAC TAGTGCAGTC CGGCGCCGAA GTGAAGAAAC CCGGTGCTTC
 51
 CGTGAAAGTC AGCTGTAAAA CTAGTGGATA CACCTTCACT GAATACACCA
 101
 TACACTGGGT TAGACAGGCC CCTGGCCAAA GGCTGGAGTG GATAGGAGGT
 151
 ATTAATCCTA ACAATGGTAT TCCTAACTAC AACCAGAAGT TCAAGGGCCG
 201
 GGTCACCAATC ACCGTAGACA CCTCTGCCAG CACCGCCTAC ATGGAAGTGT
 251
 CCAGCCTGCG CTCCGAGGAC ACTGCAGTCT ACTACTGCGC CAGAAGAAGA
 301
 ATCGCCTATG GTTACGACGA GGGCCATGCT ATGGACTACT GGGGTCAAGG
 351 372
 AACCCCTGTC ACCGTCTCCT CA

Fig. 16

1	11	21	31	41
QVQLVQSGAE	VKKPGASVKV	SCKTSGYTFT	EYTIHWVRQA	PGQRLEWIGG
51	61	71	81	91
INPNNGIPNY	NQKFKGRVTI	TVDTASASTAY	MELSSLRSED	TAVYYCARRR
101	111	121-124		
IAYGDEGHA	MDYWGQGLV	TVSS		

Fig. 17

1
DIVMSQSPSS LAVSVGEKVT MSCKSSQSLL YSRNQKNYLA WFQOKPGQSP
51
KLLIFWASTR ESGVPDRFTG SGFGTDFNLT ISSVQAEDLA VYDCQQYFSY
101
PLTFGAGTKL ELKRTVAAPS VFIFPPSDEQ LKSGTASVVC LLNNFYPREA
151
KVQWKVDNAL QSGNSQESVT EQDSKDSTYS LSSTLTLSKA DYEKHKVYAC
201
EVTHQGLSSP VTKSFNRGEC

Fig. 18

1
VQLQSGPEL VKPGASVKMS CKTSRYTFTE YTIHWVRQSH GKSLEWIGGI
51
NPNNGIPNYN QKFKGRATLT VGKSSSTAYM ELRSLTSEDS AVYFCARRRI
101
AYGYDEGHAM DYWGQGTSVT VSSASTKGPS VFPLAPSSKS TSGGTAALGC
151
LVKDYFPEPV TVSWNSGALT SGVHTFPAVL QSSGLYSLSS VVTVPSSSLG
201
TQTYICNVNH KPSNTKVDKK VEPKSCDKTH TCPPCPAPEL LGGPSVFLFP
251
PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV HNAKTKPREE
301
QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR
351
EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT
401
PPVLDSGGSF FLYSKLTVDK SRWQQGNVFS CSVMHEALHN HYTQKSLSL
451
PGK

Fig. 19

340	350	360	370	380
CGTACTGTGG	CTGCACCATC	TGTCTTCATC	TTCCCGCCAT	CTGATGAGCA
390	400	410	420	430
GTTGAAATCT	GGAAGTGCCT	CTGTTGTGTG	CCTGCTGAAT	AACTTCTATC
440	450	460	470	480
CCAGAGAGGC	CAAAGTACAG	TGGAAGGTGG	ATAACGCCCT	CCAATCGGGT
490	500	510	520	530
AACTCCCAGG	AGAGTGTCAC	AGAGCAGGAC	AGCAAGGACA	GCACCTACAG
540	550	560	570	580
CCTCAGCAGC	ACCCTGACGC	TGAGCAAAGC	AGACTACGAG	AAACACAAAG
590	600	610	620	630
TCTACGCCTG	CGAAGTCACC	CATCAGGGCC	TGAGCTCGCC	CGTCACAAAG
640	650	660		
AGCTTCAACA	GGGGAGAGTGT			

Fig. 20

114	124	134	144	154
RTVAAPSVFI	FPPSDEQLKS	GTASVVCLLN	NFYPPREKVQ	WKVDNALQSG
164	174	184	194	204
NSQESVTEQD	SKDSTYSLSS	TLTLSKADYE	KHKVYACEVT	HQGLSSPVTK
214-220				
SFNRGEC				

Fig. 21

373
 GCCTCCACCA AGGGCCCATC GGTCTTCCCC CTGGCACCCCT CCTCCAAGAG
 423
 CACCTCTGGG GGCACAGCGG CCCTGGGCTG CCTGGTCAAG GACTACTTCC
 473
 CCGAACCGGT GACGGTGTCTG TGGAACTCAG GCGCCCTGAC CAGCGGCGTG
 523
 CACACCTTCC CGGCTGTCCT ACAGTCCTCA GGACTCTACT CCCTCAGCAG
 573
 CGTGGTGACC GTGCCCTCCA GCAGCTTGGG CACCCAGACC TACATCTGCA
 623
 ACGTGAATCA CAAGCCCAGC AACACCAAGG TGGACAAGAA AGTTGAGCCC
 673
 AAATCTTGTG ACAAAACTCA CACATGCCCA CCGTGCCCAG CACCTGAACT
 723
 CCTGGGGGGA CCGTCAGTCT TCCTCTTCCC CCCAAAACCC AAGGACACCC
 773
 TCATGATCTC CCGGACCCCT GAGGTCACAT GCGTGGTGGT GGACGTGAGC
 823
 CACGAAGACC CTGAGGTCAA GTTCAACTGG TACGTGGACG GCGTGGAGGT
 873
 GCATAATGCC AAGACAAAGC CGCGGGAGGA GCAGTACAAC AGCACGTACC
 923
 GGGTGGTCAG CGTCCTCACC GTCCTGCACC AGGACTGGCT GAATGGCAAG
 973
 GAGTACAAGT GCAAGGTCTC CAACAAAGCC CTCCCAGCCC CCATCGAGAA
 1023
 AACCATCTCC AAAGCCAAAG GGCAGCCCCG AGAACCACAG GTGTACACCC
 1073
 TGCCCCCATC CCGGGAGGAG ATGACCAAGA ACCAGGTCAG CCTGACCTGC
 1123
 CTGGTCAAAG GCTTCTATCC CAGCGACATC GCCGTGGAGT GGGAGAGCAA
 1173
 TGGGCAGCCG GAGAACAAC TACAAGACCAC GCCTCCCGTG CTGGACTCCG
 1223
 ACGGCTCCTT CTTCTCTAC AGCAAGCTCA CCGTGGACAA GAGCAGGTGG
 1273
 CAGCAGGGGA ACGTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA
 1323 1362
 CCACTACACG CAGAAGAGCC TCTCCCTGTC TCCGGGTAAA

Fig. 22

125
 ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV
 175
 HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP
 225
 KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMIS RTP EVTCVVVDVS
 275
 HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK
 325
 EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC
 375
 LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW
 425 454
 QQGNVFSCSV MHEALHNHYT QKSLSLSPGK

Fig. 23A




Fig. 23B




Fig. 24

HindIII
 1 aagcttGCCGCCACCatggattcacaggcccaggttcttatgttactgccgctatgggta
 -----+-----+-----+-----+-----+-----+
ttcgaacGGCGGGTGGtacctaagtgtccgggtccaagaatacaatgacggcgatacccat
 Kozak sequence
 M D S Q A Q V L M L L P L W V

61 tctggtagctgtggggacattgtgatgtcacagtctccatcctccctagctgtgtcagtt
 -----+-----+-----+-----+-----+-----+
 agaccatggacacccctgtaacactacagtgtcagaggtaggagggatcgacacagtcaa
 S G T C G D I V M S Q S P S S L A V S V

121 ggagagaagggttactatgagctgcaagtccagtcagagccttttatatagtcgtaatcaa
 -----+-----+-----+-----+-----+-----+
 cctctcttccaatgatactcgacgttcaggtcagtcctcgaaaatatatcagcattagtt
 G E K V T M S C K S S Q S L L Y S R N Q
 CDR 1

181 aagaactacttggcctggttccagcagaagccagggcagtcctcctaaactgctgattttc
 -----+-----+-----+-----+-----+-----+
 ttcttgatgaaccggaccaaggtcgtcttcgggtcccgtcagaggatttgacgactaaaag
 K N Y L A W F Q Q K P G Q S P K L L I F

241 tgggcatccactaggggaatctgggggtccctgatcgcttcacaggcagtggttgggacg
 -----+-----+-----+-----+-----+-----+
 acccgtaggtgatcccttagaccccagggactagcgaagtgtccgtcacctaaaccctgc
 W A S T R E S G V P D R F T G S G F G T
 CDR 2

301 gatttcaatctcaccatcagcagtggtgcaggctgaggacctggcagtttatgactgtcag
 -----+-----+-----+-----+-----+-----+
 ctaaagttagagtggtagtcgtcacacgtccgactcctggaccgtcaaatactgacagtc
 D F N L T I S S V Q A E D L A V Y D C Q

361 caatatttttagctatccgctcacgttcggtgctgggaccaagctggagctgaAACGTGAG
 -----+-----+-----+-----+-----+-----+
 gttataaaatcgataggcgagtgcaagccacgaccctgggttcgacctcgactTTGCACTG
 splice donor site

Q Y F S Y P L T F G A G T K L E L K
 CDR 3

BamHI
 Tggatcc
 421 ----- 427
 Acctagg

Fig. 25

HindIII
AAGCTTGCCGCCACCATGGGATGGAGCTGGGTCTTTCTCTTTCTCCTGTCAGGAACTGCA
 1 -----+-----+-----+-----+-----+-----+-----+
TTCGAACGGCGGTGGTACCCTACCTCGACCCAGAAAGAGAAAGAGGACAGTCCTTGACGT
Kozak sequence
 M G W S W V F L F L L S G T A

GGTGTCTCTCTGAGGTCCAGCTGCAACAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCT
 61 -----+-----+-----+-----+-----+-----+-----+
 CCACAGGAGAGACTCCAGGTCGACGTTGTCAGACCTGGACTCGACCACTTCGGACCCCGA
 G V L S E V Q L Q Q S G P E L V K P G A

TCAGTAAAGATGTCCTGCAAGACTTCTAGATACACATTCAGTGAATACACCATACTGG
 121 -----+-----+-----+-----+-----+-----+-----+
 AGTCATTTCTACAGGACGTTCTGAAGATCTATGTGTAAGTGACTTATGTGGTATGTGACC
 S V K M S C K T S R Y T F T E Y T I H W
 CDR 1

GTGAGACAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAGGTATTAATCCTAACAATGGT
 181 -----+-----+-----+-----+-----+-----+-----+
 CACTCTGTCTCGGTACCTTTCTCGGAACTCACCTAACCTCCATAATTAGGATTGTTACCA
 V R Q S H G K S L E W I G G I N P N N G
 CDR 2

ATTCCTAACTACAACCAGAAGTTCAAGGGCAGGGCCACATTGACTGTAGGCAAGTCCTCC
 241 -----+-----+-----+-----+-----+-----+-----+
 TAAGGATTGATGTTAGTCTTCAAGTTCCCGTCCCGGTGTAAGTACATCCGTTCCAGGAGG
 I P N Y N Q K F K G R A T L T V G K S S

AGCACCGCCTACATGGAGCTCCGCAGCCTGACATCTGAGGATTCTGCGGTCTATTTCTGT
 301 -----+-----+-----+-----+-----+-----+-----+
 TCGTGGCGGATGTACCTCGAGGCGTCGGACTGTAGACTCCTAAGACGCCAGATAAAGACA
 S T A Y M E L R S L T S E D S A V Y F C

GCAAGAAGAAGAATCGCCTATGGTTACGACGAGGGCCATGCTATGGACTACTGGGGTCAA
 361 -----+-----+-----+-----+-----+-----+-----+
 CGTTCTTCTTCTTAGCGGATACCAATGCTGCTCCCGGTACGATACCTGATGACCCCACTT
 A R R R I A Y G Y D E G H A M D Y W G Q
 CDR 3

BamHI
 GGAACCTCAGTCACCGTCTCCTCAGGTGAGTGGATCC
 421 -----+-----+-----+-----+-----+-----+-----+ 468
 CCTTGAGTCACTGGCAGAGGAGTCCACTCACCTAGG
 splice donor site
 G T S V T V S S

Fig. 26 /1

1 gaattccagc acactggcgg ccggttACTAG TTATTAATAG TAATCAATTA
 Spe I
 51 CGGGGTCATT AGTTCATAGC CCATATATGG AGTTCCGCGT TACATAACTT
 101 ACGGTAAATG GCCCGCCTGG CTGACCGCCC AACGACCCCC GCCCATTGAC
 151 GTCAATAATG ACGTATGTTC CCATAGTAAC GCCAATAGGG ACTTTCATT
 201 GACGTCAATG GGTGGAGTAT TTACGGTAAA CTGCCCACTT GGCAGTACAT
 251 CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA ATGACGGTAA
 301 ATGGCCCGCC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA
 SnaB I
 351 CTTGGCAGTA CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG
 401 TTTTGGCAGT ACATCAATGG GCGTGGATAG CGGTTTGACT CACGGGGATT
 451 TCCAAGTCTC CACCCCATTG ACGTCAATGG GAGTTTGTTT TGGCACCAAA
 501 ATCAACGGGA CTTTCCAAAA TGTCGTAACA ACTCCGCCCC ATTGACGCAA
 551 ATGGGCGGTA GGCGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT
 601 AGTGAACCGT CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC
 Sac II
 651 ATAGAAGACA CCGGGACCGA TCCAGCCTCC GCGGCCGGA ACGGTGCATT
 701 GGAACGCGGA TTCCCCGTGC CAAGAGTGAC GTAAGTACCG CCTATAGAGT
 751 CTATAGGCCC ACCCCCTTGG CTTCTTATGC ATGCTATACT GTTTTTGGCT
 801 TGGGGTCTAT ACACCCCCGC TTCCTCATGT TATAGGTGAT GGTATAGCTT
 851 AGCCTATAGG TGTGGGTTAT TGACCATTAT TGACCACTCC CCTATTGGTG
 901 ACGATACTTT CCATTACTAA TCCATAACAT GGCTCTTTGC CACAACCTCT
 951 TTTATTGGCT ATATGCCAAT ACACTGTCCT TCAGAGACTG ACACGGACTC
 1001 TGTATTTTAA CAGGATGGGG TCTCATTTAT TATTTACAAA TTCACATATA
 1051 CAACACCACC GTCCCCAGTG CCCGCAGTTT TTATTAAACA TAACGTGGGA
 BspE I
 1101 TCTCCACGCG AATCTCGGGT ACGTGTTCCG GACATGGGCT CTTCTCCGGT
 1151 AGCGGCGGAG CTTCTACATC CGAGCCCTGC TCCCATGCCT CCAGCGACTC
 1201 ATGGTCGCTC GGCAGCTCCT TGCTCCTAAC AGTGGAGGCC AGACTTAGGC
 1251 ACAGCACGAT GCCCACCACC ACCAGTGTGC CGCACAAGGC CGTGGCGGTA

Fig. 26 /3

2351 TGAGCAAAGC AGACTACGAG AAACACAAAG TCTACGCCTG CGAAGTCACC
 L S K A D Y E K H K V Y A C E V T
 2401 CATCAGGGCC TGAGCTCGCC CGTCACAAAG AGCTTCAACA GGGGAGAGTG
 H Q G L S S P V T K S F N R G E C
 2451 TTAGAGGGGAG AAGTGCCCCC ACCTGCTCCT CAGTTCCAGC CTGACCCCTT
 *
 2501 CCCATCCTTT GGCCTCTGAC CCTTTTCCA CAGGGGACCT ACCCCTATTG
 2551 CGGTCCTCCA GTCATCTTT CACCTCACCC CCCTCCTCCT CCTTGGCTTT
 2601 AATTATGCTA ATGTTGGAGG AGAATGAATA AATAAAGTGA ATCTTTGCAC
 2651 CTGTGGTGGG TCTAATAAAA GATATTTATT TTCATTAGAT ATGTGTGTTG
 2701 GTTTTTTTGTG TGCAGTGCCT CTATCTGGAG GCCAGGTAGG GCTGGCCTTG
 2751 GGGGAGGGGG AGGCCAGAAT GACTCCAAGA GCTACAGGAA GGCAGGTCAG
 2801 AGACCCCACT GGACAAACAG TGGCTGGACT CTGCACCATA ACACACAATC
 2851 AACAGGGGAG TGAGCTGGAA ATTTGCTAGC GAATTCTTGA AGACGAAAGG
 2901 GCCTCGTGAT ACGCCTATTT TTATAGGTTA ATGTCATGAT AATAATGGTT
 2951 TCTTAGACGT CAGGTGGCAC TTTTCGGGGA AATGTGCGCG GAACCCCTAT
 3001 TTGTTTATTT TTCTAAATAC ATTCAAATAT GTATCCGCTC ATGAGACAAT
 3051 AACCCTGATA AATGCTTCAA TAATATTGAA AAAGGAAGAG TATGAGTATT
 3101 CAACATTTCC GTGTCGCCCT TATTCCTTTT TTTGCGGCAT TTTGCCTTCC
 3151 TGTTTTTGCT CACCCAGAAA CGCTGGTGAA AGTAAAAGAT GCTGAAGATC
 3201 AGTTGGGTGC ACGAGTGGGT TACATCGAAC TGGATCTCAA CAGCGGTAAG
 3251 ATCCTTGAGA GTTTTCGCCC CGAAGAACGT TTTCCAATGA TGAGCACTTT
 3301 TAAAGTTCTG CTATGTGGCG CGGTATTATC CCGTGTTGAC GCCGGGCAAG
 3351 AGCAACTCGG TCGCCGCATA CACTATTCTC AGAATGACTT GGTGAGTAC
 3401 TCACCAGTCA CAGAAAAGCA TCTTACGGAT GGCATGACAG TAAGAGAATT
 3451 ATGCAGTGCT GCCATAACCA TGAGTGATAA CACTGCGGCC AACTTACTTC
 Pvu I
 3501 TGACAACGAT CGGAGGACCG AAGGAGCTAA CCGCTTTTTT GCACAACATG
 3551 GGGGATCATG TAACTCGCCT TGATCGTTGG GAACCGGAGC TGAATGAAGC
 3601 CATACCAAAC GACGAGCGTG ACACCACGAT GCCTGCAGCA ATGGCAACAA

Fig. 26 /4

3651 CGTTGCGCAA ACTATTA ACT GGC GAACTAC T TACTCTAGC TTCCCGGCAA
 3701 CAATTAATAG ACTGGATGGA GCGGATAAA GTTGCAGGAC CACTTCTGCG
 3751 CTCGGCCCTT CCGGCTGGCT GGT TTATTGC TGATAAATCT GGAGCCGGTG
 3801 AGCGTGGGTC TCGCGGTATC ATTGCAGCAC TGGGGCCAGA TG GTAAGCCC
 3851 TCCCGTATCG TAGTTATCTA CACGACGGGG AGTCAGGCAA CTATGGATGA
 3901 ACGAAATAGA CAGATCGCTG AGATAGGTGC CTCACTGATT AAGCATTGGT
 3951 AACTGTCAGA CCAAGTTTAC TCATATATAC TTTAGATTGA TTTAAAACCT
 4001 CATTTTTAAT TTAAAAGGAT CTAGGTGAAG ATCCTTTTGT ATAATCTCAT
 4051 GACCAAAATC CCTTAACGTG AGTTTTCGTT CCACTGAGCG TCAGACCCCG
 4101 TAGAAAAGAT CAAAGGATCT TCTTGAGATC CTTTTTTTCT GCGCGTAATC
 4151 TGCTGCTTGC AAACAAAAAA ACCACCGCTA CCAGCGGTGG TTTGTTTGCC
 4201 GGATCAAGAG CTACCAACTC TTTTTCGAA GGTAAC TGGC TTCAGCAGAG
 4251 CGCAGATACC AAATACTGTC CTTCTAGTGT AGCCGTAGTT AGGCCACCAC
 4301 TTCAAGAACT CTGTAGCACC GCCTACATAC CTCGCTCTGC TAATCCTGTT
 4351 ACCAGTGGCT GCTGCCAGTG GCGATAAGTC GTGTCTTACC GGGTTGGACT
 4401 CAAGACGATA GTTACCGGAT AAGGCGCAGC GGTCGGGCTG AACGGGGGGT
 4451 TCGTGACAC AGCCAGCTT GGAGCGAACG ACCTACACCG AACTGAGATA
 4501 CCTACAGCGT GAGCTATGAG AAAGCGCCAC GCTTCCCGAA GGGAGAAAGG
 4551 CGGACAGGTA TCCGGTAAGC GGCAGGGTCG GAACAGGAGA GCGCACGAGG
 4601 GAGCTTCCAG GGGGAAACGC CTGGTATCTT TATAGTCCTG TCGGGTTTCG
 4651 CCACCTCTGA CTTGAGCGTC GATTTTTGTG ATGCTCGTCA GGGGGGCGGA
 4701 GCCTATGGAA AAACGCCAGC AACGCGGCCT TTTTACGGTT CCTGGCCTTT
 4751 TGCTGGCCTT TTGCTCCACAT GTTCTTTCCT GCGTTATCCC CTGATTCTGT
 4801 GGATAACCGT ATTACCGCCT TTGAGTGAGC TGATACCGCT CGCCGCAGCC
 4851 GAACGACCGA GCGCAGCGAG TCAGTGAGCG AGGAAGCGGA AGAGCGCCTG
 4901 ATGCGGTATT TTCTCCTTAC GCATCTGTGC GGTATTTTAC ACCGCATATG

BspLU11I

Fig. 26 /5

Bst1107I

4951 GTGCACTCTC AGTACAATCT GCTCTGATGC CGCATAGTTA AGCCAGTATA

5001 CACTCCGCTA TCGCTACGTG ACTGGGTCAT GGCTGCGCCC CGACACCCGC

5051 CAACACCCGC TGACGCGCCC TGACGGGCTT GTCTGCTCCC GGCATCCGCT

5101 TACAGACAAG CTGTGACCGT CTCCGGGAGC TGCATGTGTC AGAGGTTTTC

5151 ACCGTCATCA CCGAAACGCG CGAGGCAGCT GTGGAATGTG TGTCAGTTAG

5201 GGTGTGGAAA GTCCCCAGGC TCCCCAGCAG GCAGAAGTAT GCAAAGCATG

5251 CATCTCAATT AGTCAGCAAC CAGGCTCCCC AGCAGGCAGA AGTATGCAAA

5301 GCATGCATCT CAATTAGTCA GCAACCATAG TCCCGCCCCT AACTCCGCCC

5351 ATCCCGCCCC TAACTCCGCC CAGTTCCGCC CATTCTCCGC CCCATGGCTG

Sfi I

5401 ACTAATTTTT TTTATTTATG CAGAGGCCGA GGCCGCTCG GCCTCTGAGC

Stu I/Avr II

5451 TATTCCAGAA GTAGTGAGGA GGCTTTTTTG GAGGCCTAGG CTTTTCGAAA

5501 AAGCTAGCTT CACGCTGCCG CAAGCACTCA GGGCGCAAGG GCTGCTAAAG

5551 GAAGCGGAAC ACGTAGAAAG CCAGTCCGCA GAAACGGTGC TGACCCCGGA

5601 TGAATGTCAG CTACTGGGCT ATCTGGACAA GGGAAAACGC AAGCGCAAAG

5651 AGAAAGCAGG TAGCTTGAGG TGGGCTTACA TGGCGATAGC TAGACTGGGC

5701 GGTTTTATGG ACAGCAAGCG AACCGBAATT GCCAGCTGGG GCGCCCTCTG

5751 GTAAGGTTGG GAAGCCCTGC AAAGTAACT GGATGGCTTT CTGCGGCCA

Bgl II/Bcl I

5801 AGGATCTGAT GGCAGAGGGG ATCAAGATCT GATCAAGAGA CAGGATGAGG

5851 ATCGTTTCGC ATGATTGAAC AAGATGGATT GCACGCAGGT TCTCCGGCCG

5901 CTTGGGTGGA GAGGCTATTC GGCTATGACT GGGCACAACA GACAATCGGC

5951 TGCTCTGATG CCGCCGTGTT CCGGCTGTCA GCGCAGGGGC GCCCGGTTCT

6001 TTTTGTCAAG ACCGACCTGT CCGGTGCCCT GAATGAACTG CAGGACGAGG

Msc I

6051 CAGCGCGGCT ATCGTGGCTG GCCACGACGG GCGTTCCTTG CGCAGCTGTG

6101 CTCGACGTTG TCACTGAAGC GGAAGGGAC TGGCTGCTAT TGGGCGAAGT

6151 GCCGGGGCAG GATCTCCTGT CATCTCACCT TGCTCCTGCC GAGAAAGTAT

6201 CCATCATGGC TGATGCAATG CGGCGGCTGC ATACGCTTGA TCCGGCTACC

Fig. 26 /6

6251 TGCCCATTCG ACCACCAAGC GAAACATCGC ATCGAGCGAG CACGTACTCG

6301 GATGGAAGCC GGTCTTGTCG ATCAGGATGA TCTGGACGAA GAGCATCAGG

6351 GGCTCGCGCC AGCCGAACTG TTCGCCAGGC TCAAGGCGCG CATGCCCCGAC

6401 GGCGAGGATC TCGTCGTGAC CCATGGCGAT GCCTGCTTGC CGAATATCAT

6451 GGTGGA^{Rsr II}AAAT GGCCGCTTTT CTGGATTCAT CGACTGTGGC CGGCTGGGTG

6501 TGGCGGACCG CTATCAGGAC ATAGCGTTGG CTACCCGTGA TATTGCTGAA

6551 GAGCTTGGCG GCGAATGGGC TGACCGCTTC CTCGTGCTTT ACGGTATCGC

6601 CGCTCCCGAT TCGCAGCGCA TCGCCTTCTA TCGCCTTCTT GACGAGTTCT

6651 TCTGAGCGGG ACTCTGGGGT ^{Nsp V}TCGAATGAC CGACCAAGCG ACGCCCAACC

6701 TGCCATCACG AGATTTCGAT TCCACCGCCG CCTTCTATGA AAGGTTGGGC

6751 TTCGGAATCG TTTTCCGGGA CGCCGGCTGG ATGATCCTCC AGCGCGGGGA

6801 TCTCATGCTG GAGTTCTTCG CCCACCCCGG ^{Sma I}GCTCGATCCC ^{Nru I}CTCGCGAGTT

6851 GGTTCAGCTG CTGCCTGAGG CTGGACGACC TCGCGGAGTT CTACCGGCAG

6901 TGCAAATCCG TCGGCATCCA GGAAACCAGC AGCGGCTATC CGCGCATCCA

6951 TGCCCCCGAA CTGCAGGAGT GGGGAGGCAC GATGGCCGCT TTGGTCCCGG

7001 ATCTTTGTGA AGGAACCTTA CTTCTGTGGT GTGACATAAT TGGACAAACT

7051 ACCTACAGAG ATTTAAAGCT CTAAGGTAAA TATAAAATTT TTAAGTGTAT

7101 AATGTGTTAA ACTACTGATT CTAATTGTTT GTGTATTTTA GATTCCAACC

7151 TATGGA^{Rsr II}ACTG ATGAATGGGA GCAGTGGTGG AATGCCTTTA ATGAGGAAAA

7201 CCTGTTTTGC TCAGAAGAAA TGCCATCTAG TGATGATGAG GCTACTGCTG

7251 ACTCTCAACA TTCTACTCCT CCAAAAAAGA AGAGAAAGGT AGAAGACCCC

7301 AAGGACTTTC CTTCAGAATT GCTAAGTTTT TTGAGTCATG CTGTGTTTAG

7351 TAATAGA^{Rsr II}ACT CTTGCTTGCT TTGCTATTTA CACCACAAAG GAAAAAGCTG

7401 CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC CTTTATAAGT

7451 AGGCATAACA GTTATAATCA TAACATACTG TTTTTTCTTA CTCCACACAG

7501 GCATAGAGTG TCTGCTATTA ATA^{Rsr II}ACTATGC TCAAAAAATTG TGTACCTTTT

Fig. 26 /7

7551 GCTTTTTAAT TTGTAAAGGG GTTAATAAGG AATATTTGAT GTATAGTGCC
 7601 TTGACTAGAG ATCATAATCA GCCATACCAC ATTTGTAGAG GTTTTACTTG
 7651 CTTTAAAAAA CCTCCACAC CTCCCCCTGA ACCTGAAACA TAAATGAAT
 Mun I
 7701 GCAATTGTTG TTGTAACTT GTTTATTGCA GCTTATAATG GTTACAAATA
 7751 AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTTTT TCACTGCATT
 7801 CTAGTTGTGG TTTGTCCAAA CTCATCAATG TATCTTATCA TGTCTGGATC
 7851 TAATAAAAGA TATTTATTTT CATTAGATAT GTGTGTTGGT TTTTGTGTG
 7901 CAGTGCCTCT ATCTGGAGGC CAGGTAGGGC TGGCCTTGGG GGAGGGGGAG
 7951 GCCAGAAATGA CTCCAAGAGC TACAGGAAGG CAGGTCAGAG ACCCCACTGG
 8001 ACAACAGTG GCTGGACTCT GCACCATAAC ACACAATCAA CAGGGGAGTG
 8051 AGCTGGAAAT TTGCTAGC

Fig. 27/1

1 TTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAAT
 61 GGTTCCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTT
 121 ATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCT
 181 TCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTC
 241 CTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTAAA
 301 AGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGG
 361 TAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGT
 421 TCTGCTATGTGGCGCGGTATTATCCCGTGTTGACGCCGGGCAAGAGCAACTCGGTGCGCG
 481 CATACTATTCTCAGAATGACTTGTTGAGTACTCACCAGTCACAGAAAAGCATCTTAC
 541 GGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACTGCG
 Pvu I
 601 GGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAA
 661 CATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACC
 Fsp I
 721 AAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTTGCGCAAACTATT
 781 AACTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGA

Fig. 27 /2

841 TAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAA
 901 ATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAA
 961 GCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAA
 1021 TAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTGACACCAAGT
 1081 TTACTCATATATACTTTAGATTGATTTAAACTTCATTTTTTAATTTAAAGGATCTAGGT
 1141 GAAGATCCTTTTTGATAATCTCATGACCAAATCCCTTAACGTGAGTTTCGTTCCACTG
 1201 AGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTCTGCGCGT
 1261 AATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCA
 1321 AGAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATAC
 1381 TGTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTAC
 1441 ATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCT
 1501 TACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCCGGCTGAACGGG
 1561 GGGTTCGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACA
 1621 GCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGT
 1681 AAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTA
 1741 TCTTTATAGTCCTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTC
 1801 GTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTACGGTTCCTGGC
 1861 CTTTTGCTGGCCTTTTGCTCCACATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAA
 1921 CCGTATTACCGCCTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAG
 1981 CGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCTGATGCGGTATTTCTCCTTACGCATCT
 2041 GTGCGGTATTTACACCGCATATGGTGCCTCTCAGTACAATCTGCTCTGATGCCGCATA
 2101 GTTAAGCCAGTATACACTCCGCTATCGCTACGTGACTGGGTCATGGCTGCGCCCCGACAC
 2161 CCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGA
 2221 CAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTGTCAGAGGTTTTCACCGTCATCACCGAAA
 2281 CGCGCGAGGCAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACCTCGGCC
 2341 CATCCCGCCCCCTAACTCCGCCCAGTTCGCCCCATTCTCCGCCCCATGGCTGACTAATTTT
 2401 TTTTATTTATGCAGAGGCCGAGGGCCGCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGG
 2461 AGGCTTTTTTGGAGGCCTAGGCTTTTTGCAAAAAGCTAGCTTACAGCTCAGGGCTGCGATT

BspLU11I

Bst1107 I

Sfi I

Stu I/Avr II

Fig. 27 /3

2521 TCGCGCCAACTTGACGGCAATCCTAGCGTGAAGGCTGGTAGGATTTTATCCCCGCTGCC
 2581 ATCATGGTTCGACCATTGAACTGCATCGTCGCCGTGTCCCAAAATATGGGGATTGGCAAG
 2641 AACGGAGACCTACCCTGGCCTCCGCTCAGGAACGAGTTCAAGTACTTCCAAAGAATGACC
 2701 ACAACCTCTTCAGTGGAAGGTAAACAGAATCTGGTGATTATGGGTAGGAAAACCTGGTTC
 2761 TCCATTCCCTGAGAAGAATCGACCTTTAAAGGACAGAATTAATATAGTTCTCAGTAGAGAA
 2821 CTCAAAGAACCACCACGAGGAGCTCATTTTCTTGCCAAAAGTTTGGATGATGCCTTAAGA
 2881 CTTATTGAACAACCGGAATTGGCAAGTAAAGTAGACATGGTTTGGATAGTCGGAGGCAGT
 2941 TCTGTTTACCAGGAAGCCATGAATCAACCAGGCCACCTCAGACTCTTGTGACAAGGATC
 3001 ATGCAGGAATTTGAAAGTGACACGTTTTTCCCAGAAATTGATTTGGGGAAATATAAACTT
 3061 CTCCCAGAATACCCAGGCGTCCTCTCTGAGGTCCAGGAGGAAAAGGCATCAAGTATAAG
 3121 TTTGAAGTCTACGAGAAGAAAGACTAACAGGAAGATGCTTCAAGTTCTCTGCTCCCCTC
 3181 CTAAAGCTATGCATTTTATAAGACCATGGGACTTTTGCTGGCTTTAGATCTTTGTGAAG
 3241 GAACCTTACTTCTGTGGTGTGACATAATTGGACAACTACCTACAGAGATTTAAAGCTCT
 3301 AAGGTAAATATAAAATTTTAAAGTGTATAATGTGTAACTACTGATTCTAATTGTTTGT
 3361 GTATTTTAGATTCCAACCTATGGAAGTATGAATGGGAGCAGTGGTGGAAATGCCTTTAAT
 3421 GAGGAAAACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGAC
 3481 TCTCAACATTCTACTCCTCCAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCT
 3541 TCAGAATTGCTAAGTTTTTTGAGTCATGCTGTGTTTAGTAATAGAACTCTTGCTTGCTTT
 3601 GCTATTTACACCACAAAGGAAAAGCTGCACTGCTATACAAGAAAATTATGGAAAATAT
 3661 TCTGTAACCTTTATAAGTAGGCATAACAGTTATAATCATAACTACTGTTTTTCTTACT
 3721 CCACACAGGCATAGAGTGTCTGCTATTAATAACTATGCTCAAAAATTGTGTACCTTTAGC
 3781 TTTTAAATTTGTAAAGGGTTAATAAGGAATATTTGATGTATAGTGCCTTGACTAGAGAT
 BsaB I
 3841 CATAATCAGCCATACCACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTCCCACACCT
 Mun I
 3901 CCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTAACTTGTTTATTGCAGC
 3961 TTATAATGGTTACAAATAAAGCAATAGCATCACAAATTCACAAATAAAGCATTTTTTTC
 4021 ACTGCATTCTAGTTGTGGTTTGTCCAACTCATCAATGTATCTTATCATGTCTGGATCTA
 4081 ATAAAAGATATTTATTTTCATTAGATATGTGTGTTGGTTTTTTGTGTGCAGTGCCTCTAT
 4141 CTGGAGGCCAGGTAGGGCTGGCCTTGGGGGAGGGGAGGCCAGAATGACTCCAAGAGCTA
 4201 CAGGAAGGCAGGTCAGAGACCCCACTGGACAAACAGTGGCTGGACTCTGCACCATAACAC

3

Fig. 27 /5

(Pvu II)

5941 GTGTCCTCTCTGAGGTCCAGCTGCAACAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTT
 G V L S E V Q L Q Q S G P E L V K P G A
 Xba I Dra III

6001 CAGTAAAGATGTCCTGCAAGACTTCTAGATACACATTCAGTGAATACACCATACTGGG
 S V .K M S C K T S R Y T F T E Y T I H W
 CDR 1

6061 TGAGACAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAGGTATTAATCCTAACAAATGGTA
 V R Q S H G K S L E W I G G I N P N N G
 6121 TTCCTAACTACAACCAGAAGTTCAAGGGCAGGGCCACATTGACTGTAGGCAAGTCCTCCA
 I P N Y N Q K F K G R A T L T V G K S S
 CDR 2

6181 GCACCGCCTACATGGAGCTCCGCAGCCTGACATCTGAGGATTCTGCGGTCTATTTCTGTG
 S T A Y M E L R S L T S E D S A V Y F C

6241 CAAGAAGAAGAATCGCCTATGGTTACGACGAGGGCCATGCTATGGACTACTGGGGTCAAG
 A R R R I A Y G Y D E G H A M D Y W G Q
 CDR 3 BamH I

6301 GAACCTCAGTCACCGTCTCCTCAGGTGAGTGGATCCTCTGCGCCTGGGCCAGCTCTGTC
 G T S V T V S S

6361 CCACACCGCGGTACATGGCACCACTCTCTTGACGCTCCACCAAGGGCCCATCGGTCT
 S T K G P S V

6421 TCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGG
 F P L A P S S K S T S G G T A A L G C L
 Age I

6481 TCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTTGAACTCAGGCGCCCTGACCAGCG
 V K D Y F P E P V T V S W N S G A L T S

6541 GCGTGACACCTTCCCGGCTGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGG
 G V H T F P A V L Q S S G L Y S L S S V
 BstE II

6601 TGACCGTGGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGC
 V T V P S S S L G T Q T Y I C N V N H K

6661 CCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAATCTTGTGACAAAACCTCACACAT
 P S N T K V D K K V E P K S C D K T H T

6721 GCCCACCCTGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCCCCCAA
 C P P C P A P E L L G G P S V F L F P P

6781 AACCCAAGGACACCCTCATGATCTCCCGACCCCTGAGGTCACATGCGTGGTGGTGGACG
 K P K D T L M I S R T P E V T C V V V D

6841 TGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA
 V S H E D P E V K F N W Y V D G V E V H

6901 ATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCC
 N A K T K P R E E Q Y N S T Y R V V S V

6961 TCACCGTCTGTCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA
 L T V L H Q D W L N G K E Y K C K V S N

Fig. 27 /6

7021 AAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC
 K A L P A P I E K T I S K A K G Q P R E
 7081 CACAGGTGTACACCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGA
 P Q V Y T L P P S R E E M T K N Q V S L
 7141 CCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGC
 T C L V K G F Y P S D I A V E W E S N G
 7201 AGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTCTTCC
 Q P E N N Y K T T P P V L D S D G S F F
 7261 TCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCT
 L Y S K L T V D K S R W Q Q G N V F S C
 7321 CCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
 S V M H E A L H N H Y T Q K S L S L S P
 NgoM I
 7381 GTAAATGAGTGCGACGGCCGGCAAGCCCCGCTCCCCGGGCTCTCGCGGTTCGACGAGGAT
 G K *
 7441 GCTTGGCACGTACCCCCTGTACATACTTCCCGGGCGCCAGCATGGAAATAAAGCACCGG
 7501 ATCTAATAAAAGATATTTATTTTCATTAGATATGTGTGTTGGTTTTTTTGTGTGCAGTGCC
 7561 TCTATCTGGAGGCCAGGTAGGGCTGGCCTTGGGGGAGGGGGAGGCCAGAATGACTCCAAG
 7621 AGCTACAGGAAGGCAGGTCAGAGACCCCACTGGACAAACAGTGGCTGGACTCTGCACCAT
 7681 AACACACAATCAACAGGGGAGTGAGCTGGaaatttgctagcgaattaattc 7731

Fig. 28:




Fig. 29 /1

	1	D	I	V	M	T	Q	S	P	D	S	L	A	V	S	L	G	E	R	19
A		GAC	ATT	GTG	ATG	ACC	CAA	TCT	CCA	GAC	TCT	TTG	GCT	GTG	TCT	CTA	GGG	GAG	AGG	GCC
B	
C		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

	20	T	I	N	C	CDR1	27	A	B	C	D	E	F	28		32				
A		ACC	ATC	AAC	TGC	AAG	TCC	AGT	CAG	AGC	CTT	TTA	TAT	TCT	AGA	AAT	CAA	AAG	AAC	TAC
B	
C		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

	33	L	A	W	Y	Q	Q	K	P	G	Q	P	P	K	L	L	I	F	51	
A		TTG	GCC	TGG	TAT	CAG	CAG	AAA	CCA	GGA	CAG	CCA	CCC	AAA	CTC	CTC	ATC	TTT	TGG	GCT
B		.	.	.	F
C		---	---	---	TC	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

	52	S	T	R	E	S	CDR2	G	V	P	D	R	F	S	G	S	G	F	G	T	D	70
A		AGC	ACT	AGG	GAA	TCT	GGG	GTA	CCT	GAT	AGG	TTC	AGT	GGC	AGT	GGG	TTT	GGG	ACA	GAC		
B		
C		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	

	71	F	T	L	T	I	S	S	L	Q	A	E	D	V	A	V	Y	Y	C	88
A		TTC	ACC	CTC	ACC	ATT	AGC	AGC	CTG	CAG	GCT	GAA	GAT	GTG	GCA	GTT	TAT	TAC	TGT	
B		D	G	
C		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	

	89	Q	Q	Y	F	S	Y	P	L	T	CDR3	F	G	Q	G	T	K	V	E	I	K	107
A		CAG	CAA	TAT	TTT	AGC	TAT	CCG	CTC	ACG	TTC	GGA	CAA	GGG	ACC	AAG	GTG	GAA	ATA	AAA		
B		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
C		---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	

Fig. 30 /1

1 gaattccagc acactggcgg ccgtACTAG TTATTAATAG TAATCAATTA
 Spe I
 51 CGGGGTCATT AGTTCATAGC CCATATATGG AGTCCGCGT TACATAACTT
 101 ACGGTAAATG GCCCGCCTGG CTGACCGCCC AACGACCCCC GCCCATTGAC
 151 GTCAATAATG ACGTATGTTT CCATAGTAAC GCCAATAGGG ACTTTCCATT
 201 GACGTCAATG GGTGGAGTAT TTACGGTAAA CTGCCCACTT GGCAGTACAT
 251 CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA ATGACGGTAA
 301 ATGGCCCGCC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA
 351 CTTGGCAGTA CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG
 SnaB I
 401 TTTTGGCAGT ACATCAATGG GCGTGGATAG CGGTTTGAAT CACGGGGATT
 451 TCCAAGTCTC CACCCCATTTG ACGTCAATGG GAGTTTGTTT TGGCACCAAA
 501 ATCAACGGGA CTTTCCAAAA TGTCGTAACA ACTCCGCCCC ATTGACGCAA
 551 ATGGGCGGTA GGCCTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT
 601 AGTGAACCGT CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC
 651 ATAGAAGACA CCGGGACCGA TCCAGCCTCC GCGGCCGGGA ACGGTGCATT
 Sac II
 701 GGAACGCGGA TTCCCCGTGC CAAGAGTGAC GTAAGTACCG CCTATAGAGT
 751 CTATAGGCCC ACCCCCTTGG CTTCTTATGC ATGCTATACT GTTTTTGGCT
 801 TGGGGTCTAT ACACCCCCGC TTCCTCATGT TATAGGTGAT GGTATAGCTT
 851 AGCCTATAGG TGTGGGTTAT TGACCATTAT TGACCACTCC CCTATTGGTG
 901 ACGATACTTT CCATTACTAA TCCATAACAT GGCTCTTTGC CACAACCTC
 951 TTTATTGGCT ATATGCCAAT AACTGTCCT TCAGAGACTG ACACGGACTC
 1001 TGTATTTTTA CAGGATGGGG TCTCATTTAT TATTTACAAA TTCACATATA
 1051 CAACACCACC GTCCCCAGTG CCCGCAGTTT TTATTAAACA TAACGTGGGA
 1101 TCTCCACGCG AATCTCGGGT ACGTGTTCCG GACATGGGCT CTTCTCCGGT
 (BspE I)
 1151 AGCGGCGGAG CTTCTACATC CGAGCCCTGC TCCCATGCCT CCAGCGACTC
 1201 ATGGTCGCTC GGCAGCTCCT TGCTCCTAAC AGTGGAGGCC AGACTTAGGC

Fig. 30 /2

2151 ACAGCACGAT GCCCACCACC ACCAGTGTGC CGCACAAGGC CGTGCGGTA
1301 GGGTATGTGT CTGAAAATGA GCTCggggag cgggcttgca ccgctgacgc
Afl II
1351 atttgaaga cttaaggcag cggcagaaga agatgcaggc agctgagtg
1401 ttgtgtctg ataagagtca gaggtaactc ccgttgccgt gctgttaacg
1451 gtggagggca gtgtagtctg agcagtactc gttgctccg cgcgcgccac
1501 cagacataat agctgacaga ctaacagact gttcctttcc atgggtcttt
Mlu I Hind III
1551 tctgcagtca ccgtcctga cacgcgtctc gggaagcttG CCGCCACCAT
M
1601 GGAGACAGAC ACACTCCTGC TATGGGTGCT GCTGCTCTGG GTTCCAGGTT
E T D T L L L W V L L L W V P G
(BspE I)
1651 CCTCCGGAGA CATTGTGATG ACCCAATCTC CAGACTCTTT GGCTGTGTCT
S S G D I V M T Q S P D S L A V S
1701 CTAGGGGAGA GGGCCACCAT CAACTGCAAG TCCAGTCAGA GCCTTTTATA
L G E R A T I N C K S S Q S L L Y
XbaI CDR 1
1751 TTCTAGAAAT CAAAAGAACT ACTTGGCCTG GTATCAGCAG AAACCAGGAC
S R N Q K N Y L A W Y Q Q K P G
KpnI
1801 AGCCACCCAA ACTCCTCATC TTTTGGGCTA GCACTAGGGA ATCTGGGGTA
Q P P K L L I F W A S T R E S G V
CDR 2
1851 CCTGATAGGT TCAGTGGCAG TGGGTTTGGG ACAGACTTCA CCCTCACCAT
P D R F S G S G F G T D F T L T I
1901 TAGCAGCCTG CAGGCTGAAG ATGTGGCAGT TTATTACTGT CAGCAATATT
S S L Q A E D V A V Y Y C Q Q Y
1951 TTAGCTATCC GCTCACGTTT GGACAAGGGA CCAAGGTGGA AATAAAACGT
F S Y P L T F G Q G T K V E I K R
CDR 3
BamH I
2001 GAGTggatcc ATCTGGGATA AGCATGCTGT TTTCTGTCTG TCCCTAACAT
2051 GCCCTGTGAT TATGCGCAAA CAACACACCC AAGGGCAGAA CTTTGTTACT
2101 TAAACACCAT CCTGTTIGCT TCTTTCCTCA GGAACTGTGG CTGCACCATC
T V A A P S
2151 TGTCTTCATC TTCCCGCCAT CTGATGAGCA GTTGAAATCT GGAAGTGCCT
V F I F P P S D E Q L K S G T A
2201 CTGTTGTGTG CCTGCTGAAT AACTTCTATC CCAGAGAGGC CAAAGTACAG
S V V C L L N N F Y P R E A K V Q
2251 TGGAAGGTGG ATAACGCCCT CCAATCGGGT AACTCCCAGG AGAGTGTCAC
W K V D N A L Q S G N S Q E S V T
2301 AGAGCAGGAC AGCAAGGACA GCACCTACAG CCTCAGCAGC ACCCTGACGC

Fig. 30 /3

E Q D S K D S T Y S L S S T L T
 2351 TGAGCAAAGC AGACTACGAG AAACACAAAG TCTACGCCTG CGAAGTCACC
 L S K A D Y E K H K V Y A C E V T
 2401 CATCAGGGCC TGAGCTCGCC CGTCACAAAG AGCTTCAACA GGGGAGAGTG
 H Q G L S S P V T K S F N R G E C
 2451 TTAGAGGGAG AAGTGCCCCC ACCTGCTCCT CAGTTCCAGC CTGACCCCCCT
 Psp5 II
 2501 CCCATCCTTT GGCCTCTGAC CCTTTTTCCA CAGGGGACCT ACCCCTATTG
 2551 CGGTCCTCCA GCTCATCTTT CACCTCACCC CCCTCCTCCT CCTTGGCTTT
 2601 AATTATGCTA ATGTTGGAGG AGAATGAATA AATAAAGTGA ATCTTTGCAC
 2651 CTGTGGTGGA TCTAATAAAA GATATTTATT TTCATTAGAT ATGTGTGTTG
 2701 GTTTTTTG TG CAGTGCCT CTATCTGGAG GCCAGGTAGG GCTGGCCTTG
 2751 GGGGAGGGGG AGGCCAGAAT GACTCCAAGA GCTACAGGAA GGCAGGTCAG
 2801 AGACCCCACT GGACAAACAG TGGCTGGACT CTGCACCATA ACACACAATC
 2851 AACAGGGGAG TGAGCTGGAA ATTTGCTAGC GAATTCTTGA AGACGAAAGG
 2901 GCCTCGTGAT ACGCCTATTT TTATAGGTTA ATGTCATGAT AATAATGGTT
 2951 TCTTAGACGT CAGGTGGCAC TTTTCGGGGA AATGTGCGCG GAACCCCTAT
 3001 TTGTTTATTT TTCTAAATAC ATTCAAATAT GTATCCGCTC ATGAGACAAT
 3051 AACCTTGATA AATGCTTCAA TAATATTGAA AAAGGAAGAG TATGAGTATT
 3101 CAACATTTCC GTGTCGCCCT TATTCCTTT TTTGCGGCAT TTTGCCTTCC
 3151 TGTTTTTGCT CACCCAGAAA CGCTGGTGAA AGTAAAAGAT GCTGAAGATC
 3201 AGTTGGGTGC ACGAGTGGGT TACATCGAAC TGGATCTCAA CAGCGGTAAG
 3251 ATCCTTGAGA GTTTTCGCCC CGAAGAACGT TTTCCAATGA TGAGCACTTT
 3301 TAAAGTTCTG CTATGTGGCG CGGTATTATC CCGTGTTGAC GCCGGGCAAG
 3351 AGCAACTCGG TCGCCGCATA CACTATTCTC AGAATGACTT GGTGAGTAC
 3401 TCACCAGTCA CAGAAAAGCA TCTTACGGAT GGCATGACAG TAAGAGAATT
 3451 ATGCAGTGCT GCCATAACCA TGAGTGATAA CACTGCGGCC AACTTACTTC
 Pvu I
 3501 TGACAACGAT CGGAGGACCG AAGGAGCTAA CCGCTTTTTT GCACAACATG
 3551 GGGGATCATG TAACTCGCCT TGATCGTTGG GAACCGGAGC TGAATGAAGC

Fig. 30 /4

3601 CATACCAAAC GACGAGCGTG ACACCACGAT GCCTGCAGCA ATGGCAACAA
 3651 CGTTGCGCAA ACTATTAAC TGGCAACTAC TTA CTCTAGC TTCCCGGCAA
 3701 CAATTAATAG ACTGGATGGA GGC GGATAAA GTTGCAGGAC CACTTCTGCG
 3751 CTCGGCCCTT CCGGCTGGCT GGT TTATTGC TGATAAATCT GGAGCCGGTG
 3801 AGCGTGGGTC TCGCGGTATC ATTGCAGCAC TGGGGCCAGA TGGTAAGCCC
 3851 TCCCGTATCG TAGTTATCTA CACGACGGGG AGTCAGGCAA CTATGGATGA
 3901 ACGAAATAGA CAGATCGCTG AGATAGGTGC CTCACTGATT AAGCATTGGT
 3951 AACTGTCAGA CCAAGTTTAC TCATATATAC TTTAGATTGA TTTAAACTT
 4001 CATTTTAAAT TTAAGGAT CTAGGTGAAG ATCCTTTTG ATAATCTCAT
 4051 GACCAAAATC CCTTAACGTG AGTTTTCGTT CCACTGAGCG TCAGACCCCG
 4101 TAGAAAAGAT CAAAGGATCT TCTTGAGATC CTTTTTTTCT GCGCGTAATC
 4151 TGCTGCTTGC AAACAAAAAA ACCACCGCTA CCAGCGGTGG TTTGTTGCC
 4201 GGATCAAGAG CTACCAACTC TTTTCCGAA GGTAAGTGGC TTCAGCAGAG
 4251 CGCAGATACC AAATACTGTC CTTCTAGTGT AGCCGTAGTT AGGCCACCAC
 4301 TTCAAGAACT CTGTAGCACC GCCTACATAC CTCGCTCTGC TAATCCTGTT
 4351 ACCAGTGGCT GCTGCCAGTG GCGATAAGTC GTGTCTTACC GGGTTGGA CT
 4401 CAAGACGATA GTTACCGGAT AAGGCGCAGC GGTCGGGCTG AACGGGGGGT
 4451 TCGTGACAC AGCCCAGCTT GGAGCGAACG ACCTACACCG AACTGAGATA
 4501 CCTACAGCGT GAGCTATGAG AAAGCGCCAC GCTTCCCGAA GGGAGAAAGG
 4551 CGGACAGGTA TCCGGTAAGC GGCAGGGTCG GAACAGGAGA GCGCACGAGG
 4601 GAGCTTCCAG GGGGAAACGC CTGGTATCTT TATAGTCCTG TCGGGTTTCG
 4651 CCACCTCTGA CTTGAGCGTC GATTTTGTG ATGCTCGTCA GGGGGGCGGA
 4701 GCCTATGGAA AAACGCCAGC AACGCGGCCT TTTACGGTT CCTGGCCTTT
 4751 TGCTGGCCTT TTGCTCACAT GTCTTTTCCT GCGTTATCCC CTGATTCTGT
 4801 GGATAACCGT ATTACCGCCT TTGAGTGAGC TGATACCGCT CGCCGCAGCC

Fig. 30 /5

4851 GAACGACCGA GCGCAGCGAG TCAGTGAGCG AGGAAGCGGA AGAGCGCCTG
 4901 ATGCGGTATT TTCTCCTTAC GCATCTGTGC GGTATTTAC ACCGCATATG
 4951 GTGCACTCTC AGTACAATCT GCTCTGATGC CGCATAGTTA AGCCAGTATA ^{Bst1107I}
 5001 CACTCCGCTA TCGCTACGTG ACTGGGTCAT GGCTGCGCCC CGACACCCGC
 5051 CAACACCCGC TGACGCGCCC TGACGGGCTT GTCTGCTCCC GGCATCCGCT
 5101 TACAGACAAG CTGTGACCGT CTCCGGGAGC TGCATGTGTC AGAGGTTTTC
 5151 ACCGTCATCA CCGAAACGCG CGAGGCAGCT GTGGAATGTG TGTCAGTTAG
 5201 GGTGTGGAAA GTCCCCAGGC TCCCCAGCAG GCAGAAGTAT GCAAAGCATG
 5251 CATCTCAATT AGTCAGCAAC CAGGCTCCCC AGCAGGCAGA AGTATGCAAA
 5301 GCATGCATCT CAATTAGTCA GCAACCATAG TCCCGCCCCT AACTCCGCCC
 5351 ATCCCGCCCC TAACTCCGCC CAGTTCCGCC CATTCTCCGC CCCATGGCTG
 5401 ACTAATTTTT TTTATTTATG CAGAGGCCGA GGCCGCTCG GCCTCTGAGC ^{Sfi I}
 5451 TATTCCAGAA GTAGTGAGGA GGCTTTTTTG GAGGCCTAGG ^{Stu I/Avr II} CTTTGGCAA
 5501 AAGCTAGCTT CACGCTGCCG CAAGCACTCA GGGCGCAAGG GCTGCTAAAG
 5551 GAAGCGGAAC ACGTAGAAAG CCAGTCCGCA GAAACGGTGC TGACCCCGGA
 5601 TGAATGTCAG CTA CTG GGGCT ATCTGGACAA GGGAAAACGC AAGCGCAAAG
 5651 AGAAAGCAGG TAGCTTGAG TGGGCTTACA TGGCGATAGC TAGACTGGGC
 5701 GGTTTTATGG ACAGCAAGCG AACCGGAATT GCCAGCTGGG GCGCCCTCTG
 5751 GTAAGGTGG GAAGCCCTGC AAAGTAACT GGATGGCTTT CTTGCCGCCA
 5801 AGGATCTGAT GGCGCAGGGG ATCAAGATCT GATCAAGAGA CAGGATGAGG ^{Bgl II/Bcl I}
 5851 ATCGTTTCGC ATGATTGAAC AAGATGGATT GCACGCAGGT TCTCCGGCCG
 5901 CTTGGGTGGA GAGGCTATTC GGCTATGACT GGGCACAACA GACAATCGGC
 5951 TGCTCTGATG CCGCCGTGTT CCGGCTGTCA GCGCAGGGGC GCCCGGTTCT
 6001 TTTTGTCAAG ACCGACCTGT CCGGTGCCCT GAATGAACTG CAGGACGAGG
 6051 CAGCGCGGCT ATCGTGGCTG GCCACGACGG GCGTTCCTTG CGCAGCTGTG ^{Msc I}

Fig. 30 /6

6101 CTCGACGTTG TCACTGAAGC GGGAAGGGAC TGGCTGCTAT TGGGCGAAGT
 6151 GCCGGGGCAG GATCTCCTGT CATCTCACCT TGCTCCTGCC GAGAAAGTAT
 6201 CCATCATGGC TGATGCAATG CGGCGGCTGC ATACGCTTGA TCCGGCTACC
 6251 TGCCCATTCG ACCACCAAGC GAAACATCGC ATCGAGCGAG CACGTACTCG
 6301 GATGGAAGCC GGTCTTGTCG ATCAGGATGA TCTGGACGAA GAGCATCAGG
 6351 GGCTCGCGCC AGCCGAACTG TTCGCCAGGC TCAAGGCGCG CATGCCCGAC
 6401 GGCGAGGATC TCGTCGTGAC CCATGGCGAT GCCTGCTTGC CGAATATCAT
 6451 GGTGGAAAAT GGCCGCTTTT CTGGATTCAT CGACTGTGGC CGGCTGGGTG
 6501 TGGCGGACCG CTATCAGGAC ATAGCGTTGG CTACCCGTGA TATTGCTGAA
 6551 GAGCTTGGCG GCGAATGGGC TGACCGCTTC CTCGTGCTTT ACGGTATCGC
 6601 CGCTCCCGAT TCGCAGCGCA TCGCCTTCTA TCGCCTTCTT GACGAGTTCT
 6651 TCTGAGCGGG ACTCTGGGGT TCGAAATGAC CGACCAAGCG ACGCCCAACC
 6701 TGCCATCACG AGATTTGAT TCCACCGCCG CCTTCTATGA AAGGTTGGGC
 6751 TTCGGAATCG TTTTCCGGGA CGCCGGCTGG ATGATCCTCC AGCGCGGGGA
 6801 TCTCATGCTG GAGTTCTTCG CCCACCCCGG GCTCGATCCC CTCGCGAGTT
 6851 GGTTAGCTG CTGCCTGAGG CTGGACGACC TCGCGGAGTT CTACCGGCAG
 6901 TGCAAATCCG TCGGCATCCA GGAAACCAGC AGCGGCTATC CGCGCATCCA
 6951 TGCCCCCGAA CTGCAGGAGT GGGGAGGCAC GATGGCCGCT TTGGTCCCGG
 7001 ATCTTTGTGA AGGAACCTTA CTTCTGTGGT GTGACATAAT TGGACAACT
 7051 ACCTACAGAG ATTTAAAGCT CTAAGGTAAA TATAAAATTT TTAAGTGTAT
 7101 AATGTGTAA ACTACTGATT CTAATTGTTT GTGTATTTTA GATTCCAACC
 7151 TATGGAAGT ATGAATGGGA GCAGTGGTGG AATGCCTTTA ATGAGGAAAA
 7201 CCTGTTTTGC TCAGAAGAAA TGCCATCTAG TGATGATGAG GCTACTGCTG
 7251 ACTCTCAACA TTCTACTCCT CCAAAAAAGA AGAGAAAGGT AGAAGACCCC
 7301 AAGGACTTTC CTTCAGAATT GCTAAGTTTT TTGAGTCATG CTGTGTTTAG

Fig. 30 /7

7351 TAATAGAACT CTTGCTTGCT TTGCTATTTA CACCACAAAG GAAAAAGCTG
7401 CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC CTTTATAAGT
7451 AGGCATAACA GTTATAATCA TAACATACTG TTTTTTCTTA CTCCACACAG
7501 GCATAGAGTG TCTGCTATTA ATAACTATGC TCAAAAATTG TGTACCTTTA
7551 GCTTTTAAAT TTGTAAAGGG GTTAATAAGG AATATTTGAT GTATAGTGCC
7601 TTGACTAGAG ATCATAATCA GCCATACCAC ATTTGTAGAG GTTTTACTTG
7651 CTTTAAAAAA CCTCCACAC CTCCCCCTGA ACCTGAAACA TAAATGAAT
Mun I
7701 GCAATTGTTG TTGTAACTT GTTTATTGCA GCTTATAATG GTTACAAATA
7751 AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTTTTT TCACTGCATT
7801 CTAGTTGTGG TTTGTCCAAA CTCATCAATG TATCTTATCA TGTCTGGATC
7851 TAATAAAAGA TATTTATTTT CATTAGATAT GTGTGTTGGT TTTTGTGTG
7901 CAGTGCCTCT ATCTGGAGGC CAGGTAGGGC TGGCCTTGGG GGAGGGGGAG
7951 GCCAGAATGA CTCCAAGAGC TACAGGAAGG CAGGTCAGAG ACCCCACTGG
8001 ACAAACAGTG GCTGGACTCT GCACCATAAC ACACAATCAA CAGGGGAGTG
8051 AGCTGGAAAT TTGCTAGC

Fig. 31




Fig. 32 /1

	1	Q	V	Q	L	V	Q	S	G	A	10	E	V	K	K	P	G	A	S	V	19	K		
A	CAG	GTG	CAA	CTA	GTG	CAG	TCC	GGC	GCC	GAA	GTG	AAG	AAA	CCC	GGT	GCT	TCC	GTG	AAA					
B		
C		
D		
E		
	20	V	S	C	K	T	S	R	Y	T	F	T	30	CDR1							38	W	V	R
A	GTC	AGC	TGT	AAA	ACT	AGT	AGA	TAC	ACC	TTC	ACT	GAA	TAC	ACC	ATA	CAC	TGG	GTT	AGA					
B
C
D
E	G
							G																	
	39	Q	A	P	G	Q	R	L	E	W	I	G	49	52 A 53							56	G	I	I
A	CAG	GCC	CCT	GGC	CAA	AGG	CTG	GAG	TGG	ATA	GGA	GGT	ATT	AAT	CCT	AAC	AAT	GGT	ATT					
B
C
D
E
	57	CDR2												70	75									
A	P	N	Y	N	Q	K	F	K	G	R	A	T	L	T	V	G	K	S	A					
A	CCT	AAC	TAC	AAC	CAG	AAG	TTC	AAG	GGC	CGG	GCC	ACC	TTG	ACC	GTA	GGC	AAG	TCT	GCC					
B
C	V	.	I	.	.	D	T
D	T	.	A-C	.	.	A	CC
E	V	.	I	.	.	D	T
											T	.	A-C	.	.	A	CC
	76	S	T	A	Y	M	E	L	S	B	C	83	S	E	D	T	A	V	Y	Y				
A	AGC	ACC	GCC	TAC	ATG	GAA	CTG	TCC	AGC	CTG	CGC	TCC	GAG	GAC	ACT	GCA	GTC	TAC	TAC					
B
C
D
E

Fig. 32 /2

	92			CDR3					100	A	B	C	D	I	J	K	101		103
	C	A	R	R	R	I	A	Y	G	Y	D	E	G	H	A	M	D	Y	W
A	TGC	GCC	AGA	AGA	AGA	ATC	GCC	TAT	GGT	TAC	GAC	GAG	GGC	CAT	GCT	ATG	GAC	TAC	TGG
B
C
D
E

	104									113
	G	Q	G	T	L	V	T	V	S	S
A	GGT	CAA	GGA	ACC	CTT	GTC	ACC	GTC	TCC	TCA
B
C
D
E

Fig. 33 /1

1 TTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAAT
 61 GGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTT
 121 ATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCT
 181 TCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCC
 241 CTTTTTTGCGGCATTTTGCCTTCCTGTTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAA
 301 AGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAACTGGATCTCAACAGCGG
 361 TAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGT
 421 TCTGCTATGTGGCGCGGTATTATCCCGTGTGACGCCGGGCAAGAGCAACTCGGTGCGCCG
 481 CATACACTATTCTCAGAATGACTTGTTGAGTACTCACCAGTCACAGAAAAGCATCTTAC
 541 GGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACAATGC
 Pvu I
 601 GGCCAACTTACTTCTGACAACCGATCGGAGGACCGAAGGAGCTAACCCTTTTTTGCACAA
 661 CATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACC
 Fsp I
 721 AAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTGCGCAAACTATT
 781 AACTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGA

Fig. 33 /2

841 TAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTCCGGCTGGCTGGTTTATTGCTGATAA
 901 ATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAA
 961 GCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAA
 1021 TAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGT
 1081 TTAATCATATATACTTTAGATTGATTTAAACTTCATTTTAAATTTAAAGGATCTAGGT
 1141 GAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGTCCACTG
 1201 AGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTCTGCGCGT
 1261 AATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCA
 1321 AGAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATAC
 1381 TGTCTTCTAGTGAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTAC
 1441 ATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCT
 1501 TACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGGGCTGAACGGG
 1561 GGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACA
 1621 GCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGT
 1681 AAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACGCCTGGTA
 1741 TCTTTATAGTCCTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTC
 1801 GTCAGGGGGGCGGAGCCTATGGAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGC
 1861 CTTTTGCTGGCCTTTTGCTCACATGTTCCTTCTGCGTTATCCCCTGATTCTGTGGATAA
 1921 CCGTATTACCGCCTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAG
 1981 CGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCTGATGCGGTATTTCTCCTTACGCATCT
 2041 GTGCGGTATTTACACCGCATATGGTGCCTCTCAGTACAATCTGCTCTGATGCCGCATA
 2101 GTTAAGCCAGTATACACTCCGCTATCGCTACGTGACTGGGTATGGCTGCGCCCCGACAC
 2161 CCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGA
 2221 CAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTGAGAGGTTTACCGTCAATCACCAGAA
 2281 CGCGCGAGGCAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACCTCGCC
 2341 CATCCCCGCCCTAACTCCGCCAGTTCGCCCATTTCTCCGCCCATGGCTGACTAATTTT
 2401 TTTTATTTATGCAGAGGCCGAGGCCCTCGGCCCTGAGCTATTCAGAAAGTAGTGAGG
 2461 AGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTAGCTTACAGCTCAGGGCTGCGATT

BspLU11I

Bst1107 I

Sfi I

Stu I/Avr II

Fig. 33 /2

2521 TCGCGCCAAACTTGACGGCAATCCTAGCGTGAAGGCTGGTAGGATTTTATCCCCGCTGCC
 2581 ATCATGGTTCGACCATTGAACTGCATCGTCGCCGTGTCCCAAATATGGGGATTGGCAAG
 2641 AACGGAGACCTACCCCTGGCCTCCGCTCAGGAACGAGTTCAAGTACTTCCAAAGAATGACC
 2701 ACAACCTCTTCAGTGAAGGTAAACAGAATCTGGTGATTATGGGTAGGAAAACCTGGTTC
 2761 TCCATTCTTGAGAAGAATCGACCTTTAAAGGACAGAATTAATATAGTTCTCAGTAGAGAA
 2821 CTCAAAGAACCACCACGAGGAGCTCATTTTCTTGCCAAAAGTTTGGATGATGCCTTAAGA
 2881 CTTATTGAACAACCGGAATTGGCAAGTAAAGTAGACATGGTTTGGATAGTCGGAGGCAGT
 2941 TCTGTTTACCAGGAAGCCATGAATCAACCAGGCCACCTCAGACTCTTTGTGACAAGGATC
 3001 ATGCAGGAATTTGAAAGTGACACGTTTTTCCCAGAAATTGATTTGGGGAAATATAAACTT
 3061 CTCCCAGAATACCCAGGCGTCCTCTCTGAGGTCCAGGAGGAAAAAGGCATCAAGTATAAG
 3121 TTTGAAGTCTACGAGAAGAAAGACTAACAGGAAGATGCTTTCAAGTTCTCTGCTCCCCTC
 3181 CTAAAGCTATGCATTTTTTATAAGACCATGGGACTTTTGCTGGCTTTAGATCTTTGTGAAG
 3241 GAACCTTACTTCTGTGGTGTGACATAATTGGACAAACTACCTACAGAGATTTAAAGCTCT
 3301 AAGGTAAATATAAAATTTTTAAGTGTATAATGTGTTAACTACTGATTCTAATTGTTTGT
 3361 GTATTTTAGATTCCAACCTATGGAAGTATGAATGGGAGCAGTGGTGAATGCCTTTAAT
 3421 GAGGAAAACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGAC
 3481 TCTCAACATTCTACTCCTCCAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCT
 3541 TCAGAATTGCTAAGTTTTTTGAGTCATGCTGTGTTTAGTAATAGAACTCTTGCTTGCTTT
 3601 GCTATTTACACCACAAAGGAAAAAGCTGCACCTGCTATACAAGAAAATTATGGAAAAATAT
 3661 TCTGTAACCTTTATAAGTAGGCATAACAGTTATAATCATAACATACTGTTTTTCTTACT
 3721 CCACACAGGCATAGAGTGTCTGCTATTAATAACTATGCTCAAAAATTGTGTACCTTTAGC
 3781 TTTTTAATTTGTAAAGGGTTAATAAGGAATATTTGATGTATAGTGCCTTGACTAGAGAT
 3841 CATAATCAGCCATACCACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTCCCACACCT
 3901 CCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTGTTAACTTGTTTATTGCAGC
 3961 TTATAATGGTTACAAATAAAGCAATAGCATCACAATTTACAAATAAAGCATTTTTTTC
 4021 ACTGCATTCTAGTTGTGGTTTGTCCAACTCATCAATGTATCTTATCATGTCTGGATCTA
 4081 ATAAAAGATATTTATTTTCATTAGATATGTGTGTTGGTTTTTTGTGTGCAGTGCCTCTAT
 4141 CTGGAGGCCAGGTAGGGCTGGCCTTGGGGGAGGGGAGGCCAGAATGACTCCAAGAGCTA

Bgl II

BsaB I

Mun I

4201 CAGGAAGGCAGGTCTAGAGACCCCACTGGACAAACAGTGGCTGGACTCTGCACCATAACAC
EcoR I

4261 ACAATCAACAGGGGAGTGAGCTGGAAATTTGCTAGCGAATTCcagcacactggcgccgct
(Spe I)

4321 tACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTT

4381 CCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCC

4441 ATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACG

4501 TCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATAT

4561 GCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCA
SnaB I

4621 GTACATGACCTTATGGGACTTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTAT

4681 TACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTTGACTCAGG

4741 GGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTGGTGGCACCAAAATCA

4801 ACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCG

4861 TGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAG

4921 ACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGCGG

4981 CCGGGAACGGTGCATTGGAACGCGGATTCCCCGTGCCAAGAGTGACGTAAGTACCGCCTA

5041 TAGAGTCTATAGGCCCCACCCCTTGGCTTCTTATGCATGCTATACTGTTTTTGGCTTGGG
Bpu1102I

5101 GTCTATACACCCCGCTTCCTCATGTTATAGGTGATGGTATAGCTTAGCCTATAGGTGTG
Xcm I

5161 GGTATTGACCATTATTGACCCACTCCCTATTGGTGACGATACTTTCCATTACTAATCCA

5221 TAACATGGCTCTTTGCCACAACCTCTCTTTATTGGCTATATGCCAATACACTGTCCTTCAG

5281 AGACTGACACGGACTCTGTATTTTTACAGGATGGGGTCTCATTATTTATTTACAAATTCA

5341 CATATACAACACCACCGTCCCCAGTGCCCGCAGTTTTTTATTAACATAACGTGGGATCTC
BspE I

5401 CACGCGAATCTCGGGTACGTGTTCCGGACATGGGCTCTTCTCCGGTAGCGGCGGAGCTTC

5461 TACATCCGAGCCCTGCTCCCATGCCTCCAGCGACTCATGGTCGCTCGGCAGCTCCTTGCT

5521 CCTAACAGTGAGAGGCCAGACTTAGGCACAGCACGATGCCACCACCACCAGTGTGCCGCA

5581 CAAGGCCGTGGCGGTAGGGTATGTGTCTGAAAATGAGCTCggggagcgggcttgaccgc
(Pvu II)

5641 tgacgcatttggaagacttaaggcagcggcagaagaagatgcaggcagctgaggttgttgt

5701 gttctgataagagtcagaggttaactccggttgcggtgctgttaacggtggagggcagtg

5761 agtctgagcagtactcggttgcgcgcgcgcaccagacataatagctgacagactaa
Mlu I

5821 cagactgttcctttccatgggtcttttctgcagtcaccgtccttgacACGCGTCTCGGGA

Hind III

Fig. 33 /5

6961 TCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA
 L T V L H Q D W L N G K E Y K C K V S N

7021 AAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC
 K A L P A P I E K T I S K A K G Q P R E

7081 CACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGA
 P Q V Y T L P P S R E E M T K N Q V S L

7141 CCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGC
 T C L V K G F Y P S D I A V E W E S N G

7201 AGCCGGAGAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCC
 Q P E N N Y K T T P P V L D S D G S F F

7261 TCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCT
 L Y S K L T V D K S R W Q Q G N V F S C

7321 CCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
 S V M H E A L H N H Y T Q K S L S L S P

Ngm I

7381 GTAAATGAGTGCGACGGCCGGCAAGCCCCGCTCCCCGGGCTCTCGCGGTGCGACGAGGAT
 G K *

7441 GCTTGGCACGTACCCCTGTACATACTTCCCGGGCGCCAGCATGGAAATAAAGCACCGG

7501 ATCTAATAAAAGATATTTATTTTCATTAGATATGTGTGTTGGTTTTTTTGTGTGCAGTGCC

7561 TCTATCTGGAGGCCAGGTAGGGCTGGCCTTGGGGGAGGGGGAGGCCAGAATGACTCCAAG

7621 AGCTACAGGAAGGCAGGTCAGAGACCCCACTGGACAAACAGTGGCTGGACTCTGCACCAT

7681 AACACACAATCAACAGGGGAGTGAGCTGGaaatttgctagcgaattaattc 7731

Fig. 34 A





Fig. 34 B



⇓



⇓



Fig. 35




Fig. 36





Fig. 37





European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 98 10 7925

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	WELT ET AL.: "Antibody targeting in metastatic colon cancer: a phase I study of monoclonal antibody F19 against a cell-surface protein of reactive tumor stromal fibroblasts" JOURNAL OF CLINICAL ONCOLOGY, vol. 12, no. 6, June 1994, pages 1193-1203, XP002088696 * abstract * * page 1193, column 1, line 1 - page 1194, column 2, line 4 * * page 1202, column 2, paragraph 2 * ---	1-65	C12N15/13 C07K16/40 C07K16/46 C12N15/62 C12N15/85 C12N5/10 C07K19/00 A61K47/48 A61K51/10 A61K39/395 G01N33/577 G01N33/574
Y	WO 93 05804 A (SLOAN KETTERING INST CANCER) 1 April 1993 * abstract; claims 1-23 * ---	1-65	
Y	US 5 693 761 A (SCHNEIDER WILLIAM P ET AL) 2 December 1997 * abstract * * examples 3-9 * * column 2, line 36 - column 3, line 59 * --- -/--	1-65	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search MUNICH		Date of completion of the search 21 December 1998	Examiner Muller-Thomalla, K
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 02 (P4/C07)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 98 10 7925

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.CI.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	STUDNICKA G M ET AL: "Human-engineered monoclonal antibodies retain full specific binding activity by preserving non- CDR complementarity-modulating residues." PROTEIN ENGINEERING, (1994 JUN) 7 (6) 805-14. JOURNAL CODE: PR1. ISSN: 0269-2139., XP000447301 ENGLAND: United Kingdom * page 805, column 1, line 1 - page 806, column 2, paragraph 1 * * page 808, column 2, paragraph 2 - page 812, column 1, paragraph 1 * * page 813, column 2, paragraph 1 *	1-65	
Y	WRIGHT A ET AL: "Genetically engineered antibodies: progress and prospects." CRITICAL REVIEWS IN IMMUNOLOGY, (1992) 12 (3-4) 125-68. REF: 252 JOURNAL CODE: AF1. ISSN: 1040-8401., XP000616488 United States * page 139, column 2, paragraph 3 - page 141, column 1, paragraph 3 * * page 157, column 2, paragraph 3 - page 158, column 1, paragraph 1 *	1-65	TECHNICAL FIELDS SEARCHED (Int.CI.6)
A	WO 94 05690 A (SMITHKLINE BEECHAM CORP ;US ARMY (US); GROSS MITCHELL STUART (US);) 17 March 1994 * claim 5: figure 3 *	14-17	



European Patent
Office

**INCOMPLETE SEARCH
SHEET C**

Application Number
EP 98 10 7925

Although claims 50-52,54,55,57,61,62,65 are directed to a method of treatment of the human/animal body and/or a diagnostic method practised on the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ ~~FADED TEXT OR DRAWING~~
- ☒ ~~BLURRED OR ILLEGIBLE TEXT OR DRAWING~~
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ ~~LINES OR MARKS ON ORIGINAL DOCUMENT~~
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.